



**HAL**  
open science

# Hamiltonian Evolution of Monokinetic Measures with Rough Momentum Profile

Claude Bardos, François Golse, Peter Markowich, Thierry Paul

► **To cite this version:**

Claude Bardos, François Golse, Peter Markowich, Thierry Paul. Hamiltonian Evolution of Monokinetic Measures with Rough Momentum Profile. 2012. hal-00706180v1

**HAL Id: hal-00706180**

**<https://hal-polytechnique.archives-ouvertes.fr/hal-00706180v1>**

Preprint submitted on 9 Jun 2012 (v1), last revised 26 Apr 2013 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# HAMILTONIAN EVOLUTION OF MONOKINETIC MEASURES WITH ROUGH MOMENTUM PROFILE

CLAUDE BARDOS, FRANÇOIS GOLSE, PETER MARKOWICH, AND THIERRY PAUL

ABSTRACT. Consider in the phase space  $\mathbf{R}_x^N \times \mathbf{R}_\xi^N$  a probability density carried by the graph of a vector field  $U^{in}$  on  $\mathbf{R}^N$ , i.e. a Radon measure of the form  $\mu^{in} = \rho^{in}(x)\delta(\xi - U^{in}(x))$ . Let  $\Phi_t$  be a Hamiltonian flow on  $\mathbf{R}_x^N \times \mathbf{R}_\xi^N$ . In this paper, we study the structure of the transported measure  $\mu(t) := \Phi_t\#\mu^{in}$  and of its integral in the  $\xi$  variable denoted  $\rho(t)$ . In particular, we (a) provide estimates on the number of folds in  $\text{supp}(\mu(t)) = \Phi_t(\text{graph of } U^{in})$ , (b) establish a decomposition of  $\mu(t)$  into a “regular” component whose integral in the  $\xi$  variable is absolutely continuous with respect to the Lebesgue measure  $\mathcal{L}^N$ , (c) discuss the possibility of atoms for the measure  $\rho(t)$  and (d) construct an example in which  $\rho(t)$  is singular with respect to  $\mathcal{L}^N$  and diffuse. We conclude our study by explaining how our results can be applied to the classical limit of the Schrödinger equation by using the formalism of Wigner measures. Our results hold for initial momentum profiles  $U^{in}$  less regular than  $C^1$ , for which the usual notion of caustic is not relevant. The proofs of these results is based on the area formula of geometric measure theory.

## 1. INTRODUCTION

The subject of this article is the propagation of a certain class of positive Radon measures by Hamiltonian flows.

Let  $H \equiv H(x, \xi)$  be a Hamiltonian of class  $C^2$  on  $T^*\mathbf{R}^N = \mathbf{R}_x^N \times \mathbf{R}_\xi^N$ . Assume that the system of Hamilton’s equations

$$(1) \quad \begin{cases} \dot{X}_t = \nabla_\xi H(X_t, \Xi_t), & X_0(x, \xi) = x, \\ \dot{\Xi}_t = -\nabla_x H(X_t, \Xi_t), & \Xi_0(x, \xi) = \xi, \end{cases}$$

generates a global flow

$$(2) \quad \Phi_t(x, \xi) = (X_t(x, \xi), \Xi_t(x, \xi)).$$

Let  $\mu^{in}$  be a monokinetic measure on  $\mathbf{R}_x^N \times \mathbf{R}_\xi^N$  i.e. a Radon measure of the form

$$(3) \quad \mu^{in}(x, \xi) = \rho^{in}(x)\delta_{U^{in}(x)}(\xi)$$

where  $U^{in}$  is a vector field on  $\mathbf{R}^N$  and  $\rho^{in} \in L^1(\mathbf{R}^N)$ . Define

$$(4) \quad \mu(t) = \Phi_t\#\mu^{in}$$

---

1991 *Mathematics Subject Classification.* 81Q20, 81S30, 35Q40, 35L03, 28A75.

*Key words and phrases.* Wigner measure, Liouville equation, Schrödinger equation, WKB method, Caustic, Area formula, Coarea formula .

the push-forward of  $\mu^{in}$  under  $\Phi_t$ . Equivalently,  $\mu(t) \in C_b(\mathbf{R}_+; w - \mathcal{M}(\mathbf{R}^N \times \mathbf{R}^N))$  is the unique weak solution of the Liouville equation

$$(5) \quad \begin{cases} \partial_t \mu + \{H, \mu\} = 0, \\ \mu|_{t=0} = \mu^{in}, \end{cases}$$

where  $\{\cdot, \cdot\}$  designates the Poisson bracket

$$\{f, g\} = \nabla_\xi f \cdot \nabla_x g - \nabla_x f \cdot \nabla_\xi g.$$

Our purpose is to study the structure of  $\mu(t)$  and to deduce from it some information on its support

$$\text{supp}(\mu(t)) \subset \Lambda_t := \Phi_t(\{(y, U^{in}(y)) \mid y \in \mathbf{R}^N\}).$$

While  $\Lambda_t$  is the image under  $\Phi_t$  of the graph of  $U^{in}$ , it is not in general the graph of a vector field on  $\mathbf{R}^N$  for all values of  $t$ .

When  $U^{in} = \nabla S^{in}$  is a smooth gradient field on  $\mathbf{R}^N$ , then  $\Lambda_t$  is the union of graphs of  $x \mapsto \nabla_x S_j(t, x)$ , where  $S_j$  is a solution of the Hamilton-Jacobi equation

$$(6) \quad \partial_t S(t, x) + H(x, \nabla_x S(t, x)) = 0$$

defined on some open set of  $\mathbf{R}_t \times \mathbf{R}^N$ . The graphs of  $\nabla_x S_j$  are glued along submanifolds of  $\Lambda_t$  where the restriction of the canonical projection

$$\Pi : T^*\mathbf{R}^N = \mathbf{R}_x^N \times \mathbf{R}_\xi^N \ni (x, \xi) \mapsto x \in \mathbf{R}^N$$

is not smooth: see §8 in [4] and §46-47 in [5]. A natural question is to compute, or at least estimate, the number of solutions  $S_j$  of the Hamilton-Jacobi equation needed to obtain  $\Lambda_t$ .

Equivalently, the restriction to  $\Lambda_t$  of the canonical projection  $\Pi$  is in general not one-to-one for all  $t$ . The question above reduces to estimating the number  $\mathcal{N}(t, x)$  of points in  $\Lambda_t$  whose image under  $\Pi$  is  $x$ . Thus, the function  $\mathcal{N}$  describes the number of folds in  $\Lambda_t$  induced by the Hamiltonian dynamics  $\Phi_t$  on the graph of  $U^{in}$  — even when  $U^{in}$  is not a gradient field.

Another natural question is to study the structure of the push-forward  $\rho(t)$  of  $\mu(t)$  under the canonical projection  $\Pi$  — equivalently, of the first marginal of the measure  $\mu(t)$  in the product space  $\mathbf{R}_x^N \times \mathbf{R}_\xi^N$ . As we shall see, both questions are intimately related.

The mathematical problem described above appears in a great variety of contexts. It appeared first in the theory of geometric optics, in the works of Fermat and Huygens: see for instance chapter VII in [14], chapter III in [10] and chapter 12 §2 in [12]. It appears in the classical limit of quantum mechanics: see for instance chapter VII in [15] or [22] — we shall give more details on this case below.

In both examples above, the fact that  $U^{in}$  is a gradient field is important. There are however other types of physical models leading to the same mathematical problem even when  $U^{in}$  is not a gradient field. Indeed, the Maxwell distribution with density  $\rho$ , bulk velocity  $U$  and temperature  $\theta$ , i.e.

$$\frac{\rho(x)}{(2\pi\theta)^{N/2}} e^{-|\xi - U(x)|^2/2\theta}$$

converges weakly to the monokinetic measure  $\rho(x)\delta_{U(x)}(\xi)$  as  $\theta \rightarrow 0^+$ . The propagation of this class of measures by the flow  $\Phi_t$  generated by the free Hamiltonian  $H(x, \xi) := \frac{1}{2}|\xi|^2$ , i.e.  $\Phi_t(x, \xi) = (x + t\xi, \xi)$  can be viewed as the kinetic theory of

pressureless gases, and appears for instance in a cosmological model due to Zel'dovich [24, 8, 7, 6]. When  $U^{in}$  is a gradient field, the Liouville equation (5) can therefore be viewed as a kinetic formulation of the Hamilton-Jacobi equation (6).

While the classical mathematical theory of geometric optics or of the semi-classical limit of quantum mechanics is centered on the geometry of  $\Lambda_t$  in the case where both the Hamiltonian  $H$  and  $U^{in} = \nabla S^{in}$  are smooth, our approach of the mathematical problem stated above is centered on the propagation of the monokinetic measure  $\rho(x)\delta_{U^{in}(x)}(\xi)$  by the Hamiltonian flow. Besides, our analysis on the propagation problem is focussed on mathematical methods and results in which the initial momentum profile  $U^{in}$  is not required to be everywhere differentiable — so that  $\Lambda_t$  is not even a  $C^1$ -manifold. Likewise, the Hamiltonian nature of the dynamics is of limited importance in our analysis, and there is no need for the momentum profile  $U^{in}$  to be a gradient — or, equivalently, for  $\Lambda_t$  to be a Lagrangian submanifold of  $T^*\mathbf{R}^N$ .

However, we assume that the vector field generating the dynamics is at least of class  $C^1$  — or equivalently that the Hamiltonian  $H$  is at least of class  $C^2$  on  $\mathbf{R}_x^N \times \mathbf{R}_\xi^N$  — so that the existence, uniqueness and regularity of the flow  $\Phi_t$  results from the classical Cauchy-Lipschitz theory. The classical limit of quantum dynamics with rough potentials, for which the existence and uniqueness of the classical Hamiltonian flow does not follow from the Cauchy-Lipschitz theory, has been recently studied in [1]. Our viewpoint in the present paper is different and in some sense complementary: we focus our attention to the special class of monokinetic measures and to their propagation by smooth Hamiltonian flows, but obtain detailed information on the structure of the propagated measure  $\mu(t)$  even for rough initial momentum profiles  $U^{in}$ .

The outline of the paper is as follows. In section 2 are gathered our assumptions on the Hamiltonian  $H$  with some elementary estimates on the flow  $\Phi_t$  that are crucial in the sequel. Section 3 is focussed on the problem of estimating the number of folds in the support  $\Lambda_t$  of the propagated measure  $\mu(t)$ , while the structure of  $\mu(t)$  itself is studied in section 5. Section 4 gathers together a few examples showing that the results in section 3 are sharp. In section 6, we study different exceptional sets that appear naturally in connection with the structure of the projected measure  $\rho(t) := \Pi\#\mu(t)$ , and explain how these sets are related to the traditional notion of “caustic” — introduced by Tschirnhaus in the 17th century in the context of geometric optics). Section 7 discusses various applications of the theory presented in sections 3-5, with an emphasis on the classical limit of quantum mechanics.

## 2. ON THE HAMILTONIAN FLOW

Let  $H \equiv H(x, \xi) \in \mathbf{R}$  be a  $C^2$  function on  $\mathbf{R}^N \times \mathbf{R}^N$  satisfying the following assumptions: there exists  $\kappa > 0$ , and a function  $h \in C(\mathbf{R}; \mathbf{R}_+)$  that is sublinear at infinity, i.e.

$$\frac{h(r)}{r} \rightarrow 0 \quad \text{as } r \rightarrow +\infty$$

such that

$$(7) \quad \begin{aligned} |\nabla_\xi H(x, \xi)| &\leq \kappa(1 + |\xi|) \\ |\nabla_x H(x, \xi)| &\leq h(|x|) + \kappa|\xi| \\ |\nabla^2 H(x, \xi)| &\leq \kappa \end{aligned}$$

for all  $(x, \xi) \in \mathbf{R}^N \times \mathbf{R}^N$ .

**Lemma 2.1.** *Any Hamiltonian  $H \in C^2(\mathbf{R}^N \times \mathbf{R}^N)$  satisfying (7) generates a global Hamiltonian flow  $\Phi_t$  on  $\mathbf{R}^N \times \mathbf{R}^N$ . The map*

$$\mathbf{R} \times \mathbf{R}^N \times \mathbf{R}^N \ni (t, x, \xi) \mapsto \Phi_t(x, \xi) \in \mathbf{R}^N \times \mathbf{R}^N$$

is of class  $C^1$ . Moreover, for each  $\eta > 0$ , there exists  $C_\eta > 0$  such that

$$\sup_{|t| \leq T} |X_t(x, \xi) - x| \leq C_\eta(1 + |\xi|) + \eta|x|$$

for each  $x, \xi \in \mathbf{R}^N$ , and

$$|D\Phi_t(x, \xi) - \text{Id}_{\mathbf{R}^N \times \mathbf{R}^N}| \leq e^{\kappa|t|} - 1$$

for all  $t \in \mathbf{R}$ .

*Proof.* By (7), one has the following a priori estimates, with the notation (2).

First

$$|X_t(x, \xi) - x| \leq \int_0^t |\nabla_\xi H(\Phi_s(x, \xi))| ds \leq \kappa t + \kappa \int_0^t |\Xi_s(x, \xi)| ds$$

and

$$\begin{aligned} |\Xi_t(x, \xi)| &\leq |\xi| + \int_0^t |\nabla_x H(\Phi_s(x, \xi))| ds \\ &\leq |\xi| + \kappa \int_0^t |\Xi_s(x, \xi)| ds + \int_0^t h(X_s(x, \xi)) ds. \end{aligned}$$

By Gronwall's inequality, for all  $0 \leq s \leq t$

$$(8) \quad |\Xi_s(x, \xi)| \leq \left( |\xi| + \int_0^t h(X_\tau(x, \xi)) d\tau \right) e^{\kappa s},$$

so that

$$\begin{aligned} |X_t(x, \xi) - x| &\leq \kappa t + \kappa \int_0^t e^{\kappa s} ds \left( |\xi| + \int_0^t h(X_\tau(x, \xi)) d\tau \right) \\ &\leq \kappa t + e^{\kappa t} \left( |\xi| + \int_0^t h(X_\tau(x, \xi)) d\tau \right). \end{aligned}$$

Since  $h$  is sublinear at infinity, we have, for every  $R > 0$

$$(9) \quad h(r) \leq \mathbf{1}_{[0, R]}(r) \sup_{0 \leq r \leq R} h(r) + \mathbf{1}_{(R, +\infty)}(r) r \sup_{r > R} \frac{h(r)}{r} \leq M_R + r m_R,$$

where

$$M_R = \sup_{0 \leq r \leq R} h(r) \quad \text{and} \quad m_R = \sup_{r > R} \frac{h(r)}{r},$$

so that

$$m_R \rightarrow 0 \quad \text{as } R \rightarrow +\infty.$$

Therefore

$$\begin{aligned} |X_t(x, \xi) - x| &\leq (\kappa + M_R e^{\kappa t}) t + e^{\kappa t} |\xi| + m_R e^{\kappa t} \int_0^t |X_s(x, \xi)| ds \\ &\leq (\kappa + M_R e^{\kappa t}) t + e^{\kappa t} |\xi| + m_R e^{\kappa t} |x| + m_R e^{\kappa t} \int_0^t |X_s(x, \xi) - x| ds. \end{aligned}$$

By Gronwall's inequality,

$$(10) \quad \begin{aligned} |X_t(x, \xi) - x| &\leq ((\kappa + M_R e^{\kappa t})t + e^{\kappa t}|\xi| + m_R e^{\kappa t}|x|)e^{tm_R e^{\kappa t}} \\ &\leq \kappa t e^{tm_R e^{\kappa t}} + M_R t e^{t(\kappa + m_R e^{\kappa t})} + |\xi| e^{t(\kappa + m_R e^{\kappa t})} + m_R |x| e^{t(\kappa + m_R e^{\kappa t})}. \end{aligned}$$

The same estimates hold for  $-T \leq t \leq 0$  after substituting  $|t|$  to  $t$ .

In view of (10)-(8), for each  $(x, \xi) \in \mathbf{R}^N \times \mathbf{R}^N$ , the trajectory  $(x, \xi) \mapsto \Phi_t(x, \xi)$  cannot escape to infinity in finite time, and is therefore globally defined.

Besides, since  $m_R \rightarrow 0$  as  $R \rightarrow +\infty$ , the estimate (10) obviously implies the first inequality in the lemma with

$$\eta := m_R e^{T(\kappa + m_R e^{\kappa T})} \quad \text{and} \quad C_\eta := (1 + \kappa T + M_R T) e^{T(\kappa + m_R e^{\kappa T})}.$$

Since  $H \in C^2(\mathbf{R}^N \times \mathbf{R}^N)$ , the map  $(t, x, \xi) \mapsto \Phi_t(x, \xi)$  is of class  $C^1$  on its domain of definition  $\mathbf{R} \times \mathbf{R}^N \times \mathbf{R}^N$ . Differentiating the Hamilton equations (1) with respect to the initial condition, one finds that

$$\begin{cases} \dot{D}X_t = +\nabla_{x, \xi}^2 H(\Phi_t) \cdot DX_t + \nabla_{\xi, \xi}^2 H(\Phi_t) \cdot D\xi_t, \\ \dot{D}\xi_t = -\nabla_{x, x}^2 H(\Phi_t) \cdot DX_t - \nabla_{x, \xi}^2 H(\Phi_t) \cdot D\xi_t, \end{cases}$$

so that

$$|D\Phi_t - \text{Id}_{\mathbf{R}^N \times \mathbf{R}^N}| \leq \kappa \int_0^{|t|} |D\Phi_s| ds.$$

The second inequality in the lemma follows from Gronwall's inequality.  $\square$

### 3. ON THE NUMBER OF FOLDS IN $\Lambda_t$

Let  $U^{in} \in C(\mathbf{R}^N; \mathbf{R}^N)$  satisfy the condition

$$(11) \quad \frac{|U^{in}(y)|}{|y|} \rightarrow 0 \quad \text{as } |y| \rightarrow 0.$$

The present section is focussed on the structure of the set  $\Lambda_t$  in the introduction, defined as

$$(12) \quad \Lambda_t := \Phi_t(\{(y, U^{in}(y)) \mid y \in \mathbf{R}^N\}), \quad t \in \mathbf{R}.$$

Consider the map

$$(13) \quad F_t : \mathbf{R}^N \ni y \mapsto F_t(y) = X_t(y, U^{in}(y)) \in \mathbf{R}^N.$$

Whenever  $U^{in}$  is differentiable at  $y$ , the map  $F_t$  is also differentiable at  $y$  by the chain rule; for any such  $y$ , define absolute value of the Jacobian determinant

$$(14) \quad J_t(y) = |\det(DF_t(y))|.$$

We also introduce the set

$$(15) \quad \begin{aligned} C &:= \{(t, x) \in \mathbf{R} \times \mathbf{R}^N \mid F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\}) \neq \emptyset\}, \\ C_t &:= \{x \in \mathbf{R}^N \mid (t, x) \in C\}. \end{aligned}$$

For want of a better terminology and by analogy with geometric optics,  $C$  will be referred to as the ‘‘caustic’’ set.

**Proposition 3.1.** *Under the conditions above, for each  $t \in \mathbf{R}$ , the map  $F_t$  is proper and onto. Moreover*

$$(16) \quad \sup_{|t| \leq T} \frac{|F_t(y) - y|}{|y|} \rightarrow 0 \quad \text{as } |y| \rightarrow \infty.$$

The proof of this proposition and the next one will use the following topological argument.

**Lemma 3.2.** *Let  $g : \mathbf{R}^N \rightarrow \mathbf{R}^N$  be a continuous map satisfying the following condition: for some  $R > 0$*

$$(g(x)|x) > 0 \quad \text{for all } x \in \mathbf{R}^N \text{ such that } |x| = R.$$

*Then*

- a) *there exists  $x \in \mathbf{R}^N$  such that  $|x| \leq R$  and  $g(x) = 0$ ;*
- b) *if  $g$  is of class  $C^1$  on  $\mathbf{R}^N$  and  $0$  is a regular value of  $g$ , then  $g^{-1}(\{0\}) \cap B(0, R)$  is finite and  $\#(g^{-1}(\{0\}) \cap B(0, R))$  is odd.*

*Proof.* Consider the homotopy  $G \in C([0, 1] \times \mathbf{R}^N; \mathbf{R}^N)$  defined by

$$G(t, x) = tx + (1 - t)g(x).$$

One has

$$G(t, x) \neq 0 \quad \text{whenever } t \in [0, 1] \text{ and } |x| = R.$$

Indeed,  $G(1, x) = x \neq 0$  if  $|x| = R > 0$ ; besides, if  $t \in [0, 1[$  and  $G(t, x) = 0$ , one has

$$g(x) = -\frac{t}{1-t}x \quad \text{so that } (g(x)|x) = -\frac{t}{1-t}|x|^2 = -\frac{t}{1-t}R^2 < 0$$

for all  $x \in \mathbf{R}^N$  such that  $|x| = R$ , which contradicts our assumption.

By the homotopy invariance of the degree (see Properties 7, 8 and Theorem 12.7 in chapter 12, §A of [21])

$$d(g, B(0, R), 0) = d(I, B(0, R), 0) = 1.$$

This implies a). Moreover, if  $g$  is of class  $C^1$  on  $\mathbf{R}^N$  and  $0$  is a regular value of  $g$ , all the elements of  $\overline{g^{-1}(\{0\})}$  are isolated points by the implicit function theorem, so  $g^{-1}(\{0\}) \cap \overline{B(0, R)}$  is finite. Besides (see Property 2 in chapter 12, §A of [21])

$$d(g, B(0, R), 0) = \sum_{x \in g^{-1}(\{0\}) \cap B(0, R)} \text{sign}(\det Dg(x)) = 1.$$

Therefore, there exists an integer  $m \in \mathbf{N}$  such that

$$\begin{aligned} \#\{x \in B(0, R) \mid g(x) = 0 \text{ and } \det(Dg(x)) > 0\} &= m + 1 \\ \#\{x \in B(0, R) \mid g(x) = 0 \text{ and } \det(Dg(x)) < 0\} &= m \end{aligned}$$

so that  $\#(g^{-1}(\{0\}) \cap B(0, R)) = 2m + 1$ , which proves b).  $\square$

*Proof of Proposition 3.1.* By the first inequality in Lemma 2.1 and the condition (11 on  $U^{in}$ , for each  $\eta > 0$ , one has

$$\overline{\lim}_{|y| \rightarrow +\infty} \sup_{|t| \leq T} \frac{|F_t(y) - y|}{|y|} \leq \eta,$$

so that (16) holds.

Because of (16), the continuous map  $F_t$  satisfies

$$(17) \quad (F_t(y) - x|y) = |y|^2 + o(|y|^2) \quad \text{as } |y| \rightarrow +\infty$$

so that  $F_t$  is onto by applying Lemma 3.2 to the map  $g : y \mapsto F_t(y) - x$ . On the other hand

$$|F_t(y)| \rightarrow +\infty \quad \text{as } |y| \rightarrow +\infty$$

so that  $F_t$  is proper.  $\square$

By Proposition 3.1, we know that, for each  $(t, x) \in \mathbf{R} \times \mathbf{R}^N$ , the equation  $F_t(y) = x$  has at least one solution  $y \in \mathbf{R}^N$  when  $U^{in}$  is a continuous vector field sublinear at infinity, i.e. satisfying (11). In the next proposition, we study the number  $\mathcal{N}(t, x)$  of solutions of this equation in the case where  $U^{in}$  is of class  $C^1$  at least. Equivalently,  $\mathcal{N}(t, x)$  is the number of intersections of the manifold  $\Lambda_t$  with  $T_x^* \mathbf{R}^N \simeq \{x\} \times \mathbf{R}^N$ . Therefore, the integer-valued function  $\mathcal{N}$  measures the number of folds in the manifold  $\Lambda_t$  resulting from the interaction of the Hamiltonian flow  $\Phi_t$  with the initial profile  $U^{in}$ .

**Proposition 3.3.** *[Smooth case] Assume that (11) holds for  $U^{in} \in C^1(\mathbf{R}^N, \mathbf{R}^N)$  and that*

$$|DU^{in}(y)| = O(1) \quad \text{as } |y| \rightarrow +\infty.$$

- a) For each  $t \in \mathbf{R}$ , one has  $\mathcal{L}^N(C_t) = 0$ .
- b) The set  $C$  is closed in  $\mathbf{R} \times \mathbf{R}^N$ .
- c) For each  $(t, x) \in \mathbf{R} \times \mathbf{R}^N \setminus C$ , the set  $F_t^{-1}(\{x\})$  is finite, and henceforth denoted by

$$\{y_j(t, x), j = 1, \dots, \mathcal{N}(t, x)\}.$$

The integer  $\mathcal{N}$  is a constant function of  $(t, x)$  in each connected component of  $\mathbf{R} \times \mathbf{R}^N \setminus C$  and, for each  $j \geq 1$ , the map  $y_j$  is of class  $C^1$  on each connected component of  $\mathbf{R} \times \mathbf{R}^N \setminus C$  where  $\mathcal{N} \geq j$ .

- d) There exists  $a < 0 < b$  such that  $C \cap ((a, b) \times \mathbf{R}^N) = \emptyset$  and  $\mathcal{N} = 1$  on  $(a, b) \times \mathbf{R}^N$ .
- e)  $\mathcal{N}(t, x)$  is odd for each  $(t, x) \in \mathbf{R} \times \mathbf{R}^N \setminus C$ .

*Proof.* If  $U^{in} \in C^1(\mathbf{R}^N)$ , the map  $F_t$  is of class  $C^1$  from  $\mathbf{R}^N$  to  $\mathbf{R}^N$ , being the composition of  $C^1$  maps. Since  $C_t$  is the set of critical values of  $F_t$ , one has  $\mathcal{L}^N(C_t) = 0$  by Sard's Theorem (in the equal dimension case), which proves a).

Pick any sequence  $(t_n, x_n) \in C$  such that  $(t_n, x_n) \rightarrow (t, x)$  in  $\mathbf{R} \times \mathbf{R}^N$  as  $n \rightarrow \infty$ . By definition of  $C$ , there exists  $y_n$  such that  $F_{t_n}(y_n) = x_n$ . Since the sequences  $t_n$  and  $x_n$  converge, they are both bounded. Let  $T = \sup_n |t_n|$ ; assume that the sequence  $y_n$  is unbounded; if so there exists a subsequence  $y_{n_l}$  such that  $|y_{n_l}| \rightarrow \infty$ . By (16)

$$|x_{n_l} - y_{n_l}| = |F_{t_{n_l}}(y_{n_l}) - y_{n_l}| \leq \sup_{|t| \leq T} |F_t(y_{n_l}) - y_{n_l}| = o(|y_{n_l}|)$$

so that  $|x_{n_l}| \sim |y_{n_l}|$  as  $n_l \rightarrow +\infty$ . Since this contradicts the fact that the sequence  $x_n$  is bounded, we conclude that the sequence  $y_n$  is bounded. Therefore, there exists a convergent subsequence  $y_{n_k}$  of  $y_n$ ; call  $y$  its limit as  $n_k \rightarrow +\infty$ . Passing to the limit in both relations

$$F_{t_{n_k}}(y_{n_k}) = x_{n_k} \quad \text{and} \quad J_{t_{n_k}}(y_{n_k}) = |\det(\nabla F_{t_{n_k}}(y_{n_k}))| = 0,$$

we conclude that  $(t, x, y)$  satisfies

$$F_t(y) = x \quad \text{and} \quad J_t(y) = 0$$

and therefore that  $(t, x) \in C$ , which proves b).



For  $(t, x) \in \mathbf{R} \times \mathbf{R}^N \setminus C$ , all the solutions of the equation  $F_t(y) - x = 0$  are isolated by the implicit function theorem. The set of all such points,  $F_t^{-1}(\{x\})$ , is therefore compact since  $F_t$  is proper. By the implicit function theorem, the integer  $\mathcal{N}$  is a locally constant function of  $(t, x) \in \mathbf{R} \times \mathbf{R}^N \setminus C$ , and is therefore constant on each connected component of  $\mathbf{R} \times \mathbf{R}^N \setminus C$ . Let  $j \in \mathbf{N}^*$ , and let  $\Omega$  be a connected component of  $\mathbf{R} \times \mathbf{R}^N \setminus C$ ; by the implicit function theorem  $y_j \in C^1(\Omega)$ . This proves c).

Assume  $\inf\{t > 0 \mid C_t \neq \emptyset\} = 0$ . Then, there exists  $(t_n, x_n, y_n)$  such that

$$t_n \rightarrow 0^+, \quad F_{t_n}(y_n) = x_n, \quad \text{and } J_{t_n}(y_n) = 0.$$

Assume that a subsequence  $y_{n_k}$  of the sequence  $y_n$  is bounded. Up to extraction of a subsequence, one can assume that  $y_{n_k} \rightarrow y$ , so that  $0 = J_{t_{n_k}}(y_{n_k}) \rightarrow J_0(y)$ . But since  $F_t = \text{Id}_{\mathbf{R}^N}$ , one has  $J_0(y) = 1$ . Therefore  $|y_n| \rightarrow +\infty$ . By the second inequality in Lemma 2.1

$$\begin{aligned} |D_x X_{t_n}(y_n, U^{in}(y_n)) - \text{Id}_{\mathbf{R}^N}| &\leq e^{\kappa|t_n|} - 1, \\ |D_\xi X_{t_n}(y_n, U^{in}(y_n)) \cdot DU^{in}(y_n)| &= O\left(e^{\kappa|t_n|} - 1\right), \end{aligned}$$

so that

$$\begin{aligned} 0 = J_{t_n}(y_n) &= |\det(D_x X_{t_n}(y_n, U^{in}(y_n)) + D_\xi X_{t_n}(y_n, U^{in}(y_n)) \cdot DU^{in}(y_n))| \\ &\rightarrow |\det(\text{Id}_{\mathbf{R}^N})| = 1 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

Thus the assumption  $t_n \rightarrow 0$  leads to a contradiction. Therefore,

$$\inf\{t > 0 \mid C_t \neq \emptyset\} = b > 0.$$

By the same token,

$$\sup\{t < 0 \mid C_t \neq \emptyset\} = a < 0.$$

Thus  $(a, b) \times \mathbf{R}^N$  is contained in the connected component of  $\{0\} \times \mathbf{R}^N$  in  $\mathbf{R} \times \mathbf{R}^N \setminus C$ . Since  $F_0 = \text{Id}_{\mathbf{R}^N}$ , one has  $\mathcal{N}(0, x) = 1$  for all  $x \in \mathbf{R}^N$ , and since  $\mathcal{N}$  is constant on each connected component of  $\mathbf{R} \times \mathbf{R}^N \setminus C$ , one concludes  $\mathcal{N} = 1$  on  $(a, b) \times \mathbf{R}^N$ , which proves d).

If  $(t, x) \in \mathbf{R} \times \mathbf{R}^N \setminus C$ , the point  $x$  is a regular value of  $F_t$ . Since  $F_t$  is proper,  $F_t^{-1}(\{x\})$  is compact, and therefore bounded. Pick  $R > 0$  such that

$$F_t^{-1}(\{x\}) \subset B(0, R).$$

By (17) and Lemma 3.2 b) applied to the map  $g : y \mapsto F_t(y) - x$

$$\#F_t^{-1}(\{x\}) = \#(F_t^{-1}(\{x\}) \cap B(0, R)) \text{ is odd,}$$

which proves e). □

When  $U^{in}$  is not of class  $C^1$ , the arguments used to prove Proposition 3.3 are no longer valid. However one can still obtain some information on the number  $\mathcal{N}(t, x)$  of solutions  $y$  of the equation  $F_t(y) = x$  by a completely different method, involving the area — or co-area formula.

Assume that  $U^{in} \in C(\mathbf{R}^N; \mathbf{R}^N)$  satisfies (11) and that its gradient (in the sense of distributions)  $DU^{in}$  satisfies the condition

$$(18) \quad \partial_t U_k^{in}|_\Omega \in L^{N,1}(\Omega) \text{ for each bounded open } \Omega \subset \mathbf{R}^N,$$

for all  $k, l = 1, \dots, N$ . We recall that a measurable function  $f : \Omega \rightarrow \mathbf{R}$  belongs to the Lorentz space  $L^{N,1}(\Omega)$  if

$$\int_0^\infty (\mathcal{L}^N(\{x \in \Omega \mid |f(x)| \geq \lambda\}))^{1/N} d\lambda < \infty.$$

By Theorem B in [13], the vector field  $U^{in}$  is differentiable a.e. on  $\mathbf{R}^N$ . Let  $E$  be the  $\mathcal{L}^N$ -negligible set defined as

$$(19) \quad E := \{y \in \mathbf{R}^N \mid U^{in} \text{ is not differentiable at } y\}.$$

By the chain rule, the absolute value of the Jacobian determinant

$$(20) \quad J_t(y) = |\det(D_x X_t(y, U^{in}(y)) + D_\xi X_t(y, U^{in}(y)) D U^{in}(y))|$$

is defined for all  $(t, y) \in \mathbf{R} \times (\mathbf{R}^N \setminus E)$ .

Henceforth, the notation  $J_t^{-1}(\{0\})$  designates the set

$$(21) \quad J_t^{-1}(\{0\}) := \{y \in \mathbf{R}^N \setminus E \mid J_t(y) = 0\},$$

and we shall also consider the set

$$(22) \quad Z_t := J_t^{-1}(\{0\}) \cup E.$$

**Theorem 3.4.** [*Rough case*] Assume that  $U^{in} \in C(\mathbf{R}^N; \mathbf{R}^N)$  satisfies (11) and (18).

a) for all  $t \in \mathbf{R}$ , the function  $J_t \in L^1_{loc}(\mathbf{R}^N)$ ;

b) for all  $t \in \mathbf{R}$ , there are finitely many solutions  $y$  of the equation

$$(23) \quad F_t(y) = x$$

for a.e.  $x \in \mathbf{R}^N$  — in other words,  $\mathcal{N}(t, x)$  is finite for a.e.  $x \in \mathbf{R}^N$ , for all  $t \in \mathbf{R}$ ;

c) for each bounded  $B \subset \mathbf{R}^N$ , denote by  $\mathcal{N}_B(t, x)$  the number of solutions  $y \in B$  of (23); then, whenever  $B$  is bounded, for each  $t \in \mathbf{R}$  and each  $n \in \mathbf{N}$ , one has

$$\begin{aligned} \mathcal{L}^N(\{x \in \mathbf{R}^N \mid \mathcal{N}_B(t, x) \geq n\}) &\leq \frac{1}{n} \int_B J_t(y) dy \\ &\leq \frac{1}{n} e^{N\kappa|t|} \|1 + (1 - e^{-\kappa|t|})\| \|D U^{in}\|_{L^N(B)}^N; \end{aligned}$$

d) let  $t \in \mathbf{R}$ ; then for a.e.  $x \in \mathbf{R}^N$ , all solutions  $y$  of (23) satisfy

$$y \in \mathbf{R}^N \setminus E \text{ and } J_t(y) > 0;$$

e) for all  $T > 0$

$$\mathcal{H}^1(\{(t, y) \in [-T, T] \times \mathbf{R}^N \mid F_t(y) = x\}) < +\infty$$

for a.e.  $x \in \mathbf{R}^N$ .

**Remarks.**

a) By the first statement in Proposition 3.1,  $\mathcal{N}(t, x) = \mathcal{N}_{\mathbf{R}^N}(t, x) \geq 1$  for all  $(t, x) \in \mathbf{R} \times \mathbf{R}^N$ .

b) Even in the smooth case, i.e. assuming in addition that  $U^{in} \in C^1(\mathbf{R}^N)$ , statement c) in Theorem 3.4 provides information on the number of folds of  $\Lambda_t$ , that is the image under  $\Phi_t$  of the graph of  $U^{in}$ , which seems to be new at the time of this writing.

*Proof.* By the second estimate in Lemma 2.1

$$(24) \quad |D_x X_t(y, U^{in}(y)) + D_\xi X_t(y, U^{in}(y)) \cdot DU^{in}(y)| \leq e^{\kappa|t|} + (e^{\kappa|t|} - 1)|DU^{in}(y)|$$

so that

$$(25) \quad \begin{aligned} J_t(y) &= |\det(D_x X_t(y, U^{in}(y)) + D_\xi X_t(y, U^{in}(y)) \cdot DU^{in}(y))| \\ &\leq e^{N\kappa|t|}(1 + (1 - e^{-\kappa|t|})|DU^{in}(y)|)^N \end{aligned}$$

by Hadamard's inequality. Since  $U^{in}$  satisfies (18) and since  $L^{N,1}(\Omega) \subset L^N(\Omega)$ , this inequality implies statement a).

Since the map  $F_t$  is proper by Proposition 3.1, the set  $K_{t,R} = F_t^{-1}(\overline{B(0,R)})$  is compact for each  $R > 0$ . By the area formula (see Theorem 3.4 in [18] and Theorem A in [13])

$$\int_{\mathbf{R}^N} \#(F_t^{-1}(\{x\}) \cap K_{t,R}) dx = \int_{K_{t,R}} J_t(y) dy < +\infty.$$

Therefore  $\#F_t^{-1}(\{x\}) < \infty$  for a.e.  $x \in \overline{B(0,R)}$ ; since this is true for all  $R \in \mathbf{N}$ , one concludes that  $\#F_t^{-1}(\{x\}) < \infty$  for a.e.  $x \in \mathbf{R}^N$ , so that b) holds.

Let  $B$  be a measurable subset of  $\mathbf{R}^N$ ; applying again the area formula shows that

$$\int_{\mathbf{R}^N} \mathcal{N}_B(t, x) dx = \int_B J_t(y) dy.$$

By the Bienaymé-Chebyshev inequality, for each  $n \geq 1$

$$\mathcal{L}^N(\{x \in \mathbf{R}^N \mid \mathcal{N}_B(t, x) \geq n\}) \leq \frac{1}{n} \int_B J_t(y) dy,$$

which is precisely the first inequality in c). The second inequality follows from (25) and Hölder's inequality.

Let  $n = 1$  and  $Z_t := J_t^{-1}(\{0\}) \cup E$ . Let  $B = Z_t \cap K_{t,R}$ ; the set  $B$  is measurable and bounded. Then

$$\int_B J_t(y) dy = 0$$

since  $J_t(y) = 0$  for all  $y \in B \setminus E$  and  $\mathcal{L}^N(E) = 0$ . Applying c) shows that

$$\mathcal{L}^N(\{x \in \mathbf{R}^N \mid |x| \leq R \text{ and } \mathcal{N}_{Z_t}(t, x) \geq 1\}) = 0,$$

which entails d) by monotone convergence, letting  $R \in \mathbf{N}$  tend to infinity.

Consider next the continuous map

$$F : [-T, T] \times \mathbf{R}^N \ni (t, y) \mapsto F(t, y) \in \mathbf{R}^N.$$

In view of (16),  $|F(t, y)| \rightarrow \infty$  as  $|y| \rightarrow +\infty$  uniformly in  $t \in [-T, T]$ . Therefore, the set  $K_R := F^{-1}(\overline{B(0,R)})$  is compact for each  $R > 0$ . Then, for each  $t \in [-T, T]$  and each  $y \in \mathbf{R}^N \setminus E$ , the Jacobian  $DF(t, y)$  is the column-wise partitioned matrix

$$DF(t, y) = [V(t, y), M(t, y)],$$

with

$$V(t, y) = \nabla_\xi H(\Phi_t(y, U^{in}(y)))$$

and

$$M(t, y) := D_x X_t(y, U^{in}(y)) + D_\xi X_t(y, U^{in}(y)) DU^{in}(y).$$

Therefore,

$$DF(t, y) DF(t, y)^T = V(t, y) V(t, y)^T + M(t, y) M(t, y)^T$$

so that, by the co-area formula (Theorem 1.3 in [19])

$$\begin{aligned} & \int_{\mathbf{R}^N} \mathcal{H}^1(F^{-1}(\{x\}) \cap K_R) dx \\ &= \int_{K_R} \sqrt{\det(V(t, y)V(t, y)^T + M(t, y)M(t, y)^T)} dt dy. \end{aligned}$$

By Lemma 2.1,  $(t, x, \xi) \mapsto \Phi_t(x, \xi)$  is of class  $C^1$  on  $\mathbf{R} \times \mathbf{R}^N \times \mathbf{R}^N$ , so that the map  $(t, y) \mapsto V(t, y)$  is continuous on  $\mathbf{R} \times \mathbf{R}^N$ , and therefore bounded on the compact  $K_R$ . On the other hand, by (24)

$$\sup_{|t| \leq T} |M(t, y)| \leq e^{\kappa T} + (e^{\kappa T} - 1) |DU^{in}(y)| \in L_{loc}^N(\mathbf{R}^N),$$

since  $U^{in}$  satisfies (18). Denoting

$$K'_R := \{y \in \mathbf{R}^N \mid \text{there exists } t \in [-T, T] \text{ s.t. } (t, y) \in K_R\}$$

that is compact in  $\mathbf{R}^N$  (being the projection of the compact  $K_R$  on the second factor in  $\mathbf{R} \times \mathbf{R}^N$ ), one has

$$\begin{aligned} \|VV^T + MM^T\|_{L^{N/2}(K_R)}^{N/2} &\leq 2^{N/2-1} \|V\|_{L^\infty(K_R)}^N \mathcal{L}^{N+1}(K_R) + 2^{N/2} T \|M\|_{L^N(K'_R)}^N \\ &< \infty. \end{aligned}$$

Therefore  $\mathcal{H}^1(F^{-1}(\{x\}) \cap K_R) < +\infty$  is finite for a.e.  $x \in \overline{B(0, R)}$ , and since this is true for all  $R \in \mathbf{N}$ , one concludes that  $\mathcal{H}^1(F^{-1}(\{x\})) < +\infty$  for a.e.  $x \in \mathbf{R}^N$ , which is statement e).  $\square$

Theorem 3.4 suggests considering the sets

$$(26) \quad C'_t = \{x \in \mathbf{R}^N \mid F_t^{-1}(\{x\}) \text{ is infinite}\}$$

for all  $t \in \mathbf{R}$ .

On the other hand, the definition of the caustic (15) in the smooth case should be modified as follows when the continuous vector field  $U^{in}$  is not of class  $C^1$  but satisfies (11) and (18):

$$(27) \quad \begin{aligned} C_t &:= \{x \in \mathbf{R}^N \mid F_t^{-1}(\{x\}) \cap Z_t \neq \emptyset\}, \\ C &:= \bigcup_{t \in \mathbf{R}} C_t, \end{aligned}$$

where  $Z_t$  is defined in (22). Notice that this definition coincides with (15) whenever  $U^{in}$  is differentiable everywhere. However we do not know whether  $C$  or  $C_t$  are closed whenever  $U^{in}$  is not of class  $C^1$ .

The first part of statement c) in Proposition 3.3 is equivalent to the inclusion

$$(28) \quad C'_t \subset C_t \quad \text{for all } t \in \mathbf{R}.$$

Statements b) and d) in Theorem 3.4 can be recast as follows

$$(29) \quad \mathcal{L}^N(C'_t) = 0 \text{ and } \mathcal{L}^N(C_t) = 0, \quad \text{for all } t \in \mathbf{R}.$$

This is the analogue of statement a) in Proposition 3.3 in the case of a rough  $U^{in}$ . Notice that (29) is a consequence of the area formula while statement a) in Proposition 3.3 follows from Sard's theorem (see Remark 2.97 on pp. 103–104 in [2], discussing the relation between Sard's theorem and the co-area formula).

## 4. SOME EXAMPLES

While the notion of caustic  $C$  naturally occurs in studying the geometry of  $\Lambda_t$  in the smooth case, its relation to the number of solutions  $y$  of the equation  $F_t(y) = x$  is slightly less obvious, as shown by the following examples. In all these examples, the Hamiltonian is

$$H(x, \xi) = \frac{1}{2}\xi^2$$

generating the free flow

$$\Phi_t : \mathbf{R} \times \mathbf{R} \ni (x, \xi) \mapsto (x + t\xi, \xi) \in \mathbf{R} \times \mathbf{R}.$$

Thus

$$F_t : \mathbf{R} \ni y \mapsto y + tU^{in}(y) \in \mathbf{R}.$$

The first example below shows that the set  $F_t^{-1}(\{x\})$  may be finite even if  $x \in C_t$ . In other words, it may happen that the inclusion (28) is strict for all  $t \in \mathbf{R}$ .

**Example 1.** Set  $N = 1$ , and let  $U^{in}$  be real analytic on  $\mathbf{R}$  and satisfy (11). Therefore  $F_t$  is real analytic on  $\mathbf{R}$  for each  $t \in \mathbf{R}$ . For each  $x \in \mathbf{R}$ , the set  $F_t^{-1}(\{x\})$  is the set of zeros of the analytic function  $y \mapsto F_t(y) - x$ . Since  $U^{in}$  satisfies (11), one has  $F_t(y) \sim y$  as  $|y| \rightarrow \infty$ , so that, for each  $t \in \mathbf{R}$ , the function  $F_t$  is not a constant. Therefore the zeros of  $y \mapsto F_t(y) - x$  are isolated for each  $(t, x) \in \mathbf{R} \times \mathbf{R}$ . Equivalently, the set  $F_t^{-1}(\{x\})$  consists of isolated points. On the other hand, for each  $(t, x) \in \mathbf{R}^2$ , the set  $F_t^{-1}(\{x\})$  is compact since  $F_t$  is proper by Proposition 3.1. Therefore the set  $F_t^{-1}(\{x\})$  is finite for all  $(t, x) \in \mathbf{R}^2$ .

The next example shows that the set  $F_t^{-1}(\{x\})$  may be infinite for infinitely many times  $t$  — in this case, for all  $t$  in a nonempty, open interval of  $\mathbf{R}$ .

**Example 2.** Set  $N = 1$ , and let  $U^{in}$  be defined by

$$U^{in}(z) := \begin{cases} \tanh(z) \sin(\ln |z|) & \text{if } z > 0, \\ 0 & \text{if } z = 0. \end{cases}$$

Clearly  $U^{in} \in C^1(\mathbf{R}^*)$  and, for all  $z \in \mathbf{R}^*$ , one has

$$(U^{in})'(z) = (1 - \tanh^2(z)) \sin(\ln |z|) + \frac{\tanh(z)}{z} \cos(\ln |z|).$$

Observe that  $(U^{in})'(z) = \sqrt{2} \sin(\ln |z| + \frac{\pi}{4}) + O(z^2)$  does not have a limit for  $z \rightarrow 0$ , so that  $U^{in} \notin C^1(\mathbf{R})$ . On the other hand

$$\sup_{z \in \mathbf{R}} |U^{in}(z)| = 1 \quad \text{and} \quad \sup_{z \neq 0} |(U^{in})'(z)| \leq 2$$

so that  $U^{in}$  is Lipschitz continuous on  $\mathbf{R}$  and the map  $F_t$  is proper on  $\mathbf{R}$ .

Let  $x = 0$ ; for each  $t$  such that  $|t| > 1$ , the set  $F_t^{-1}(\{0\})$  is infinite.

Indeed, the set  $F_t^{-1}(\{0\})$  obviously contains  $z = 0$ ; besides,  $F_t^{-1}(\{0\})$  contains also a decreasing sequence  $y_n \rightarrow 0$  as  $n \rightarrow +\infty$ , which satisfies

$$\sin(\ln |y_n|) \rightarrow -1/t \quad \text{as } n \rightarrow \infty.$$

Therefore  $F_t^{-1}(\{0\})$  is countably infinite whenever  $|t| > 1$ .

Notice that, in example 2, whenever  $T > 1$

$$\{(t, y) \in [-T, T] \times \mathbf{R} \mid F_t(y) = 0\} = ([-T, -1] \cup [1, T]) \times (\{0\} \cup \{y_n \mid n \geq 1\})$$

so that

$$\mathcal{H}^1(\{(t, y) \in [-T, T] \times \mathbf{R} \mid F_t(y) = 0\}) = +\infty.$$

In other words, the set of points  $x$  for which

$$\mathcal{H}^1(\{(t, y) \in [-T, T] \times \mathbf{R} \mid F_t(y) = x\}) = +\infty,$$

while  $\mathcal{L}^1$ -negligible by statement e) of Theorem 3.4, may be non empty.

There is another interesting observation in connection with Example 2. Observe that  $0 \in C'_t$  and that

$$F_t^{-1}(\{0\}) = \{0\} \cup \{y_n \mid n \geq 1\}, \quad \text{whenever } |t| > 1.$$

Nevertheless, for each  $n \geq 1$

$$(U^{in})'(y_n) = \sin(\ln |y_n|) + \cos(\ln |y_n|) + O(|y_n|^2) \rightarrow -\frac{1}{t} + \sqrt{1 - \frac{1}{t^2}} \neq -\frac{1}{t}$$

if  $|t| > 1$ , and  $F_t$  is not differentiable at  $y = 0$ . Thus  $y = 0$  is in the exceptional set  $E$  where the Lipschitz continuous function  $U^{in}$  is not differentiable, so that  $0 \in C_t$  with the definition (27) although

$$F_t^{-1}(\{0\}) \cap J_t^{-1}(\{0\}) = \emptyset,$$

so that 0 would not belong to  $C_t$  had we kept the classical definition (15) in the case of non everywhere differentiable  $U^{in}$  profiles.

The next two examples show that  $F_t^{-1}(\{x\})$  can even be uncountably infinite, even for a smooth profile  $U^{in}$ .

**Example 3.** Set  $N = 1$ , and let  $U^{in}$  be defined by

$$U^{in}(z) := \begin{cases} +1 & \text{if } z < -1, \\ -z & \text{if } |z| \leq 1, \\ -1 & \text{if } z > 1. \end{cases}$$

Consider the equation

$$F_t(y) := y + tU^{in}(y) = x$$

with unknown  $y$ ; for  $t < 1$ , its solution is unique and given by

$$y = \begin{cases} x - t & \text{if } x < t - 1, \\ \frac{x}{1 - t} & \text{if } |x| \leq 1 - t, \\ x + t & \text{if } x > 1 - t. \end{cases}$$

For  $t = 1$ , the solution is

$$y = \begin{cases} x - t & \text{if } x < 0, \\ \text{any } y \in [-1, 1] & \text{if } x = 0, \\ x + 1 & \text{if } x > 0. \end{cases}$$

For  $t > 1$ , the solution is

$$y = \begin{cases} x - t & \text{if } x < 1 - t, \\ x - t, \frac{x}{1 - t} \text{ and } x + t & \text{if } 1 - t \leq x \leq t - 1, \\ x + t & \text{if } x > t - 1. \end{cases}$$

Now  $F_t^{-1}(\{x\})$  is finite for all  $x \in \mathbf{R}$  whenever  $t \neq 1$ , while  $F_1^{-1}(\{0\}) = [-1, 1]$ .

**Example 4.** In the previous example,  $U^{in}$  is Lipschitz continuous but not of class  $C^1$ . Yet the same phenomenon can be observed by smoothing  $U^{in}$  near  $z = \pm 1$ . Regularize  $U^{in}$  and obtain  $U_\epsilon^{in} \in C^\infty(\mathbf{R})$  so that

$$\begin{aligned} \text{supp}(U_\epsilon^{in} - U^{in}) &\subset [-1 - \epsilon, -1 + \epsilon] \cup [1 - \epsilon, 1 + \epsilon], \quad \text{and} \\ U_\epsilon^{in} &\leq U^{in} \text{ on } [-1 - \epsilon, -1 + \epsilon] \text{ and } U^{in} \leq U_\epsilon^{in} \text{ on } [1 - \epsilon, 1 + \epsilon]. \end{aligned}$$

In that case

$$F_1^{-1}(\{0\}) = [-1 + \epsilon, 1 - \epsilon].$$

Indeed, since  $U_\epsilon^{in} \leq U^{in}$  on  $[-1 - \epsilon, -1 + \epsilon]$ , all the points on the graph of  $U_\epsilon^{in}$  with abscissa in  $(-1 - \epsilon, -1 + \epsilon)$  will reach  $x = 0$  after  $t = 1$ . The same is true of the points on the graph of  $U_\epsilon^{in}$  with abscissa in  $(1 - \epsilon, 1 + \epsilon)$ . Thus the regularization does not affect the dynamics of the points with abscissa in  $(-1 + \epsilon, 1 - \epsilon)$  for all  $t \in [0, 1]$ , and in particular for  $t = 1$ .

## 5. ON THE STRUCTURE OF $\mu(t)$

Throughout this section, we assume that  $U^{in} \in C(\mathbf{R}^N; \mathbf{R}^N)$  satisfies the sublinearity condition (11) at infinity and the regularity condition (18).

Consider a monokinetic measure  $\mu^{in}$  of the form (3) with  $\rho^{in} \in L^1(\mathbf{R}^N)$ , whose action on a test function  $\chi \in C_b(\mathbf{R}^N \times \mathbf{R}^N)$  is given by the formula

$$\langle \mu^{in}, \chi \rangle := \int_{\mathbf{R}^N} \chi(y, U^{in}(y)) \rho^{in}(y) dy.$$

In other words

$$\mu^{in} = \mathcal{L}_x^N \otimes \rho^{in}(x) \delta_{U^{in}(x)}^{in}.$$

Let  $H \in C^2(\mathbf{R}^N \times \mathbf{R}^N)$  satisfy (7), and let  $\Phi_t$  be the Hamiltonian flow generated by  $H$ .

For all  $t \in \mathbf{R}$ , let  $\mu(t) = \Phi_t \# \mu^{in}$  be the push-forward of  $\mu^{in}$  under  $\Phi_t$ , defined as follows: for each test function  $\chi \in C_b(\mathbf{R}^N \times \mathbf{R}^N)$ ,

$$(30) \quad \langle \mu(t), \chi \rangle := \langle \mu^{in}, \chi \circ \Phi_t \rangle = \int_{\mathbf{R}^N} \chi(\Phi_t(y), U^{in}(y)) \rho^{in}(y) dy.$$

Finally, let  $\rho(t)$  be the measure on  $\mathbf{R}^N$  defined as

$$(31) \quad \rho(t) := \Pi \# \mu(t)$$

where  $\Pi : T^*\mathbf{R}^N \simeq \mathbf{R}^N \times \mathbf{R}^N \ni (x, \xi) \mapsto x \in \mathbf{R}^N$  is the canonical projection. In other words, for each test function  $\phi \in C_b(\mathbf{R}^N)$

$$(32) \quad \langle \rho(t), \phi \rangle = \int_{\mathbf{R}^N} \phi(X_t(y), U^{in}(y)) \rho^{in}(y) dy$$

or, equivalently

$$(33) \quad \rho(t) = F_t \# \rho^{in}.$$

We shall also use the following definition

$$(34) \quad P_t := \{y \in \mathbf{R}^N \setminus E \mid J_t(y) > 0\}.$$

Our main result in this section bears on the structure of the measures  $\mu(t)$  and its projection  $\rho(t)$ .

**Theorem 5.1.** *Under the assumptions above, let  $\rho^{in} \in L^1(\mathbf{R}^N)$  be such that  $\rho^{in} \geq 0$  a.e. on  $\mathbf{R}^N$ .*

a) for each  $t \in \mathbf{R}$ ,

$$\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N \ll J_t \mathcal{L}^N, \quad \text{and } \rho^{in} \mathbf{1}_{Z_t} \mathcal{L}^N \perp J_t \mathcal{L}^N$$

b) for each  $t \in \mathbf{R}$ ,

$$\mu(t) = \mu_a(t) + \mu_s(t) \quad \text{with } \mu_a(t) \perp \mu_s(t),$$

where

$$\begin{cases} \mu_a(t) := \mathcal{L}_x^N \otimes \sum_{y \in F_t^{-1}(\{x\})} \frac{\rho^{in} \mathbf{1}_{P_t}(y)}{J_t} \delta_{\Xi_t(y, U^{in}(y))}, \\ \mu_s(t) := \Phi_t \# (\mathcal{L}_x^N \otimes \rho^{in}(x) \mathbf{1}_{Z_t}(x) \delta_{U^{in}(x)}). \end{cases}$$

Moreover

c) for each  $t \in \mathbf{R}$ , one has

$$\text{supp}(\mu(t)) \subset \Lambda_t := \Phi_t(\{(y, U^{in}(y)) \mid y \in \mathbf{R}^N\});$$

d) for each  $t \in \mathbf{R}$ , the measure

$$\rho_a(t) := \Pi \# \mu_a(t) \ll \mathcal{L}^N;$$

with

$$\frac{d\rho_a(t)}{d\mathcal{L}^N}(x) = \mathbf{1}_{\mathbf{R}^N \setminus C_t'}(x) \sum_{y \in F_t^{-1}(\{x\})} \frac{d(\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N)}{d(J_t \mathcal{L}^N)}(y)$$

where  $C_t'$  is defined in (26);

e) for each  $t \in \mathbf{R}$ , the measure

$$\rho_s(t) := \Pi \# \mu_s(t) \text{ is carried by } C_t;$$

in particular

$$\rho_a(t) \perp \rho_s(t).$$

A few remarks are in order before we give the proof of Theorem 5.1. In the smooth case — i.e. whenever  $U^{in} \in C^1(\mathbf{R}^N; \mathbf{R}^N)$  and satisfies (11), we recall that  $C_t$  is closed in  $\mathbf{R}^N$  for each  $t \in \mathbf{R}$ , by statement b) of Proposition 3.3. For any given  $t \in \mathbf{R}$ , let  $\chi \equiv \chi(x, \xi)$  be a test function in  $C_c(\mathbf{R}^N \times \mathbf{R}^N)$  such that  $\Pi \text{supp}(\chi) \cap C_t = \emptyset$ . By (30)

$$\langle \mu(t), \chi \rangle = \int_{\mathbf{R}^N} \chi(\Phi_t(y, U^{in}(y))) \rho^{in}(y) dy.$$

Since  $\Pi(\text{supp}(\chi))$  is compact and included in the open set  $\mathbf{R}^N \setminus C_t$ , it intersects at most finitely many connected components of  $\mathbf{R}^N \setminus C_t$ . Assume without loss of generality that  $\Pi(\text{supp}(\chi))$  is connected, so that it intersects exactly one connected component  $\Omega$  of  $\mathbf{R}^N \setminus C_t$ ; on  $\Omega$ , the integer-valued function  $\mathcal{N}$  is a constant denoted by  $\mathcal{N}^\Omega$ , by statement c) of Proposition 3.3. With the notation used in that proposition, for each  $x \in \Pi(\text{supp}(\chi)) \subset \Omega$

$$F_t^{-1}(\{x\}) = \{y_j(t, x) \mid j = 1, \dots, \mathcal{N}^\Omega\},$$

and

$$y_j(t, \cdot) \in C^1(\Omega) \text{ for all } j = 1, \dots, \mathcal{N}^\Omega.$$



Therefore  $y_j(t, \cdot)$  is a  $C^1$ -diffeomorphism from  $\Omega$  on its image  $O_j$ , with inverse  $F_t$ . Thus

$$F_t^{-1}(\Omega) = \bigcup_{j=1}^{\mathcal{N}^\Omega} O_j \quad \text{and } O_i \cap O_j = \emptyset \text{ if } i \neq j,$$

so that

$$\iint_{\Omega \times \mathbf{R}^N} \chi(t, x) \mu(t, dx d\xi) = \sum_{j=1}^{\mathcal{N}^\Omega} \int_{O_j} \chi(F_t(y), \Xi_t(y, U^{in}(y))) \rho^{in}(y) dy.$$

In each of the integrals on the right hand side,  $F_t$  is a  $C^1$ -diffeomorphism mapping  $O_j$  on  $\Omega$ , so that, changing variables, we see that

$$\begin{aligned} & \int_{O_j} \chi(F_t(y), \Xi_t(y, U^{in}(y))) \rho^{in}(y) dy \\ &= \int_{\Omega} \chi(x, \Xi_t(y_j(t, x), U^{in}(y_j(t, x)))) \rho^{in}(y_j(t, x)) |\det(\nabla_x y_j(t, x))| dx. \end{aligned}$$

Since

$$|\det(D_x y_j(t, x))| = J_t(y_j(t, x))^{-1},$$

we conclude that the restriction of  $\mu(t)$  to  $(\mathbf{R} \times \mathbf{R}^N) \setminus C$  is a measure-valued function of  $(t, x)$  given by the following formula:

$$(35) \quad \mu(t, x, \cdot) := \sum_{j=1}^{\mathcal{N}(t, x)} \frac{\rho^{in}(y_j(t, x))}{J_t(y_j(t, x))} \delta_{\Xi_t(y_j(t, x), U^{in}(y_j(t, x)))}$$

whenever  $(t, x) \notin C$ . This formula is strikingly similar to the one giving  $\mu_a(t)$  in statement b) of Theorem 5.1. There are however subtle differences, which we shall discuss in more detail in the next section. At this point, it suffices to say that Theorem 5.1 provides a formula for  $\mu(t)$  that holds globally on  $\mathbf{R}^N \times \mathbf{R}^N$  instead of  $(\mathbf{R}^N \setminus C_t) \times \mathbf{R}^N$ , and that the argument above requires more regularity on  $U^{in}$  than assumed in Theorem 5.1.

*Proof of Theorem 5.1.* Let  $A \subset \mathbf{R}^N$ ; then the condition

$$\int_A J_t(y) dy = 0 \text{ implies that } J_t(y) = 0 \text{ for a.e. } y \in A.$$

Therefore  $\mathcal{L}^N(P_t \cap A) = 0$  so that

$$\int_A (\rho^{in} \mathbf{1}_{P_t})(y) dy = \int_{P_t \cap A} \rho^{in}(y) dy = 0.$$

Thus  $\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N \ll J_t \mathcal{L}^N$ .

On the other hand, for each  $t \in \mathbf{R}$

$$\mathbf{R}^N = P_t \cup Z_t \quad \text{with } P_t \cap Z_t = \emptyset.$$

Since

$$J_t(y) = 0 \text{ for } y \in Z_t \setminus E \quad \text{i.e. } \mathcal{L}^N\text{-a.e. on } Z_t$$

while

$$\rho^{in}(y) \mathbf{1}_{Z_t}(y) = 0 \quad \text{for all } y \in P_t,$$

we conclude that  $\rho^{in} \mathbf{1}_{Z_t} \mathcal{L}^N \perp J_t \mathcal{L}^N$ , which proves a).

Define

$$\begin{aligned}\mu_a^{in} &:= \mathcal{L}_x^N \otimes (\rho^{in} \mathbf{1}_{P_t})(x) \delta_{U^{in}(x)}^{in}, \\ \mu_s^{in} &:= \mathcal{L}_x^N \otimes (\rho^{in} \mathbf{1}_{Z_t})(x) \delta_{U^{in}(x)}^{in},\end{aligned}$$

and, for each  $t \in \mathbf{R}$

$$\mu_a(t) := \Phi_t \# \mu_a^{in}, \quad \mu_s(t) := \Phi_t \# \mu_s^{in},$$

so that one has indeed

$$\mu_a(t) + \mu_s(t) = \Phi_t \# (\mu_a^{in} + \mu_s^{in}) = \Phi_t \# \mu^{in} = \mu(t).$$

Then there exists a unique  $b \in L^1(\mathbf{R}^N; J_t \mathcal{L}^N)$  such that

$$\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N = b J_t \mathcal{L}^N$$

by the Radon-Nikodym theorem. Thus, for each  $\chi \in C_c(\mathbf{R}^N \times \mathbf{R}^N)$ , by the area formula (see Theorem 3.4 in [18] and Theorem A in [13])

$$\begin{aligned}\langle \mu_a(t), \chi \rangle &= \int_{\mathbf{R}^N} \chi(F_t(y), \Xi_t(y, U^{in}(y))) b(y) J_t(y) dy \\ &= \int_{\mathbf{R}^N} \left( \sum_{y \in F_t^{-1}(\{x\})} b(y) \chi(x, \Xi_t(y, U^{in}(y))) \right) dx \\ &= \int_{\mathbf{R}^N} \left( \sum_{y \in F_t^{-1}(\{x\})} b(y) \langle \delta_{\Xi_t(y, U^{in}(y))}, \chi(x, \cdot) \rangle \right) dx.\end{aligned}$$

In the formula above

$$b := \frac{d(\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N)}{d(J_t \mathcal{L}^N)}$$

is the Radon-Nikodym derivative of  $\rho^{in} \mathbf{1}_{P_t}$  with respect to  $J_t \mathcal{L}^N$ . Since  $J_t > 0$  on the set  $P_t$

$$(36) \quad b = \frac{\rho^{in} \mathbf{1}_{P_t}}{J_t} \text{ a.e. on } \mathbf{R}^N,$$

which proves b).

Formula (3) obviously implies that

$$\text{supp}(\mu^{in}) \subset \Lambda_0 := \{(y, U^{in}) \mid y \in \mathbf{R}^N\}.$$

Since  $\mu(t) = \Phi_t \# \mu^{in}$ , one has

$$\text{supp}(\mu(t)) \subset \Phi_t(\{(y, U^{in}) \mid y \in \mathbf{R}^N\}) = \Lambda_t,$$

which is precisely statement c).

In view of the first formula in b) and of the definition  $\rho_a(t) = \Pi \# \mu_a(t)$  of the measure  $\rho_a(t)$ , one has

$$\rho_a(t) = \left( \sum_{y \in F_t^{-1}(\{x\})} b(y) \right) \mathcal{L}^N,$$

with  $b$  as in (36). The set  $F_t^{-1}(\{x\})$  can obviously be infinite, in which case the sum above can be undefined. However, this occurs only if  $x \in C'_t$  as defined in (26). Since  $C'_t$  is  $\mathcal{L}^N$ -negligible

$$\rho_a(t) = f_t \mathcal{L}^N$$

with

$$f_t(x) := \mathbf{1}_{\mathbf{R}^N \setminus C_t'}(x) \left( \sum_{y \in F_t^{-1}(\{x\})} b(y) \right).$$

Thus,  $\rho_a(t)$  is of the form  $\rho_a(t) = f_t \mathcal{L}^N$  with  $f_t \geq 0$  measurable on  $\mathbf{R}^N$ . Besides

$$0 \leq \rho_a(t) \leq \rho(t) \quad \text{and} \quad \int_{\mathbf{R}^N} \rho(t, dx) = \int_{\mathbf{R}^N} \rho^{in}(x) dx < \infty$$

which implies in particular that  $\rho_a(t) \ll \mathcal{L}^N$  with the formula for the Radon-Nikodym derivative as in d).

Consider the measurable set  $A := \mathbf{R}^N \setminus C_t$ . By (33) applied to  $\rho^{in} \mathbf{1}_{Z_t}$  instead of  $\rho^{in}$ , one has

$$\rho_s(t)(A) = \int_{F_t^{-1}(A)} \rho^{in}(y) \mathbf{1}_{Z_t}(y) dy = \int_{F_t^{-1}(A) \cap Z_t} \rho^{in}(y) dy = 0.$$

Indeed, by definition of  $C_t$ , one has  $F_t^{-1}(A) \cap Z_t = \emptyset$ . In other words,  $\rho_s(t)$  is carried by  $C_t$ .

Finally, since  $\rho_a(t) \ll \mathcal{L}^N$  and  $\rho_s(t)$  is carried by  $C_t$  which is  $\mathcal{L}^N$ -negligible by (29), we conclude that  $\rho_a(t) \perp \rho_s(t)$  which is precisely statement e).  $\square$

## 6. ON THE CAUSTIC AND OTHER EXCEPTIONAL SETS

In the case of a smooth  $U^{in}$  profile — i.e. when  $U^{in} \in C^1(\mathbf{R}^N; \mathbf{R}^N)$  satisfies (11), the caustic  $C$  is the only exceptional set —  $C_t$  being equivalently defined as the image under the projection  $\Pi$  of the set of points in the manifold  $\Lambda_t$  in (12) where the restriction  $\Pi|_{\Lambda_t}$  is not differentiable.

When  $U^{in}$  is not everywhere differentiable, this definition of  $C_t$  does not make sense in general since  $\Lambda_t$  is not a differentiable manifold in the first place. In such cases, it is more natural to consider the measures  $\mu(t)$  and  $\rho(t)$  instead of the sets  $\Lambda_t$  and  $C_t$  — all the more so since  $C_t$  may not even be closed in  $\mathbf{R}^N$ . Thus, even though  $\rho_s(t)$  is concentrated on  $C_t$ , one cannot say that  $C_t$  is the support of  $\rho_s(t)$  as  $C_t$  may not be closed. On the other hand, the inclusion  $\text{supp } \rho_s(t) \subset \overline{C_t}$  is of little interest as  $C_t$  might be dense in some domain of  $\mathbf{R}^N$ . Although  $\rho_s(t)$  is concentrated on  $C_t$ , this obviously does not characterize  $C_t$  (if a measure is concentrated on a set, it is also concentrated on the complement in that set of any negligible set for that same measure).

There are analogous difficulties with the absolutely continuous part of the measure  $\rho_a(t)$ . In formula (35), the restriction of  $\mu_a(t)$  to  $(\mathbf{R}^N \setminus C_t) \times \mathbf{R}^N$  is viewed as a function of  $x \in \mathbf{R}^N \setminus C_t$  with values in the set of Radon measures in the variable  $\xi \in \mathbf{R}^N$ . This viewpoint is obviously not appropriate if  $U^{in}$  is not at least of class  $C^1$  — for instance, if the set  $C_t$  is dense in some domain of  $\mathbf{R}^N$ . In Theorem 5.1, the measure  $\mu$  is a weakly continuous function of the time variable  $t$  with values in the space of Radon measures in the variables  $(x, \xi)$ , and is therefore globally defined on  $\mathbf{R}_x^N \times \mathbf{R}_\xi^N$ . Obviously, the ratio  $\rho^{in} \mathbf{1}_{P_t} / J_t$  is just one possible choice of the Radon-Nikodym derivative  $d(\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N) / d(J_t \mathcal{L}^N)$  and could be modified arbitrarily on any set of  $J_t \mathcal{L}^N$ -measure 0 — which could be of positive  $\mathcal{L}^N$ -measure,

as in Examples 3-4 above. The difference induced in the expression

$$(37) \quad \sum_{y \in F_t^{-1}(\{x\})} \frac{d(\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N)}{d(J_t \mathcal{L}^N)}(y) \delta_{U^{in}(y)}$$

by two different choices of the Radon-Nikodym derivative  $d(\rho^{in} \mathbf{1}_{P_t} \mathcal{L}^N)/d(J_t \mathcal{L}^N)$  is of the form

$$\sum_{y \in F_t^{-1}(\{x\})} m(y) \delta_{U^{in}(y)}$$

with  $m \geq 0$  being a  $J_t \mathcal{L}^N$ -measurable function such that  $m = 0$  a.e. on  $\mathbf{R}^N \setminus Z_t$ . Therefore

$$\int_{\mathbf{R}^N} m(y) J_t(y) dy = 0 = \int_{\mathbf{R}^N} \left( \sum_{y \in F_t^{-1}(\{x\})} |m(y)| \right) dx$$

so that

$$\sum_{y \in F_t^{-1}(\{x\})} m(y) = 0 \quad \mathcal{L}^N - \text{a.e. on } \mathbf{R}^N$$

and

$$\mathcal{L}^N \otimes \sum_{y \in F_t^{-1}(\{x\})} m(y) \delta_{U^{in}(y)} = 0$$

as a measure on  $\mathbf{R}^N \times \mathbf{R}^N$ .

On the other hand, even in the smooth case, formula (35) is not enough to define completely  $\mu(t)$ , as it fails to capture singular parts of the measure carried by  $C_t \times \mathbf{R}^N$ . For instance, in Example 4, for  $t = 1$ , if one computes the restriction of the measure  $\mu(1)$  to  $(\mathbf{R}^N \setminus C_1) \times \mathbf{R}^N$  by formula (35), one obtains

$$\begin{aligned} \int_{\mathbf{R}^N \setminus C_1} \left( \int_{\mathbf{R}^N} \mu(t, x, d\xi) \right) dx &= \int_{-\infty}^{-1+\epsilon} \rho^{in}(y) dy + \int_{1-\epsilon}^{+\infty} \rho^{in}(y) dy \\ &\neq \int_{\mathbf{R}^N} \rho^{in}(y) dy = \int_{\mathbf{R}^N} \left( \int_{\mathbf{R}^N} \mu^{in}(x, d\xi) \right) dx \end{aligned}$$

unless  $\rho^{in} = 0$  on  $[-1 + \epsilon, 1 - \epsilon] = J_1^{-1}(\{0\})$ . Since this is obviously incompatible with the conservation of the total mass of the measure

$$\iint_{\mathbf{R}^N \times \mathbf{R}^N} \mu(t, dx d\xi) = \iint_{\mathbf{R}^N \times \mathbf{R}^N} \Phi_t \# \mu^{in}(dx d\xi) = \iint_{\mathbf{R}^N \times \mathbf{R}^N} \mu^{in}(dx d\xi)$$

by the transportation under the flow  $\Phi_t$ , this example confirms that (35) alone is not enough to compute  $\mu(t)$ .

The expression involved in the formula for  $\mu_a$  in Theorem 5.1, i.e.

$$(38) \quad \sum_{y \in F_t^{-1}(\{x\})} \frac{\rho^{in} \mathbf{1}_{P_t}}{J_t}(y) \delta_{U^{in}(y)}$$

which is analogous to formula (35) a priori makes sense only if  $x \in \mathbf{R}^N \setminus C'_t$ . In the rough case, i.e. if  $U^{in}$  satisfies only the regularity properties in Theorem 3.4 or Theorem 5.1, it might on principle happen that

$$C'_t \cap (\mathbf{R}^N \setminus C_t) \neq \emptyset.$$

Whenever this is the case, the sum in formula (38) above may fail to be defined everywhere on  $\mathbf{R}^N \setminus C_t$ , as would occur for  $U^{in}$  of class  $C^1$ , by statement c) in

Proposition 3.3, or (28). Since this may occur at most on a  $\mathcal{L}^N$ -negligible set, the expression for  $\mu_a$  in Theorem 5.1 is well defined as a measure on  $\mathbf{R}^N \times \mathbf{R}^N$  while the one in (38) may not be defined everywhere on  $\mathbf{R}^N \setminus C_t$  as a function of  $x$  with values in the space of measures in the variable  $\xi \in \mathbf{R}^N$ .

Another strategy is to abandon the idea of viewing the caustic  $C$  as an exceptional set unambiguously defined by the measure  $\rho(t)$ , and to investigate those exceptional sets that can be uniquely defined in terms of this measure.

This leads for instance to consider the set

$$(39) \quad C_t'' := \{x \in \mathbf{R}^N \mid \mathcal{L}^N(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})) > 0\},$$

which is of particular interest, as shown by the following result.

**Theorem 6.1.** *Assume that  $H \in C^2(\mathbf{R}^N \times \mathbf{R}^N)$  satisfies the conditions (7), while  $U^{in} \in C(\mathbf{R}^N; \mathbf{R}^N)$  satisfies the sublinearity condition (11) at infinity and the regularity condition (18). Then*

- a)  $C_t'' \subset C_t \cap C_t'$  for each  $t \in \mathbf{R}$ ;
- b) for each  $t \in \mathbf{R}$  and each  $x \in C_t''$ , there exists  $\rho^{in} \in L^1(\mathbf{R}^N)$  such that  $\rho^{in} \geq 0$  a.e. and the projected measure  $\rho(t) = \Pi\#\mu(t)$  satisfies

$$\rho(t)(\{x\}) > 0;$$

- c)  $C_t''$  is at most countable for each  $t \in \mathbf{R}$ ;
- d) if  $\rho^{in} \in L^1(\mathbf{R}^N)$  is such that  $\rho^n \geq 0$  a.e. on  $\mathbf{R}^N$  and  $\rho^{in} = 0$  a.e. on  $J_t^{-1}(\mathbf{R}_+^*)$  while  $\|\rho^{in}\|_{L^1(\mathbf{R}^N)} > 0$ , then

$$\rho(t)(\{x\}) > 0 \quad \text{if and only if } (\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})) > 0.$$

*Proof.* If  $x \in C_t''$ , then  $\mathcal{L}^N(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})) > 0$  so that in particular the intersection  $F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})$  is infinite, which implies that  $x \in C_t \cap C_t'$ , thereby proving a).

Let  $R > 0$ ; since  $F_t$  is proper by Proposition 3.1,  $F_t^{-1}(\overline{B(0, R)}) = K_R$  is a compact subset of  $\mathbf{R}^N$ . Pick  $\rho^{in} = \mathbf{1}_{K_R}$ ; obviously  $\rho^{in} \in L^1(\mathbf{R}^N)$  and  $\rho^{in} \geq 0$  a.e.. Set  $\mu^{in} = \mathcal{L}^N \otimes (\mathbf{1}_{K_R} \delta_{U^{in}})$ , let  $\mu(t) = \Phi\#\mu^{in}$  be the push-forward of  $\mu^{in}$  under the Hamiltonian flow and let  $\rho(t) = \Pi\#\mu(t)$  be the projected measure. As explained above  $\rho(t) = F_t\#(\mathbf{1}_{K_R} \mathcal{L}^N)$  so that, whenever  $x \in \overline{B(0, R)} \cap C_t''$ ,

$$\begin{aligned} \rho(t)(\{x\}) &= (\mathbf{1}_{K_R} \mathcal{L}^N)(F_t^{-1}(\{x\})) = \mathcal{L}^N(K_R \cap F_t^{-1}(\{x\})) \\ &= \mathcal{L}^N(F_t^{-1}(\{x\})) \\ &\geq \mathcal{L}^N(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})) > 0. \end{aligned}$$

This immediately implies both b) and c). Indeed, b) is satisfied with  $\rho^{in} = \mathbf{1}_{K_R}$  for each  $R > 0$  such that  $x \in \overline{C_t'' \cap B(0, R)}$ . On the other hand, for each  $R > 0$ , all the points in  $C_t'' \cap \overline{B(0, R)}$  are atoms of the same Borel measure  $F_t\#(\mathbf{1}_{K_R} \mathcal{L}^N)$  on  $\mathbf{R}^N$ , which implies c).

As for d), one has again  $\rho(t) = F_t\#(\rho^{in} \mathcal{L}^N)$ , so that

$$\begin{aligned} \rho(t)(\{x\}) &= (\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\})) = (\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\}) \cap E) \\ &\quad + (\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})) \\ &\quad + (\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\}) \cap J_t^{-1}(\mathbf{R}_+^*)) \end{aligned}$$

where  $E$  is the  $\mathcal{L}^N$ -negligible set where  $F_t$  is not differentiable. The conclusion follows from the fact that  $\mathcal{L}^N(E) = 0$  and  $\rho^{in} = 0$  a.e. on  $J_t^{-1}(\mathbf{R}_+^*)$ , so that

$$\rho(t)(\{x\}) = (\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\}))$$

□

If  $\rho^{in} \in L^1(\mathbf{R}^N)$  satisfies  $\rho^{in} \geq 0$  a.e. on  $\mathbf{R}^N$  and  $\rho^{in} = 0$  a.e. on  $J_t^{-1}(\mathbf{R}_+^*)$  while  $\|\rho^{in}\|_{L^1(\mathbf{R}^N)} > 0$ , one has  $(\rho^{in} \mathcal{L}^N) \perp (J_t \mathcal{L}^N)$ , so that the projected measure  $\rho(t) = \Pi \# \mu(t)$  satisfies  $\rho(t) \perp \mathcal{L}^N$  by statement b) in Theorem 5.1. By statement d) in Theorem 6.1, the measure  $\rho(t)$  is diffuse if and only if

$$(\rho^{in} \mathcal{L}^N)(F_t^{-1}(\{x\}) \cap J_t^{-1}(\{0\})) = 0 \text{ for all } x \in C_t.$$

This may indeed happen, as shown by the following example.

**Example 5.** Assume  $N = 1$ , and set  $H(x, \xi) = \frac{1}{2}\xi^2$ , which generates the free flow  $\Phi_t : (x, \xi) \mapsto (x + t\xi, \xi)$ . By regularity of the Lebesgue measure, there exists a compact set  $K \subset (0, 1) \setminus \mathbf{Q}$  such that  $\frac{1}{2} < \mathcal{L}^1(K) < 1$ . Let  $\Omega = (0, 1) \setminus K$ ; since  $\Omega$  is open in  $(0, 1)$  and contains  $(0, 1) \cap \mathbf{Q}$ , it is a countably infinite union of disjoint nonempty open intervals:

$$\Omega = \bigcup_{n \in \mathbf{N}} I_n, \quad \text{so that } \mathcal{L}^1(\Omega) = \sum_{n \in \mathbf{N}} \mathcal{L}^1(I_n).$$

In particular  $\lambda := \mathcal{L}^1(\Omega) > 0$  since  $\mathcal{L}^1(I_n) > 0$  for each  $n \in \mathbf{N}$  (indeed each  $I_n$  is an open interval that contains at least one rational). Set  $t = 1$  and

$$F_1(y) := \begin{cases} y, & \text{if } y \leq 0, \\ \int_0^y \mathbf{1}_\Omega(z) dz, & \text{if } y \in (0, 1), \\ \lambda + (y - 1), & \text{if } y \geq 1. \end{cases}$$

Define

$$U^{in}(y) := F_1(y) - y.$$

The function  $F_1$  is Lipschitz continuous on  $\mathbf{R}$ , being the indefinite integral of a bounded measurable function. Therefore  $U^{in}$  is also Lipschitz continuous on  $\mathbf{R}$ .

The function  $F_1$  is increasing on  $\mathbf{R}$  — this being obvious on  $\mathbf{R}_-^*$  and on  $(1, \infty)$ . Indeed, if  $0 < y_1 < y_2 < 1$ , the interval  $(y_1, y_2)$  contains at least one rational number  $r$ , so that

$$F_1(y_2) - F_1(y_1) = \mathcal{L}^1(\Omega \cap (y_1, y_2)) > 0,$$

since the open set  $\Omega \cap (y_1, y_2)$  contains the intersection of the connected component of  $r$  in  $\Omega$  with  $(y_1, y_2)$ , which is a nonempty open interval. Since  $F_1(y) \rightarrow \pm\infty$  as  $y \rightarrow \pm\infty$ , we conclude that  $F_1 : \mathbf{R} \rightarrow \mathbf{R}$  is one-to-one and onto.

By the Lebesgue differentiation theorem (Theorem 7.11 in [20]),  $F_1$  is differentiable a.e. on  $\mathbf{R}$  (in fact it is even of class  $C^\infty$  on  $\mathbf{R}_-^* \cup (1, \infty)$ ) and  $F_1'(y) = \mathbf{1}_\Omega(y)$  for a.e.  $y \in (0, 1)$ . Thus, denoting by  $E$  the  $\mathcal{L}^1$ -negligible set on which  $F_1$  is not differentiable, one has  $(F_1')^{-1}(\{0\}) \cup E = K$ . Since  $F_1$  is one-to-one and onto,  $C_1 = F_1(K)$ .

Let  $\rho^{in} = \mathbf{1}_K$ ; thus  $\|\rho^{in}\|_{L^1} = \mathcal{L}^1(K) > 0$ , so that  $(\rho^{in} \mathcal{L}^1) \perp (|F'_1| \mathcal{L}^1)$ . The measure  $\rho(t) = \Pi \# \mu(t)$  where  $\mu(t) = \Phi \# \mu^{in}$  with  $\mu^{in} := \mathcal{L}_x^1 \otimes \rho^{in}(x) \delta_{U^{in}(x)}$  satisfies  $\rho(1) \perp \mathcal{L}^N$  by statement b) in Theorem 5.1

$$\rho(1)(\mathbf{R}) = \rho(1)(C_1) = \|\rho^{in}\|_{L^1}^2 = \mathcal{L}^1(K) > 0.$$

On the other hand, since  $F_1$  is one-to-one and onto, for each  $x \in \mathbf{R}$ , one has  $\#F_1^{-1}(\{x\}) = 1$ . In particular, for each  $x \in C_1 = F_1(K)$ , one has

$$F_1^{-1}(\{x\}) \subset (F'_1)^{-1}(\{0\}) \quad \text{so that } \#(F_1^{-1}(\{x\}) \cap (F'_1)^{-1}(\{0\})) = \#F_1^{-1}(\{x\}) = 1.$$

In particular, for each  $x \in C_1$ , by statement d) of Theorem 6.1, one has

$$(\rho^{in} \mathcal{L}^1)(F_1^{-1}(\{x\}) \cap (F'_1)^{-1}(\{0\})) = 0,$$

so that

$$\rho(1)(\{x\}) = 0.$$

Hence  $\rho(1) \perp \mathcal{L}^1$  and is diffuse, while  $\rho(1) \neq 0$  since  $\rho(1)(C_1) > 0$ .

## 7. APPLICATION TO THE CLASSICAL LIMIT OF QUANTUM MECHANICS

In this section, we apply the results obtained above to the classical limit of the Schrödinger equation.

**7.1. The classical scaling.** Consider the evolution Schrödinger equation

$$i\hbar \partial_t \psi = -\frac{1}{2} m \hbar^2 \Delta_x \psi + V(x) \psi$$

for the wave function  $\psi$  of a point particle of mass  $m$  subject to the action of an external potential  $V \equiv V(x) \in \mathbf{R}$ .

Choosing “appropriate” units of time  $T$  and length  $L$ , we recast the Schrödinger equation in terms of dimensionless variables  $\hat{t} := t/T$  and  $\hat{x} := x/L$ . We define a rescaled wave function  $\hat{\psi}$  and a rescaled, dimensionless potential  $\hat{V}$  by the formulas

$$\hat{\psi}(\hat{t}, \hat{x}) := \psi(t, x) \quad \text{and} \quad \hat{V}(\hat{x}) := \frac{T^2}{mL^2} V(x).$$

In these dimensionless variables, the Schrödinger equation takes the form

$$i \partial_{\hat{t}} \hat{\psi} = -\frac{\hbar T}{2mL^2} \Delta_{\hat{x}} \hat{\psi} + \frac{mL^2}{\hbar T} \hat{V}(\hat{x}) \hat{\psi}.$$

The dimensionless number  $2\pi\hbar T/mL^2$  is the ratio of the Planck constant to  $mL^2/T$ , that is (twice) the action of a classical particle of mass  $m$  moving at speed  $L/T$  on a distance  $L$ . If the scales of time  $T$  and length  $L$  have been chosen conveniently,  $L/T$  is the typical order of magnitude of the particle speed, and  $L$  is the typical length scale on which the particle motion is observed. The classical limit of quantum mechanics is defined by the scaling assumption  $2\pi\hbar \ll mL^2/T$  — i.e. the typical action of the particle considered is large compared to  $\hbar$ . Equivalently,  $mL/T$  is the order of magnitude of the particle momentum, so that  $2\pi\hbar T/mL$  is its de Broglie wavelength; the scaling assumption  $2\pi\hbar T/mL \ll L$  means that the de Broglie wavelength of the particle under consideration is small compared to the observation length scale  $L$ .

Introducing the small, dimensionless parameter  $\epsilon = \hbar T/mL^2$  and dropping hats in the dimensionless variables as well as on the rescaled wave function and dimensionless potential, we arrive at the following formulation for the Schrödinger equation in dimensionless variables

$$(40) \quad i\epsilon \partial_t \psi = -\frac{1}{2}\epsilon^2 \Delta_x \psi + V(x)\psi.$$

The WKB ansatz postulates that, at time  $t = 0$ , the wave function is of the form

$$\psi(t, x) = a^{in}(x) e^{iS^{in}(x)/\hbar}, \quad x \in \mathbf{R}^N.$$

Consistently with the scaling argument above, we set

$$\hat{a}^{in}(\hat{x}) := a^{in}(x) \text{ and } \hat{S}^{in}(\hat{x}) := TS^{in}(x)/mL^2$$

— since  $S^{in}$  has the dimension of an action — so that

$$(41) \quad \hat{\psi}(0, \hat{x}) = \hat{a}^{in}(\hat{x}) e^{i\hat{S}^{in}(\hat{x})/\epsilon}.$$

Dropping hats in the initial data as well as in the Schrödinger equation, one arrives at the following Cauchy problem for the Schrödinger equation in dimensionless variables:

$$(42) \quad \begin{cases} i\epsilon \partial_t \psi_\epsilon = -\frac{1}{2}\epsilon^2 \Delta_x \psi_\epsilon + V(x)\psi_\epsilon, & x \in \mathbf{R}^N, t \in \mathbf{R}, \\ \psi_\epsilon(0, x) = a^{in}(x) e^{iS^{in}(x)/\epsilon}. \end{cases}$$

The problem of the classical limit of the Schrödinger equation is to describe the wave function  $\psi_\epsilon$  in the limit as  $\epsilon \rightarrow 0^+$ .

**7.2. The WKB method.** The traditional procedure for describing the classical limit of the Schrödinger equation is the WKB method recalled below. First we recall some mathematical tools and elements of notation used in the presentation of that method.

Assume that  $V \in C^\infty(\mathbf{R}^N)$  satisfies

$$(43) \quad \partial^\alpha V \in L^\infty(\mathbf{R}^N) \text{ for each multi-index } \alpha \in \mathbf{N}^N$$

and

$$(44) \quad \frac{V(x)}{|x|} \rightarrow 0 \text{ as } |x| \rightarrow +\infty.$$

Then the Hamiltonian

$$H(x, \xi) := \frac{1}{2}|\xi|^2 + V(x)$$

satisfies (7) and therefore generates a global flow

$$\mathbf{R}^N \times \mathbf{R}^N \ni (x, \xi) \mapsto (X_t(x, \xi), \Xi_t(x, \xi)) \in \mathbf{R}^N \times \mathbf{R}^N$$

by Lemma 2.1.

Assume further that

$$(45) \quad \sup_{x \in \mathbf{R}^N} \int_{\mathbf{R}^N} \Gamma_\eta(x-y) V^-(y) dy \rightarrow 0 \text{ as } \eta \rightarrow 0 \quad \text{if } N \geq 2$$

with

$$\Gamma_\eta(z) = \begin{cases} \mathbf{1}_{[0, \eta]}(|z|) |z|^{2-N} & \text{if } N \geq 3, \\ \mathbf{1}_{[0, \eta]}(|z|) \ln(1/|z|) & \text{if } N = 2, \end{cases}$$



while

$$(46) \quad \sup_{x \in \mathbf{R}^N} \int_{x-1}^{x+1} V^-(y) dy < \infty \quad \text{if } N = 1.$$

Under assumptions (45)-(46), the operator  $-\frac{1}{2}\epsilon^2 \Delta_x + V$  has a self-adjoint extension on  $L^2(\mathbf{R}^N)$  that is bounded from below.

Under assumption (43), there exists a FIO that is a parametrix for the operator

$$G_\epsilon(t) := e^{i\frac{t}{\epsilon}(\frac{1}{2}\epsilon^2 \Delta_x - V)}$$

see for instance Theorem 2.1 in [16], whose main features are recalled below.

Consider the action

$$(47) \quad S(t, x, \xi) := \int_0^t \left( \frac{1}{2} |\Xi_s(x, \xi)|^2 - V(X_s(x, \xi)) \right) ds$$

Given  $T > 0$ , we shall have to deal with the class of phase functions

$$\varphi \equiv \varphi(t, x, y, \eta) \in C \text{ of class } C^\infty \text{ on } [0, T] \times \mathbf{R}^N \times \mathbf{R}^N \times \mathbf{R}^N$$

satisfying the conditions

$$(48) \quad \begin{cases} \varphi(t, X_t(y, \eta), y, \eta) = S(t, y, \eta), \\ D_x \varphi(t, X_t(y, \eta), y, \eta) = \Xi_t(y, \eta), \\ iD_x^2 \varphi(t, x, y, \eta) \leq 0 \text{ is independent of } x, \\ \det(D_{x\eta} \varphi(t, X_t(y, \eta), y, \eta)) \neq 0 \text{ for each } (t, y, \eta) \in [0, T] \times \mathbf{R}^N \times \mathbf{R}^N. \end{cases}$$

Pick  $\chi \in C_c^\infty(\mathbf{R}^N \times \mathbf{R}^N)$  and  $T > 0$ . Then, for any phase function  $\varphi$  satisfying (48) and any  $n \geq 0$ , there exists  $A_n \equiv A_n(t, y, \eta, \epsilon) \in C_c^\infty([0, T] \times \mathbf{R}^N \times \mathbf{R}^N)[\epsilon]$  such that the FIO  $G_n(t)$  with Schwartz kernel

$$G_{\epsilon, n}(t, x, y) = \int A_n(t, y, \eta, \epsilon) e^{i\varphi(t, x, y, \eta)/\epsilon} \frac{d\eta}{(2\pi\epsilon)^N}$$

satisfies

$$\sup_{0 \leq t \leq T} \|(G_\epsilon(t) - G_{\epsilon, n}(t))\chi(x, -i\epsilon\partial_x)\|_{\mathcal{L}(L^2(\mathbf{R}^N))} \leq C[V, T, \chi] \epsilon^{n-2N}.$$

In this inequality the notation  $\chi(x, -i\epsilon\partial_x)$  designates the pseudo-differential operator defined by the formula

$$\chi(x, -i\epsilon\partial_x)\phi(x) := \iint_{\mathbf{R}^N \times \mathbf{R}^N} e^{i(x-y)\cdot\eta/\epsilon} \chi(x, \eta) \phi(y) \frac{dy d\eta}{(2\pi\epsilon)^N}$$

Taking Theorem 2.1 in [16] for granted, one arrives at the following description of the classical limit of (42). It is stated without proof in Appendix 11 of [5] or as Theorem 5.1 in [3].

Let  $U^{in} = \nabla S^{in}$  and let  $C$  be defined as in (15); let  $\mathcal{N}(t, x)$  and  $y_j(t, x)$  be defined as in Proposition 3.3 for each  $(t, x) \in \mathbf{R} \times \mathbf{R}^N \setminus C$ . Let  $J_t(y)$  be defined as in (14).

**Proposition 7.1.** *Let  $a^{in} \in C_c^m(\mathbf{R}^N)$  and  $S^{in} \in C^{m+1}(\mathbf{R}^N)$  with  $m > 6N + 4$ . For all  $\epsilon > 0$  and all  $(t, x) \in \mathbf{R}_+ \times \mathbf{R}^N \setminus C$ , set*

$$(49) \quad \Psi_\epsilon(t, x) = \sum_{j=1}^{\mathcal{N}(t, x)} \frac{a^{in}(y_j(t, x))}{J_t(y_j(t, x))^{1/2}} e^{iS_j(t, x)/\epsilon} e^{i\pi\nu_j(t, x)/2},$$

where

$$S_j(t, x) := S^{in}(y_j(t, x)) + S(t, y_j(t, x), \nabla S^{in}(y_j(t, x))), \quad j = 1, \dots, \mathcal{N}(t, x)$$

$S(t, y, \xi)$  is given by (47) and  $\nu_j(t, x) \in \mathbf{Z}$  for all  $(t, x) \in \mathbf{R}_+ \times \mathbf{R}^N \setminus C$  is constant on each connected component of  $\mathbf{R} \times \mathbf{R}^N$  where  $j \leq \mathcal{N}$ .

Then the solution  $\psi_\epsilon$  of the Cauchy problem (42) satisfies

$$(50) \quad \psi_\epsilon(t, x) = \Psi_\epsilon(t, x) + R_\epsilon^1(t, x) + R_\epsilon^2(t, x)$$

for all  $T > 0$ , where

$$\sup_{0 \leq t \leq T} \|R_\epsilon^1\|_{L^2(B(0, R))} = O(\epsilon) \text{ for all } R > 0$$

and

$$\sup_{(t, x) \in K} |R_\epsilon^2(t, x)| = O(\epsilon) \text{ for each compact } K \subset \mathbf{R}_+ \times \mathbf{R}^N \setminus C$$

as  $\epsilon \rightarrow 0^+$ .

Let us remark that the integer  $\nu_j(t, x)$  is precisely defined through the proof of the Proposition, see formula 63 below. As we won't need it we will not discuss here its link with the Morse index of the path

$$[0, t] \ni s \mapsto X_s(y_j(t, x), \nabla S^{in}(y_j(t, x))),$$

see [3] for more details concerning this point.

A self-contained proof of Proposition 7.1 is given in Appendix A following the parametrix construction of [16].

**7.3. The Wigner transform.** Alternately, the asymptotic limit of  $\psi_\epsilon$  can also be investigated with the help of Wigner's transform [23, 17].

For each  $\Psi \in L^2(\mathbf{R}^N)$ , one defines the Wigner transform of  $\Psi$  at scale  $\epsilon$  by the formula

$$W_\epsilon[\Psi](x, \xi) := \frac{1}{(2\pi)^N} \int_{\mathbf{R}^N} e^{-iy \cdot \xi} \Psi(x + \frac{1}{2}\epsilon y) \overline{\Psi(x - \frac{1}{2}\epsilon y)} dy.$$

Let us define the  $\epsilon$ -Fourier transform  $\widehat{\Psi}$  as

$$\widehat{\Psi}(\xi) := \frac{1}{(2\pi\epsilon)^{N/2}} \int_{L^2(\mathbf{R}^N)} \Psi(x) e^{i\frac{x\xi}{\epsilon}} dx.$$

Straightforward computations lead easily to the following identities.

$$(51) \quad \int_{\mathbf{R}^N} W_\epsilon[\Psi](x, \xi) d\xi = |\Psi(x)|^2, \quad \int_{\mathbf{R}^N} W_\epsilon[\Psi](x, \xi) dx = |\widehat{\Psi}(\xi)|^2$$

and, for any pseudodifferential  $A$  operator of (total) Weyl symbol  $a(x, \xi)$

$$(52) \quad \int_{\mathbf{R}^N} W_\epsilon[\Psi](x, \xi) a(x, \xi) dx d\xi = \langle \Psi, A\Psi \rangle_{L^2(\mathbf{R}^N)}$$

**Lemma 7.2.** *Let  $a^{in} \in L^2(\mathbf{R}^N)$  and  $S^{in} \in W_{loc}^{1,1}(\mathbf{R}^N)$ . Then*

$$W_\epsilon[a^{in} e^{iS^{in}/\epsilon}](x, \xi) \rightarrow (a^{in})^2(x) \delta(\xi - \nabla S^{in}(x)) \text{ in } \mathcal{S}'(\mathbf{R}^N \times \mathbf{R}^N)$$

as  $\epsilon \rightarrow 0$ .

See Example III.5 in [17].

**Proposition 7.3** ([17]). *Assume that  $V \in C^{1,1}(\mathbf{R}^N)$  satisfies (45)-(46)-(43)-(44). Let*

$$\mu_\epsilon(t, x, \xi) := W_\epsilon[\psi_\epsilon(t, \cdot)](x, \xi)$$

for each  $\epsilon > 0$  and each  $t \in \mathbf{R}$ , and for a.e.  $(x, \xi) \in \mathbf{R}^N \times \mathbf{R}^N$ .

Then the family  $\mu_\epsilon$  converges in  $\mathcal{S}'(\mathbf{R} \times \mathbf{R}^N \times \mathbf{R}^N)$  as  $\epsilon \rightarrow 0$  to the unique solution of the Liouville equation

$$(53) \quad \begin{cases} \partial_t \mu + \xi \cdot \nabla_x \mu - \nabla_x V(x) \cdot \nabla_\xi \mu = 0, & x, \xi \in \mathbf{R}^N, t \in \mathbf{R}, \\ \mu(0, x, \xi) := a^{in}(x)^2 \delta_{\nabla S^{in}(x)}(\xi). \end{cases}$$

In particular,  $\mu \in C_b(\mathbf{R}, w - \mathcal{M}(\mathbf{R}^N \times \mathbf{R}^N))$  and  $\mu(t)$  is a positive Radon measure for all  $t \in \mathbf{R}$ .

This proposition is stated as Theorem IV.1 in [17], to which we refer for a complete proof.

Therefore, the structure of the measure  $\mu(t)$  is given by Theorem 5.1 with

$$\rho^{in} = (a^{in})^2 \quad \text{and} \quad U^{in} = \nabla S^{in}.$$

Likewise, Theorem 6.1 provides a precise description of the exceptional sets associated to  $\mu(t)$ .

**7.4. Link with the results of the present article.** The WKB asymptotic solution (49) obviously contains more information than the regular part  $\mu_a(t)$  of the Wigner measure  $\mu(t)$  that is the solution of the Liouville equation (53) — for the definition of  $\mu_a(t)$  and its structure, see Theorem 5.1.

For instance, the (finitely many) phase functions  $S_j(t, x)$  appearing in (49) define velocity fields  $U_j(t, x) := \nabla_x S_j(t, x)$  as in the formula giving  $\mu_a(t)$  in Theorem 5.1. On the contrary, the Morse indices  $\nu_j(t, x)$  do not appear in the formula for  $\mu_a(t)$ .

One can go a little further, and compute the Wigner measure associated to the WKB asymptotic solution (49), under the assumptions used in Proposition 7.1.

Let  $t \in \mathbf{R}$ ; for each  $x \in \mathbf{R}^N \setminus C_t$  and  $j \in \{1, \dots, \mathcal{N}(t, x)\}$ , one has

$$\begin{aligned} \nabla_x S_j(t, x) &= DS^{in}(y_j(t, x)) D_x y_j(t, x) + D_y S(t, y_j(t, x), DS^{in}(y_j(t, x))) D_x y_j(t, x) \\ &\quad + D_\eta S(t, y_j(t, x), DS^{in}(y_j(t, x))) D^2 S^{in}(y_j(t, x)) D_x y_j(t, x) \\ &= \Xi_t(y_j(t, x), DS^{in}(y_j(t, x))) \cdot (D_y X_t(y_j(t, x), DS^{in}(y_j(t, x))) D_x y_j(t, x) \\ &\quad + D_\eta X_t(y_j(t, x), DS^{in}(y_j(t, x))) D^2 S^{in}(y_j(t, x)) D_x y_j(t, x)), \end{aligned}$$

in view of formulas (3.1-2) in [16] recalled above (see (58)). By definition

$$(54) \quad X_t(y_j(t, x), DS^{in}(y_j(t, x))) = x, \quad j = 1, \dots, \mathcal{N}(t, x),$$

so that

$$\begin{aligned} &D_y X_t(y_j(t, x), DS^{in}(y_j(t, x))) D_x y_j(t, x) \\ &+ D_\eta X_t(y_j(t, x), DS^{in}(y_j(t, x))) D^2 S^{in}(y_j(t, x)) D_x y_j(t, x) = I \end{aligned}$$

which implies in turn that

$$(55) \quad \nabla_x S_j(t, x) = \Xi_t(y_j(t, x), DS^{in}(y_j(t, x))), \quad j = 1, \dots, \mathcal{N}(t, x).$$

Because of (54) and of the condition

$$y_j(t, x) \neq y_k(t, x) \quad \text{if } 1 \leq j < k \leq \mathcal{N}(t, x),$$

one has

$$\begin{aligned}\nabla_x S_j(t, x) &= \Xi_t(y_j(t, x), DS^{in}(y_j(t, x))) \\ &\neq \Xi_t(y_k(t, x), DS^{in}(y_k(t, x))) = \nabla_x S_k(t, x) \quad \text{if } 1 \leq j < k \leq \mathcal{N}(t, x),\end{aligned}$$

by uniqueness of the solution of the Cauchy problem for Hamilton's equations.

Applying Proposition 1.5 in [9] implies that

$$W_\epsilon[\Psi_\epsilon(t, \cdot)] \rightarrow \mu(t) \quad \text{in } \mathcal{D}'((\mathbf{R}^N \setminus C_t) \times \mathbf{R}^N)$$

as  $\epsilon \rightarrow 0$ , where

$$\mu(t, x, d\xi) := \sum_{j=1}^{\mathcal{N}(t, x)} \frac{|a^{in}(y_j(t, x))|^2}{J_t(y_j(t, x))} \delta_{\nabla_x S_j(t, x)}.$$

The expression in the right hand side of the equality above coincides with the regular part  $\mu_a(t)$  of the Wigner measure  $\mu(t)$  in Theorem 5.1.

Since Theorem 5.1 is valid under much weaker hypothesis on the phase of the initial WKB type initial condition, one might ask alternatively which information our results give on the solution of the Schrödinger equation itself. The fact that the Wigner measure is the limit of the Wigner transform at  $\epsilon = 0$  shows clearly that there is no simple way of pulling back any quantitative information from it to the wave function at  $\epsilon \neq 0$ . Moreover the Wigner transform is quadratic. Therefore it is natural to expect that our results carry information about the limit  $\epsilon \rightarrow 0$  of some quadratic expressions of the solution of the Schrödinger equation with rough initial data.

Using (51) and (52) we easily get the

**Proposition 7.4.** *Let  $\psi_\epsilon$  the solution of the Schrödinger equation (42) where  $a^{in}$  and  $\nabla S^{in}$  satisfy the hypothesis of Theorem 5.1. One has, for each  $t \in \mathbf{R}$ ,*

a) *for any pseudodifferential operator  $A$  of principal Weyl symbol  $a(x, \xi)$ ,*

$$\lim_{\epsilon \rightarrow 0} \langle \psi_\epsilon, A\psi_\epsilon \rangle_{L^2(\mathbf{R}^N)} = \int a\mu(t),$$

where  $\mu(t)$  is described in Theorem 5.1,

b) *weakly in  $\mathcal{M}((\mathbf{R}^N \setminus C'_t))$ ,*

$$|\psi_\epsilon(x)|^2 \rightarrow \sum_{y \in F_t^{-1}(\{x\})} \frac{|a^{in}|^2 \mathbf{1}_{P_t}(y)}{J_t} \quad \text{as } \epsilon \rightarrow 0,$$

c) *weakly in  $\mathcal{M}(\mathbf{R}^N \setminus \Xi_t(C'_t, \nabla S^{in}(C'_t)))$ ,*

$$|\widehat{\psi}_\epsilon(\xi)|^2 \rightarrow \int_{\mathbf{R}^N} \sum_{y \in F_t^{-1}(\{x\})} \frac{|a^{in}|^2 \mathbf{1}_{P_t}(y)}{J_t} \delta(\xi - \Xi_t(y, \nabla S^{in}(y))) dx \quad \text{as } \epsilon \rightarrow 0,$$

where again  $C'_t$  is defined in (26) and  $P_t$ ,  $J_t$ ,  $\Xi_t$  and  $F_t$  are as in Theorem 5.1,  $\mathcal{M}(\mathbb{X})$  denotes the cone of positive bounded measures on  $\mathbb{X}$  and  $\Xi_t(C'_t, \nabla S^{in}(C'_t))$  is meant for  $\{\Xi_t(y, \nabla S^{in}(y)), y \in C'_t\}$ .

Note that the restrictions  $x \notin C'_t$  in b) and  $\xi \notin \Xi_t(C'_t, \nabla S^{in}(C'_t))$  in c) can be released using the singular part of  $\mu(t)$  introduced in Theorem 5.1. We will not do it here, the corresponding expressions being somehow less explicit.

Let us give some final comments concerning the results contained in the present paper. Though Proposition 7.1 can be improved by adding extra hypothesis on  $a^{in}$  and  $S^{in}$ , the approximation stated *in full generality* holds in  $\mathbf{R} \times \mathbf{R}^N \setminus C$ , and only locally. In particular, the approximation of the wave function  $\psi_\epsilon$  by the WKB ansatz  $\Psi_\epsilon$  is *not uniform* as  $(t, x)$  gets near  $C$ . (For more on this well known fact in another, related context, we refer the reader to the last paragraph in §55 and on the problem in §59 in [14].) On the contrary, the result obtained in Theorem 5.1 is *global*, holds in the sense of measures and does not involve the caustic  $C$  as explained in the previous sections.

#### APPENDIX A. PROOF OF PROPOSITION 7.1

Let  $\chi(x, \xi) = \chi_1(x)\chi_2(\xi)$  with  $\chi_1, \chi_2 \in C_c^\infty(\mathbf{R}^N)$ , satisfying

$$\mathbf{1}_{B(0,R)}(x) \leq \chi_1(x) \leq \mathbf{1}_{B(0,R+1)} \text{ and } \mathbf{1}_{B(0,Q)}(\xi) \leq \chi_2(\xi) \leq \mathbf{1}_{B(0,Q+1)}$$

for all  $x, \xi \in \mathbf{R}^N$ , where  $R > 0$  and  $Q$  is to be chosen later. Pick  $n > 2N$ ; then

$$\begin{aligned} \psi_\epsilon(t, \cdot) - G_{\epsilon,n}(t)\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in} &= G_\epsilon(t)(1 - \chi(x, -i\epsilon\partial_x))\psi_\epsilon^{in} \\ &\quad + (G_\epsilon(t) - G_{\epsilon,n}(t))\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in}. \end{aligned}$$

Since  $G_\epsilon(t)$  is a unitary group on  $L^2(\mathbf{R}^N)$

$$\begin{aligned} \|\psi_\epsilon(t, \cdot) - G_{\epsilon,n}(t)\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} &\leq \|(1 - \chi(x, -i\epsilon\partial_x))\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} \\ &\quad + \|(G_\epsilon(t) - G_{\epsilon,n}(t))\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} \\ &\leq \|(1 - \chi(x, -i\epsilon\partial_x))\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} \\ &\quad + C_{T,Q}\epsilon^{n-2N}\|a^{in}\|_{L^2(\mathbf{R}^N)} \end{aligned}$$

for all  $t \in [0, T]$ , where  $C_{Q,T} = C[V, T, \chi]$ .

Now,  $\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in} = \chi_1(x)\chi_2((-i\epsilon\partial_x)\psi_\epsilon^{in})$  and since  $\text{supp}(a^{in}) \subset B(0, R)$

$$\begin{aligned} \|(1 - \chi(x, -i\epsilon\partial_x))\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} &= \|\chi_1(1 - \chi_2(-i\epsilon\partial_x))\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} \\ &\leq \|(1 - \chi_2(-i\epsilon\partial_x))\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} \\ &= (2\pi)^{-N} \|(1 - \chi_2(\epsilon\xi))\widehat{\psi_\epsilon^{in}}\|_{L^2(\mathbf{R}^N)} \\ &\leq (2\pi)^{-N} \|\mathbf{1}_{[Q/\epsilon, \infty)}(|\xi|)\widehat{\psi_\epsilon^{in}}\|_{L^2(\mathbf{R}^N)} \end{aligned}$$

Since

$$\widehat{\psi_\epsilon^{in}}(\zeta/\epsilon) = \int_{\mathbf{R}^N} e^{-i(\zeta \cdot x - S^{in}(x))/\epsilon} a^{in}(x) dx$$

we conclude from estimate (7.7.1') in [11] that

$$|\widehat{\psi_\epsilon^{in}}(\zeta/\epsilon)| \leq \frac{C\|a^{in}\|_{W^{m,\infty}(\mathbf{R}^N)}}{(|\zeta| - \|\nabla S^{in}\|_{L^\infty(B(0,R))})^m} \epsilon^m$$

provided that  $\text{supp}(a^{in}) \subset B(0, R)$  and  $|\zeta| > 1 + \|\nabla S^{in}\|_{L^\infty(B(0,R))}$ . Therefore

$$(56) \quad \begin{aligned} \|\psi_\epsilon(t, \cdot) - G_{\epsilon,n}(t)\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in}\|_{L^2(\mathbf{R}^N)} &\leq C_{T,Q}\|a^{in}\|_{L^2(\mathbf{R}^N)}\epsilon^{n-2N} \\ &\quad + C\|(1 + |\zeta|)^{-m}\|_{L^2(\mathbf{R}^N)}\|a^{in}\|_{W^{m,\infty}(\mathbf{R}^N)}\epsilon^m \end{aligned}$$

for all  $m > N/2$ .

Next we analyze the term

$$(57) \quad \begin{aligned} & G_{\epsilon,n}(t)\chi(x, -i\epsilon\partial_x)\psi_\epsilon^{in}(x) \\ &= \iiint A_n(t, y, \eta, \epsilon)a^{in}(z)\chi_2(\zeta)e^{i(\varphi(t,x,y,\eta)+\zeta\cdot(y-z)+S^{in}(z))/\epsilon}\frac{dzd\zeta dyd\eta}{(2\pi\epsilon)^{2N}} \end{aligned}$$

with the stationary phase method.

Choose  $\varphi$  of the form

$$\varphi(t, x, y, \eta) = S(t, y, \eta) + (x - X_t(y, \eta)) \cdot \Xi_t(y, \eta) + iB : (x - X_t(y, \eta))^{\otimes 2}$$

where the matrix  $B = B^T > 0$  is constant (see formula (2.7) [16] and the following Remark 2.1). Critical points of the phase in the oscillating integral (57) are defined by the system of equations

$$\begin{cases} -\zeta + DS^{in}(z) = 0, \\ y - z = 0, \\ \partial_y S(t, x, y) - \Xi_t(y, \eta) \cdot D_y X_t(y, \eta) + (x - X_t(y, \eta)) \cdot D_y \Xi_t(y, \eta) \\ - iB : (x - X_t(y, \eta)) \otimes D_y X_t(y, \eta) + \zeta = 0, \\ \partial_\eta S(t, x, y) - \Xi_t(y, \eta) \cdot D_\eta X_t(y, \eta) + (x - X_t(y, \eta)) \cdot D_\eta \Xi_t(y, \eta) \\ - iB : (x - X_t(y, \eta)) \otimes D_\eta X_t(y, \eta) + \zeta = 0. \end{cases}$$

At this point, we recall formulas (3.1-2) from [16]

$$(58) \quad \begin{aligned} \partial_y S(t, x, y) &= \Xi_t(y, \eta) \cdot D_y X_t(y, \eta) - \eta, \\ \partial_\eta S(t, x, y) &= \Xi_t(y, \eta) \cdot D_\eta X_t(y, \eta), \end{aligned}$$

together with the following definitions

$$\begin{aligned} Y(t, y, \eta) &:= D_y \Xi_t(y, \eta) - iB D_y X_t(y, \eta), \\ Z(t, y, \eta) &:= D_\eta \Xi_t(y, \eta) - iB D_\eta X_t(y, \eta). \end{aligned}$$

Thus the critical points of the phase in (57) are given by

$$\begin{cases} \zeta = DS^{in}(z), \\ y = z, \\ (x - X_t(y, \eta))^T Y(t, y, \eta) + \zeta = \eta, \\ (x - X_t(y, \eta))^T Z(t, y, \eta) = 0. \end{cases}$$

Since the matrix  $Z$  is invertible by Lemma 4.1 of [16], we conclude that the system of equations above is equivalent to

$$\begin{cases} \zeta = DS^{in}(z), \\ y = z, \\ \zeta = \eta, \\ x = X_t(y, \eta). \end{cases}$$

In other words,

$$F_t(z) = x, \quad y = z, \quad \zeta = \eta = DS^{in}(z).$$

Assuming that  $(t, x) \notin C$ , we apply Proposition 3.3 and conclude that the set of critical points of the phase in (57) is of the form

$$\begin{cases} y = z = y_j(t, x), \\ \zeta = \eta = DS^{in}(y_j(t, x)), \end{cases} \quad j = 1, \dots, \mathcal{N}(t, x).$$

At this point, we apply the stationary phase method (Theorem 7.7.5 in [11]). First we need to compute the Hessian of the phase in (57) at its critical points. One finds

$$H_j(t, x) := \begin{pmatrix} D^2S^{in} & -I & 0 & 0 \\ -I & 0 & +I & 0 \\ 0 & +I & -YD_yX_t & -YD_\eta X_t - I \\ 0 & 0 & -ZD_yX_t & -ZD_\eta X_t \end{pmatrix}_{\substack{y=y_j(t,x) \\ \eta=DS^{in}(y_j(t,x))}}$$

and it remains to compute  $\det(H_j(t, x))$ . Adding the first row of  $H_j(t, x)$  to the third row, one finds that

$$\begin{aligned} \det H_j(t, x) &= \begin{vmatrix} D^2S^{in} & -I & 0 & 0 \\ -I & 0 & +I & 0 \\ D^2S^{in} & 0 & -YD_yX_t & -YD_\eta X_t - I \\ 0 & 0 & -ZD_yX_t & -ZD_\eta X_t \end{vmatrix}_{\substack{y=y_j(t,x) \\ \eta=DS^{in}(y_j(t,x))}} \\ &= (-1)^N \begin{vmatrix} -I & +I & 0 & 0 \\ D^2S^{in} & -YD_yX_t & -YD_\eta X_t - I & 0 \\ 0 & -ZD_yX_t & -ZD_\eta X_t & 0 \end{vmatrix}_{\substack{y=y_j(t,x) \\ \eta=DS^{in}(y_j(t,x))}} \\ &= (-1)^N \begin{vmatrix} -I & 0 & 0 & 0 \\ D^2S^{in} & -YD_yX_t + D^2S^{in} & -YD_\eta X_t - I & 0 \\ 0 & -ZD_yX_t & -ZD_\eta X_t & 0 \end{vmatrix}_{\substack{y=y_j(t,x) \\ \eta=DS^{in}(y_j(t,x))}} \end{aligned}$$

where the last equality follows from adding the first column in the right hand side of the second equality to the second column. Eventually, one finds that

$$\det H_j(t, x) = \begin{vmatrix} -YD_yX_t + D^2S^{in} & -YD_\eta X_t - I \\ -ZD_yX_t & -ZD_\eta X_t \end{vmatrix}_{\substack{y=y_j(t,x) \\ \eta=DS^{in}(y_j(t,x))}}$$

which is computed as follows. First

$$\begin{vmatrix} I & -YZ^{-1} \\ 0 & I \end{vmatrix} \begin{vmatrix} -YD_yX_t + D^2S^{in} & -YD_\eta X_t - I \\ -ZD_yX_t & -ZD_\eta X_t \end{vmatrix} = \begin{vmatrix} D^2S^{in} & -I \\ -ZD_yX_t & -ZD_\eta X_t \end{vmatrix}$$

so that

$$\det H_j(t, x) = \begin{vmatrix} D^2S^{in} & -I \\ -ZD_yX_t & -ZD_\eta X_t \end{vmatrix}_{\substack{y=y_j(t,x) \\ \eta=DS^{in}(y_j(t,x))}}$$

On the other hand

$$\begin{aligned} \begin{vmatrix} D^2S^{in} & -I \\ -ZD_yX_t & -ZD_\eta X_t \end{vmatrix} &= \det(ZD_yX_t + ZD_\eta X_t D^2S^{in}) \\ &= \det(Z) \det(D_yX_t + D_\eta X_t D^2S^{in}) \end{aligned}$$

by the following elementary lemma.

**Lemma A.1.** *Let  $A, B, C, D \in M_N(\mathbf{C})$ . If  $AB = BA$ , one has*

$$\begin{vmatrix} A & B \\ C & D \end{vmatrix} = \det(DA - CB).$$

Therefore

$$(59) \quad \begin{aligned} \det H_j(t, x) &= \det(Z) \det(D_y X_t + D_\eta X_t D^2 S^{in}) \Big|_{y=y_j(t, x), \eta=DS^{in}(y_j(t, x))} \\ &= \det(Z(y_j(t, x), DS^{in}(y_j(t, x)))) \det(DF_t(y_j(t, x))) \end{aligned}$$

where  $F_t$  is defined in (13) with  $U^{in} = DS^{in}$ .

Pick a nonempty closed ball  $B \subset \mathbf{R} \times \mathbf{R}^N \setminus C$ , let  $\mathcal{N}_B = \mathcal{N}(t, x)$  for all  $(t, x) \in B$ , and let

$$K_j = \{(y_j(t, x), \nabla S^{in}(y_j(t, x))) \mid (t, x) \in B\}, \quad j = 1, \dots, \mathcal{N}_B.$$

Assuming that  $B$  is of small enough radius,  $K_j \cap K_k = \emptyset$  for  $j \neq k \in \{1, \dots, \mathcal{N}_B\}$ . Let  $\kappa_j \in C_c^\infty(\mathbf{R}^{2N})$  for all  $j = 1, \dots, \mathcal{N}_B$ , such that

$$\begin{cases} \kappa_j \geq 0 \text{ and } \kappa_j|_{K_j} = 1, & j = 1, \dots, \mathcal{N}_B, \\ \text{while } \kappa_j \kappa_k = 0 \text{ for } j \neq k \in \{1, \dots, \mathcal{N}_B\}, \end{cases}$$

Applying Theorem 7.7.1 in [11] shows that

$$(60) \quad \begin{aligned} \sup_{(t, x) \in B} \left| \iiint \iiint A_n(t, y, \eta, \epsilon) a^{in}(z) \chi_2(\zeta) e^{i(\varphi(t, x, y, \eta) + \zeta \cdot (y - z) + S^{in}(z)) / \epsilon} \frac{dz d\zeta dy d\eta}{(2\pi\epsilon)^{2N}} \right. \\ \left. - \sum_{j=1}^{\mathcal{N}_B} \iiint \iiint A_n(t, y, \eta, \epsilon) a^{in}(z) \chi_2(\zeta) \kappa_j(y, \eta) \kappa_j(z, \zeta) \right. \\ \left. e^{i(\varphi(t, x, y, \eta) + \zeta \cdot (y - z) + S^{in}(z)) / \epsilon} \frac{dz d\zeta dy d\eta}{(2\pi\epsilon)^{2N}} \right| = O(\epsilon) \end{aligned}$$

as  $\epsilon \rightarrow 0$ .

Next we set

$$\begin{aligned} I_j(t, x, \epsilon) &:= \iiint \iiint A_n(t, y, \eta, \epsilon) a^{in}(z) \chi_2(\zeta) \kappa_j(y, \eta) \kappa_j(z, \zeta) \\ &\quad \times e^{i(\varphi(t, x, y, \eta) + \zeta \cdot (y - z) + S^{in}(z)) / \epsilon} \frac{dz d\zeta dy d\eta}{(2\pi\epsilon)^{2N}} \end{aligned}$$

for  $j = 1, \dots, \mathcal{N}_B$ . By Theorem 7.7.5 in [11], we conclude that

$$(61) \quad \begin{aligned} \sup_{(t, x) \in B} |I_j(t, x, \epsilon) - A_0(t, y_j(t, x), \nabla S^{in}(y_j(t, x)), 0) a^{in}(y_j(t, x)) \chi_2(\nabla S^{in}(y_j(t, x))) \\ \times e^{i(\varphi(t, x, y_j(t, x), \nabla S^{in}(y_j(t, x))) + S^{in}(y_j(t, x))) / \epsilon} (\det H_j(t, x))^{-1/2}| = O(\epsilon) \end{aligned}$$

as  $\epsilon \rightarrow 0$ . Our choice of  $\chi_2$  and  $\varphi$  implies that  $\chi_2(\nabla S^{in}(y_j(t, x))) = 1$  and

$$\varphi(t, x, y_j(t, x), \nabla S^{in}(y_j(t, x))) = S(t, y_j(t, x), \nabla S^{in}(y_j(t, x)))$$

so that

$$(\varphi(t, x, y_j(t, x), \nabla S^{in}(y_j(t, x))) + S^{in}(y_j(t, x))) = S_j(t, x).$$

By formula (2.13) in [16]

$$(62) \quad A_0(t, y_j(t, x), \nabla S^{in}(y_j(t, x)), 0) = \sqrt[cont]{\text{Det}(Z(y_j(t, x), DS^{in}(y_j(t, x))))}$$



where the determination of  ${}^{\text{cont}}\sqrt{z}$  is obtained by continuity (let us recall that  $\text{Det}(Z(y_j(t, x), DS^{\text{in}}(y_j(t, x)))$  never vanishes), According to (62), (59), Lemma 5.1 and formula (5.15) in [16], we have

$$(63) \quad \begin{aligned} & A_0(t, y_j(t, x), \nabla S^{\text{in}}(y_j(t, x)), 0)(\det H_j(t, x))^{-1/2} \\ &= |\det(DF_t(y_j(t, x)))|^{-1/2} e^{i\pi\nu_j(t, x)/2} = J(y_j(t, x))^{-1/2} e^{i\pi\nu_j(t, x)/2}, \end{aligned}$$

where  $\nu_j(t, x)$  is an integer.

Putting together (56)-(60)-(61)-(63) concludes the proof.

*Proof of Lemma A.1.* If  $A$  is nonsingular and  $AB = BA$ , one has

$$\begin{aligned} \begin{vmatrix} A & B \\ C & D \end{vmatrix} &= (-1)^N \begin{vmatrix} A & B \\ C & D \end{vmatrix} \begin{vmatrix} A^{-1} & B \\ 0 & -A \end{vmatrix} = (-1)^N \begin{vmatrix} I & 0 \\ CA^{-1} & CB - DA \end{vmatrix} \\ &= (-1)^N \det(CB - DA) = \det(DA - CB). \end{aligned}$$

Since both sides of the identity above are continuous functions of  $A$  and the set of nonsingular matrices  $GL_N(\mathbf{C})$  is dense in  $M_N(\mathbf{C})$ , this identity holds for all  $A \in M_n(\mathbf{C})$  such that  $AB = BA$ .  $\square$

#### REFERENCES

- [1] L. Ambrosio, A. Figalli, G. Friesecke, J. Giannoulis, T. Paul: *Semiclassical limit of quantum dynamics with rough potentials and well posedness of transport equations with measure initial data*, Comm. Pure Appl. Math. **64** (2011), 1199–1242.
- [2] L. Ambrosio, N. Fusco, D. Pallara: “Functions of Bounded Variations and Free Discontinuity Problems”, Oxford University Press, Oxford 2000.
- [3] V.I. Arnold: *Characteristic class entering in quantization condition*, Func. Anal. Appl. **1** (1967), 1–14.
- [4] V.I. Arnold: “Geometrical methods of the theory of ordinary differential equations”, Springer-Verlag, New York 1997.
- [5] V.I. Arnold: “Mathematical Methods of Classical Mechanics”, Springer-Verlag, New York 1989.
- [6] Y. Brenier, U. Frisch, M. Hnon, G. Loeper, S. Matarrese, R. Moyhayae, A. Sobolevskii: *Reconstruction of the early Universe as a convex optimization problem*, Mon. Not. Royal. Astron. Soc.
- [7] Y. Brenier, E. Grenier: *Sticky particles and scalar conservation laws*, SIAM J. Numer. Anal. **35** (1998), 2317–2328.
- [8] W. E, Y. Rykov, Ya. Sinai: *Generalized variational principles, global weak solutions and behavior with random initial data for systems of conservation laws arising in adhesion particle dynamics*, Comm. Math. Phys., **177** (1996), 349–380.
- [9] P. Gérard, P. Markowich, N. Mauser, F. Poupaud: *Homogenization limit and Wigner Transforms*, Comm. on Pure and App. Math. **50** (1997), 323–379.
- [10] V. Guillemin, S. Sternberg: “Geometric Asymptotics”, Amer. Math. Soc., Providence 1977.
- [11] L. Hörmander: “The analysis of linear partial differential operators I”, Springer-Verlag, Berlin, Heidelberg 1983, 1990.
- [12] L. Hörmander: “The analysis of linear partial differential operators II”, Springer-Verlag, Berlin, Heidelberg 1983, 1990.
- [13] J. Kauhannen, P. Koskela, J. Malý: *On functions with derivatives in a Lorentz space*, Manuscripta Math. **100** (1999), 87–101.
- [14] L. Landau, E. Lifshitz: “The Classical Theory of Fields”, Butterworth-Heinemann, Oxford, 1976.
- [15] L. Landau, E. Lifshitz: “Quantum Mechanics Nonrelativistic Theory”, Butterworth-Heinemann, Oxford, 1977.
- [16] A. Laptev, I. Sigal: *Global Fourier Integral Operators and Semiclassical Asymptotics*, Review of Math. Phys. **12** (2000), 749–766.

- [17] P.-L. Lions, T. Paul: *Sur les mesures de Wigner*, Rev. Mat. Iberoamericana **9** (1993), 553–618.
- [18] J. Malý: *Absolutely Continuous Functions of Several Variables*, J. Math. Anal. Appl. **231** (1999), 492–508.
- [19] J. Malý, D. Swanson, W. Ziemer: *The Co-Area Formula for Sobolev Mappings*, Trans. Amer. Math. Soc. **355** (2002), 477–492.
- [20] W. Rudin: “Real and Complex Analysis”, McGraw Hill, Singapore, 1986.
- [21] J. Smoller: *Shock Waves and Reaction-Diffusion Equations*; Springer-Verlag, New York, 1983.
- [22] C. Sparber, P. A. Markowich, N. Mauser: *Wigner Functions versus WKB-Methods in Multivalued Geometrical Optics*, Asymptot. Anal., **33** (2003), 153–187.
- [23] E. Wigner: *On the Quantum Correction For Thermodynamic Equilibrium*, Phys. Rev. **40** (1932), 749–759.
- [24] Ya. Zeldovich: *Gravitational instability: An approximate theory for large density perturbations*, Astron. Astrophys. **5** (1970), 84–89.

(C.B.) UNIVERSITÉ PARIS-DIDEROT, LABORATOIRE J.-L. LIONS, BP187, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05 FRANCE

*E-mail address:* `claude.bardos@gmail.com`

(F.G.) ECOLE POLYTECHNIQUE, CENTRE DE MATHÉMATIQUES L. SCHWARTZ, 91128 PALAISEAU CEDEX, FRANCE & UNIVERSITÉ PARIS-DIDEROT, LABORATOIRE J.-L. LIONS, BP187, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05 FRANCE

*E-mail address:* `francois.golse@math.polytechnique.fr`

(P.M.) UNIVERSITY OF CAMBRIDGE, DAMTP, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE, CB3 0WA, UNITED KINGDOM

*E-mail address:* `P.A.Markowich@damtp.cam.ac.uk`

(T.P.) CNRS AND ECOLE POLYTECHNIQUE, CENTRE DE MATHÉMATIQUES L. SCHWARTZ, 91128 PALAISEAU CEDEX, FRANCE

*E-mail address:* `thierry.paul@math.polytechnique.fr`