

Derivation of a homogenized two-temperature model from the heat equation

Laurent Desvillettes, François Golse, Valeria Ricci

► **To cite this version:**

Laurent Desvillettes, François Golse, Valeria Ricci. Derivation of a homogenized two-temperature model from the heat equation. ESAIM: Mathematical Modelling and Numerical Analysis, EDP Sciences, 2014, 48 (2014), pp.1583-1613. 10.1051/m2an/2014011 . hal-00827912

HAL Id: hal-00827912

<https://hal-polytechnique.archives-ouvertes.fr/hal-00827912>

Submitted on 29 May 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DERIVATION OF A HOMOGENIZED TWO-TEMPERATURE MODEL FROM THE HEAT EQUATION

LAURENT DESVILLETTES, FRANÇOIS GOLSE, AND VALERIA RICCI

ABSTRACT. This work studies the heat equation in a two-phase material with spherical inclusions. Under some appropriate scaling on the size, volume fraction and heat capacity of the inclusions, we derive a coupled system of partial differential equations governing the evolution of the temperature of each phase at a macroscopic level of description. The coupling terms describing the exchange of heat between the phases are obtained by using homogenization techniques originating from [D. Cioranescu, F. Murat: Collège de France Seminar vol. 2. (Paris 1979-1980) Res. Notes in Math. vol. 60, pp. 98–138. Pitman, Boston, London, 1982.]

1. DESCRIPTION OF THE PROBLEM

1.1. The homogenized two-temperature model. The purpose of this paper is to derive a model governing the exchange of heat in a composite medium consisting of a background material with very small spherical inclusions of another material with large thermal conductivity. Specifically, we assume that the volume fraction of the inclusions is negligible, while the heat capacity of each inclusion is large.

Under some appropriate scaling assumptions on the size, volume fraction and heat capacity of the inclusions, the temperature field $T \equiv T(t, x)$ of the background material and the temperature field $\theta \equiv \theta(t, x)$ of the dispersed phase (i.e. the inclusions) satisfy

$$(1) \quad \begin{cases} \partial_t T - d\Delta_x T + 4\pi\rho d(T - \theta) = 0, \\ \frac{d}{d'}\partial_t \theta + 4\pi\rho d(\theta - T) = 0, \end{cases}$$

where $\rho \equiv \rho(x)$ is the number density of inclusions while d and d' are the heat diffusion coefficients (i.e. the ratio of the heat conductivity to the volumetric heat capacity) of the background material and the inclusions respectively.

Our work is motivated by a class of models used in the theory of multiphase flows, especially of multiphase flows in porous media. In such flows, each phase can have its own temperature (in which case the flow is said to be in thermal local non-equilibrium). Averaged equations for those temperatures similar to (1) have been proposed in [9], [6] and [2] on the basis of arguments at a macroscopic level of description. While these references address the case of complex realistic flows, our setting is purposely chosen as simple as possible. Neither convection nor phase changes are taken into account in our model. Besides we only consider two phase flows, with only one phase having a positive diffusion rate. The case of positive

1991 *Mathematics Subject Classification.* 35K05, 35B27, 76T05, (35Q79, 76M50).

Key words and phrases. Heat equation; Homogenization; Infinite diffusion limit; Thermal nonequilibrium models.

diffusion rates is considered in [9] (eq. (13) – (15) on p. 242) and in [6] (on p. 2151), while the case of phases with zero diffusion rate is considered in [2] (eqs. (1.63) and (1.74) on p. 39).

We give a rigorous derivation of the coupled system above from a model where the heat conductivity of the dispersed phase is assumed to be infinite from the outset. Our derivation is based on homogenization arguments following our earlier work in [5], inspired from [7, 4].

For the sake of being complete, we also give a rigorous derivation of the infinite conductivity model from the classical heat diffusion equation. In the next two sections, we briefly describe the heat diffusion problem in a binary composite material, and the infinite conductivity model that is our starting point for the homogenization process.

1.2. The model with finite conductivity. Consider an open domain $\Omega \subset \mathbf{R}^3$, let A be an open subset of Ω and let $B = \Omega \setminus A$ be closed in \mathbf{R}^3 . Assume that $\partial\Omega$ and ∂B are submanifolds of \mathbf{R}^3 of class C^2 , and that $B \cap \partial\Omega = \emptyset$. Notice that our results would also hold in the case $\Omega = \mathbf{R}^3$. The unit normal field on the boundary of B is oriented towards A .

The set A is occupied by a material \mathcal{A} with heat conductivity κ_A , density ρ_A and specific heat capacity C_A , while the set B is occupied by a material \mathcal{B} with heat conductivity κ_B , density ρ_B and specific heat capacity C_B . It will be assumed that $\rho_A, C_A, \kappa_A, \rho_B, C_B, \kappa_B$ are continuous positive functions on \bar{A} and B respectively. Denote by $T_A := T_A(t, x) > 0$ and $T_B := T_B(t, x) > 0$ the temperatures of \mathcal{A} and \mathcal{B} at time $t > 0$ and point $x \in A$ ($x \in B$ respectively).

Assuming that Fourier's law holds in both materials and that T_A and T_B are smooth (at least of class C^2) one has

$$(2) \quad \begin{aligned} \rho_A(x)C_A(x)\partial_t T_A(t, x) &= \operatorname{div}_x(\kappa_A(x)\nabla_x T_A(t, x)), & x \in A, t > 0, \\ \rho_B(x)C_B(x)\partial_t T_B(t, x) &= \operatorname{div}_x(\kappa_B(x)\nabla_x T_B(t, x)), & x \in \mathring{B}, t > 0. \end{aligned}$$

If there is no heat source concentrated on the interface ∂B , then the temperature varies continuously across the interface between material \mathcal{A} and material \mathcal{B} and there is no net heat flux across that same interface. In other words, assuming that T_A and T_B are smooth up to the interface ∂B between both materials

$$(3) \quad \begin{cases} T_A(t, x) = T_B(t, x), & x \in \partial B, t > 0, \\ \kappa_A(x)\frac{\partial T_A}{\partial n}(t, x) = \kappa_B(x)\frac{\partial T_B}{\partial n}(t, x), & x \in \partial B, t > 0. \end{cases}$$

Define

$$(4) \quad \rho(x) := \begin{cases} \rho_A(x) & x \in A \\ \rho_B(x) & x \in B \end{cases} \quad C(x) := \begin{cases} C_A(x) & x \in A \\ C_B(x) & x \in B \end{cases}$$

together with

$$(5) \quad \kappa(x) := \begin{cases} \kappa_A(x) & x \in A \\ \kappa_B(x) & x \in B \end{cases}$$

and

$$(6) \quad T(t, x) := \begin{cases} T_A(t, x) & x \in A \\ T_B(t, x) & x \in B \end{cases}$$

Assume that

$$(7) \quad \begin{cases} T_A \in C([0, \tau]; L^2(A)) \cap L^2(0, \tau; H^1(A)), \\ T_B \in C([0, \tau]; L^2(B)) \cap L^2(0, \tau; H^1(B)). \end{cases}$$

In that case the functions T_A and T_B have traces on ∂B denoted $T_A|_{\partial B}$ and $T_B|_{\partial B}$ belonging to $L^2(0, \tau; H^{1/2}(\partial B))$.

Moreover, if T_A and T_B satisfy (2), the vector fields

$$(\rho_A C_A T_A, -\kappa_A \nabla_x T_A) \text{ and } (\rho_B C_B T_B, -\kappa_B \nabla_x T_B)$$

are divergence free in $(0, \tau) \times A$ and $(0, \tau) \times \mathring{B}$ respectively. By statement a) in Lemma A.3, both sides of the second equality in (3) are well defined elements of $H_{00}^{1/2}((0, \tau) \times \partial B)'$. (We recall that $H_{00}^{1/2}((0, \tau) \times \partial B)$ is the Lions-Magenes subspace of functions in $H^{1/2}((0, \tau) \times \partial B)$ whose extension by 0 to $\mathbf{R} \times \partial B$ defines an element of $H^{1/2}(\mathbf{R} \times \partial B)$; the notation $H_{00}^{1/2}((0, \tau) \times \partial B)'$ designates the dual of that space.)

Lemma 1.1. *Assume that T_A and T_B satisfy assumptions (7). Let ρ, C, κ and T be defined as in (4)-(5) and (6). Then*

$$T \in C([0, \tau]; L^2(\Omega)) \cap L^2(0, \tau; H^1(\Omega))$$

and

$$(8) \quad \rho(x)C(x)\partial_t T(t, x) = \operatorname{div}_x(\kappa(x)\nabla_x T(t, x)) \quad x \in \Omega, \quad t > 0$$

holds in the sense of distributions in $(0, \tau) \times \Omega$ if and only if both (2) and (3) hold in the sense of distributions.

Proof. Under the assumption (7), the function T defined by (6) belongs to the space $L^2((0, \tau); H^1(\Omega))$ if and only if the boundary traces of T_A and T_B coincide, i.e.

$$T_A(t, \cdot)|_{\partial B} = T_B(t, \cdot)|_{\partial B} \quad \text{for a.e. } t \in [0, \tau].$$

If (8) holds in the sense of distributions on $(0, \tau) \times \Omega$, then (2) hold in the sense of distributions on $(0, \tau) \times A$ and $(0, \tau) \times B$ respectively.

For $\phi \in C_c^\infty(\Omega)$, one has

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \rho(x)C(x)T(t, x)\phi(x)dx + \int_{\Omega} \kappa(x)\nabla_x T(t, x) \cdot \nabla \phi(x)dx \\ &= \frac{d}{dt} \int_A \rho_A(x)C_A(x)T_A(t, x)\phi(x)dx + \int_A \kappa_A(x)\nabla_x T_A(t, x) \cdot \nabla \phi(x)dx \\ &+ \frac{d}{dt} \int_B \rho_B(x)C_B(x)T_B(t, x)\phi(x)dx + \int_B \kappa_B(x)\nabla_x T_B(t, x) \cdot \nabla \phi(x)dx \\ &= - \left\langle \kappa_A \frac{\partial T_A}{\partial n}, \phi \right\rangle_{H^{-1/2}(\partial B), H^{1/2}(\partial B)} + \left\langle \kappa_B \frac{\partial T_B}{\partial n}, \phi \right\rangle_{H^{-1/2}(\partial B), H^{1/2}(\partial B)} \end{aligned}$$

provided that T_A and T_B satisfy (2), by statement b) in Lemma A.3.

Thus, if T satisfies (8) in the sense of distributions on $(0, \tau) \times \Omega$, then T_A and T_B satisfy (2) on $(0, \tau) \times A$ and $(0, \tau) \times \mathring{B}$ respectively. Therefore the identity above holds with left hand side equal to 0 in the sense of distributions on $(0, \tau)$, so that

$$\left\langle \kappa_A \frac{\partial T_A}{\partial n} - \kappa_B \frac{\partial T_B}{\partial n}, \phi \right\rangle_{H^{-1/2}(\partial B), H^{1/2}(\partial B)} = 0 \quad \text{in } \mathcal{D}'((0, \tau)).$$

This implies in turn the second equality in (3).

Conversely, if T_A and T_B satisfy (2), the above identity holds with right hand side equal to 0 by the second equality in (3). Therefore

$$\frac{d}{dt} \int_{\Omega} \rho(x)C(x)T(t,x)\phi(x)dx + \int_{\Omega} \kappa(x)\nabla_x T(t,x) \cdot \nabla\phi(x)dx = 0$$

for all $\phi \in C_c^\infty(\Omega)$, which implies that (8) holds in the sense of distributions on $(0, \tau) \times \Omega$ by a classical density argument. \square

Therefore, we start from the heat equation (8) with ρ, C, κ as in (4), (5) and we assume that there is no heat flux across $\partial\Omega$, in other words that T satisfies the Neumann boundary condition

$$(9) \quad \frac{\partial T}{\partial n}(t,x) = 0, \quad x \in \partial\Omega, \quad t > 0.$$

1.3. The model with infinite conductivity. In this section we assume that B has N connected components denoted B_i for $i = 1, \dots, N$.

Our first task is to derive the governing equation for the temperature field T in Ω when the material \mathcal{B} filling B has infinite heat conductivity. In that case the temperature T instantaneously reaches equilibrium in each connected component B_i of B , so that

$$(10) \quad T(t,x) = T_i(t), \quad x \in B_i, \quad t > 0$$

for each $i = 1, \dots, N$. Therefore, the unknown for the problem with infinite conductivity is $(T_A(t,x), T_1(t), \dots, T_N(t))$, where

$$(11) \quad \begin{cases} \rho_A(x)C_A(x)\partial_t T_A(t,x) = \operatorname{div}_x(\kappa_A(x)\nabla_x T_A(t,x)), & x \in A, \quad t > 0, \\ \frac{\partial T}{\partial n}(t,x) = 0, & x \in \partial\Omega, \quad t > 0 \\ T_A(t,x) = T_i(t), & x \in \partial B_i, \quad t > 0 \end{cases}$$

This is obviously not enough to determine the evolution of T_A and of T_i for all $i = 1, \dots, N$.

For finite κ_B , the vector field

$$(t,x) \mapsto (\rho_B(x)C_B(x)T_B(t,x), -\kappa_B(x)\nabla_x T_B(t,x))$$

is divergence free in $(0, \tau) \times B_i$ for each $i = 1, \dots, N$. By statement b) in Lemma A.3 and the second equality in (3)

$$\begin{aligned} \frac{d}{dt} \int_{B_i} \rho_B(x)C_B(x)T_B(t,x)dx &= \left\langle \kappa_B(x) \frac{\partial T_B}{\partial n}(t, \cdot), 1 \right\rangle_{H^{-1/2}(\partial B_i), H^{1/2}(\partial B_i)} \\ &= \left\langle \kappa_A(x) \frac{\partial T_A}{\partial n}(t, \cdot), 1 \right\rangle_{H^{-1/2}(\partial B_i), H^{1/2}(\partial B_i)}. \end{aligned}$$

Letting $\kappa_B \rightarrow \infty$ and abusing the integral notation to designate the last duality bracket above, one uses (10) to conclude that

$$(12) \quad \dot{T}_i(t) = \frac{1}{\beta_i} \int_{\partial B_i} \kappa_A(x) \frac{\partial T_A}{\partial n}(t,x) dS(x)$$

where

$$(13) \quad \beta_i := \int_{B_i} \rho_B(x)C_B(x)dx.$$

The argument above suggests that the governing equations for the infinite conductivity problem with unknowns $(T_A(t, x), T_1(t), \dots, T_N(t))$ is the system consisting of (11) with (12) for $i = 1, \dots, N$.

2. MAIN RESULTS

2.1. Existence and uniqueness theory for the heat equation with discontinuous coefficients. Since our starting point is (2) with interface condition (3), or equivalently the heat equation (8) with discontinuous coefficients (see Lemma 1.1), we first recall the existence and uniqueness theory for (8) with Neumann boundary condition (9). Except for the possibly non smooth factor $\rho(x)C(x)$, this is a classical result. This factor can be handled with appropriate weighted Sobolev spaces; for the sake of being complete, we sketch the (elementary) argument below.

Proposition 2.1. *Let $\kappa \equiv \kappa(x)$, $\rho \equiv \rho(x)$ and $C \equiv C(x)$ be measurable functions on Ω satisfying*

$$\kappa_m \leq \kappa(x) \leq \kappa_M, \quad \rho_m \leq \rho(x) \leq \rho_M, \quad C_m \leq C(x) \leq C_M$$

for a.e. $x \in \Omega$, where $\kappa_m, \kappa_M, \rho_m, \rho_M, C_m, C_M > 0$, and let $T^{in} \in L^2(\Omega)$. There exists a unique

$$T \in C_b([0, +\infty); L^2(\Omega)) \cap L^2(0, \tau; H^1(\Omega))$$

for each $\tau > 0$ that is a weak solution of the problem

$$(14) \quad \begin{cases} \rho(x)C(x)\partial_t T(t, x) = \operatorname{div}_x(\kappa(x)\nabla_x T(t, x)), & x \in \Omega, \quad t > 0, \\ \frac{\partial T}{\partial n}(t, x) = 0, & x \in \partial\Omega, \quad t > 0, \\ T(0, x) = T^{in}(x), & x \in \Omega. \end{cases}$$

This solution satisfies

$$\rho C \partial_t T \in L^2(0, \tau; H^1(\Omega)'),$$

for each $\tau > 0$, together with the “energy” identity

$$\frac{1}{2} \int_{\Omega} \rho(x)C(x)T(t, x)^2 dx + \int_0^t \int_{\Omega} \kappa(x)|\nabla_x T(t, x)|^2 dx dt = \frac{1}{2} \int_{\Omega} \rho(x)C(x)T^{in}(x)^2 dx$$

for each $t > 0$.

We recall the weak formulation of (14): for each $w \in H^1(\Omega)$

$$\langle \rho C \partial_t T(t, \cdot), w \rangle_{H^1(\Omega)', H^1(\Omega)} + \int_{\Omega} \kappa(x)\nabla_x T(t, x) \cdot \nabla_x w(t, x) dx = 0 \text{ for a.e. } t \geq 0.$$

The Neumann condition in (14) is contained in the choice of $L^2([0, +\infty); H^1(\Omega))$ as the set of test functions in the weak formulation above, while there is no difficulty with initial condition since $T \in C([0, +\infty); L^2(\Omega))$.

2.2. The infinite conductivity limit.

2.2.1. *Variational formulation of the infinite conductivity problem.* Assume as in section 1.3 that B has N connected components denoted B_i for $i = 1, \dots, N$. The heat diffusion problem with infinite heat conductivity in B is:

$$(15) \quad \begin{cases} \rho_A C_A \partial_t T(t, x) = \operatorname{div}_x(\kappa_A \nabla_x T(t, x)), & x \in A, t > 0, \\ \frac{\partial T}{\partial n}(t, x) = 0, & x \in \partial\Omega, t > 0, \\ T(t, x) = T_i(t), & x \in \partial B_i, t > 0, 1 \leq i \leq N, \\ \beta_i \dot{T}_i(t) = \int_{\partial B_i} \kappa_A \frac{\partial T}{\partial n}(t, x) dS(x), & t > 0, 1 \leq i \leq N, \\ T(0, x) = T^{in}(x), & x \in \Omega. \end{cases}$$

Its variational formulation is as follows.

Let \mathcal{H}_N be the closed subspace of $L^2(\Omega)$ defined as

$$\mathcal{H}_N := \left\{ u \in L^2(\Omega) \text{ s.t. } u(x) = \frac{1}{|B_i|} \int_{B_i} u(y) dy \text{ for a.e. } x \in B_i, i = 1, \dots, N \right\};$$

and equipped with the inner product

$$(u|v)_{\mathcal{H}_N} = \int_{\Omega} u(x)v(x)\rho(x)C(x)dx.$$

Define

$$\mathcal{V}_N := \mathcal{H}_N \cap H^1(\Omega)$$

with the inner product

$$(u|v)_{\mathcal{V}_N} = (u|v)_{\mathcal{H}_N} + \int_A \nabla u(x) \cdot \nabla v(x) \rho_A(x) C_A(x) dx.$$

Obviously \mathcal{V}_N is a separable Hilbert space, the inclusion $\mathcal{V}_N \subset \mathcal{H}_N$ is continuous and \mathcal{V}_N is a dense subspace of \mathcal{H}_N . Besides, the map $\mathcal{H}_N \ni u \mapsto L_u \in \mathcal{V}'_N$, where L_u is the linear functional $v \mapsto (u|v)_{\mathcal{H}_N}$, identifies \mathcal{H}_N with a dense subspace of \mathcal{V}'_N .

The variational formulation of the infinite conductivity problem is as follows: a weak solution of (15) is a function

$$(16) \quad T \in C([0, \tau]; \mathcal{H}_N) \cap L^2(0, \tau; \mathcal{V}_N) \text{ such that } \rho C \partial_t T \in L^2(0, \tau; \mathcal{V}'_N)$$

satisfying the initial condition and

$$(17) \quad \begin{cases} \partial_t(T(t, \cdot)|w)_{\mathcal{H}_N} + \int_A \kappa_A(x) \nabla_x T(t, x) \cdot \nabla w(x) dx = 0 \text{ for a.e. } t \in [0, \tau] \\ \text{for each test function } w \in \mathcal{V}_N \end{cases}$$

This variational formulation is justified by the following observation.

Proposition 2.2. *Let T satisfy (16) and the initial condition in (15).*

If T satisfies the variational condition (17), then

$$(18) \quad \rho_A C_A \partial_t T = \operatorname{div}_x(\kappa_A \nabla_x T) \quad \text{in } \mathcal{D}'((0, \tau) \times A)$$

and

$$(19) \quad \kappa_A \frac{\partial T}{\partial n} \Big|_{(0, \tau) \times \partial\Omega} = 0 \quad \text{in } H_{00}^{1/2}((0, \tau) \times \partial\Omega)'$$

while

$$(20) \quad \beta_i \dot{T}_i = \left\langle \kappa_A \frac{\partial T}{\partial n} \Big|_{\partial B_i}, 1 \right\rangle_{H^{-1/2}(\partial B_i), H^{1/2}(\partial B_i)} \quad \text{in } H^{-1}((0, \tau))$$

for each $i = 1, \dots, N$, where

$$T_i(t) := \frac{1}{|B_i|} \int_{B_i} T(t, x) dx.$$

Conversely, if T satisfies both (18), (19) and (20), it must satisfy the variational formulation (17).

The existence and uniqueness of a weak solution of the infinite heat conductivity problem is given in the next proposition.

Proposition 2.3. *Assume that κ_A is a measurable function defined a.e. on A satisfying*

$$(21) \quad \kappa_m \leq \kappa_A(x) \leq \kappa_M, \quad \text{for a.e. } x \in A,$$

where κ_m and κ_M are positive numbers, while ρ and C satisfy the same assumptions as in Proposition 2.1. Then for each $T^{in} \in \mathcal{H}_N$, there exists a unique weak solution T of (15) defined for all $t \in [0, +\infty)$. This solution satisfies

$$\rho C \partial_t T \in L^2(0, \tau; \mathcal{V}'_N),$$

for all $\tau > 0$, together with the “energy” identity

$$\begin{aligned} \frac{1}{2} \int_A \rho_A(x) C_A(x) T(t, x)^2 dx + \frac{1}{2} \sum_{i=1}^N \beta_i T_i(t)^2 + \int_0^t \int_A \kappa_A(x) |\nabla_x T(t, x)|^2 dx dt \\ = \frac{1}{2} \int_A \rho_A(x) C_A(x) T^{in}(x)^2 dx + \frac{1}{2} \sum_{i=1}^N \beta_i |T_i^{in}|^2 \end{aligned}$$

for each $t > 0$, where

$$T_i(t) := \frac{1}{|B_i|} \int_{B_i} T(t, y) dy \quad \text{and} \quad T_i^{in} := \frac{1}{|B_i|} \int_{B_i} T^{in}(y) dy.$$

Notice that this existence and uniqueness result assumes that the initial temperature field T^{in} is a constant in each connected component of B . This assumption is implied by the requirement that $T^{in} \in \mathcal{H}_N$. While this restriction may seem questionable, it is very natural from the mathematical viewpoint. For general initial temperature fields T^{in} , the solution of (15) would include an initial layer corresponding with the relaxation to thermal equilibrium in each connected component of B . Such initial layers involve fast variations of the temperature field that are incompatible with the condition $\rho C \partial_t T \in L^2(0, \tau; \mathcal{V}'_N)$ in the infinite conductivity limit.

2.2.2. *Convergence to the infinite conductivity model.* For each $\eta > 0$, let κ_η be defined as follows:

$$(22) \quad \kappa_\eta(x) := \begin{cases} \kappa_A(x) & x \in A \\ \kappa_B(x)/\eta & x \in B \end{cases}$$

where κ_A and κ_B are measurable functions on A and B respectively satisfying

$$(23) \quad \kappa_m \leq \kappa_A(x) \leq \kappa_M \text{ and } \kappa_m \leq \kappa_B(y) \leq \kappa_M, \quad \text{for a.e. } x \in A \text{ and } y \in B,$$

κ_M and κ_m being two positive constants.

Theorem 2.4. *Assume that ρ and C satisfy the same assumptions as in Proposition 2.1, while κ_A and κ_B satisfy (23). Let $T^{in} \in \mathcal{H}_N$. For each $\eta > 0$, let $T_\eta \in C_b([0, +\infty); L^2(\Omega)) \cap L^2(0, \tau; H^1(\Omega))$ for all $\tau > 0$ be the weak solution of (14) with heat conductivity κ_η defined as in (22) and initial data T^{in} . Then*

$$T_\eta \rightarrow T \text{ in } L^2(0, \tau; H^1(\Omega))$$

as $\eta \rightarrow 0$ for all $\tau > 0$, where $T \in C_b([0, +\infty); \mathcal{H}_N) \cap L^2(0, \tau; \mathcal{V}_N)$ for all $\tau > 0$ is the weak solution of the infinite conductivity problem (15).

2.3. The homogenized system. Let $\sigma, \sigma' > 0$ and let $\rho \in C_b(\overline{\Omega})$ be a probability density on $\overline{\Omega}$ such that $1/\rho$ is bounded on $\overline{\Omega}$. Let $T^{in}, \vartheta^{in} \in L^2(\Omega)$. Consider the system

$$(24) \quad \begin{cases} (\partial_t - \sigma \Delta_x)T(t, x) + 4\pi\sigma(\rho(x)T(t, x) - \vartheta(t, x)) = 0, & x \in \Omega, t > 0, \\ \partial_t \vartheta(t, x) + 4\pi\sigma'(\vartheta(t, x) - \rho(x)T(t, x)) = 0, & x \in \Omega, t > 0, \\ \frac{\partial T}{\partial n}(t, x) = 0, & x \in \partial\Omega, t > 0, \\ T(0, x) = T^{in}(x), \quad \vartheta(0, x) = \vartheta^{in}(x), & x \in \Omega. \end{cases}$$

A weak solution of (24) is a pair (T, ϑ) such that

$$T \in L^\infty([0, +\infty); L^2(\Omega)) \cap L^2(0, \tau; H^1(\Omega)) \quad \text{and} \quad \vartheta \in L^\infty([0, +\infty); L^2(\Omega)),$$

for all $\tau > 0$, and

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} T(t, x)\phi(x)dx + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x)dx \\ + 4\pi\sigma \int_{\Omega} (\rho(x)T(t, x) - \vartheta(t, x))\phi(x)dx = 0 \\ \frac{d}{dt} \int_{\Omega} \vartheta(t, x)\psi(x)dx + 4\pi\sigma' \int_{\Omega} (\vartheta(t, x) - \rho(x)T(t, x))\psi(x)dx = 0 \end{aligned}$$

in the sense of distributions on $(0, +\infty)$ for each $\phi \in H^1(\Omega)$ and $\psi \in L^2(\Omega)$, together with the initial condition. Observe that the identities above imply that

$$\frac{d}{dt} \int_{\Omega} T(t, x)\phi(x)dx \quad \text{and} \quad \frac{d}{dt} \int_{\Omega} \vartheta(t, x)\psi(x)dx \in L^2([0, \tau])$$

for each $\tau > 0$, so that the functions

$$t \mapsto \int_{\Omega} T(t, x)\phi(x)dx \quad \text{and} \quad t \mapsto \int_{\Omega} \vartheta(t, x)\psi(x)dx$$

are continuous on $[0, +\infty)$. Therefore the initial condition, interpreted as

$$\int_{\Omega} T(0, x)\phi(x)dx = \int_{\Omega} T^{in}(x)\phi(x)dx, \quad \int_{\Omega} \vartheta(0, x)\psi(x)dx = \int_{\Omega} \vartheta^{in}(x)\psi(x)dx$$

for all $\phi \in H^1(\Omega)$ and all $\psi \in L^2(\Omega)$, makes perfect sense.

In the next proposition, we state the basic results concerning the existence and uniqueness of a weak solution of the initial-boundary value problem for the homogenized system. In fact, one can say more about the continuity in time of (T, ϑ) , as explained below.

Proposition 2.5. *Under the assumptions above, any weak solution of (24) satisfies*

$$\partial_t T \in L^2(0, \tau; H^1(\Omega)') \quad \text{and} \quad \partial_t \vartheta \in L^2(0, \tau; L^2(\Omega)),$$

and (up to modification on some negligible t -set)

$$T, \vartheta \in C_b([0, +\infty); L^2(\Omega)).$$

Moreover, there exists a unique weak solution of the system (24). It is a solution of the partial differential equations

$$\begin{cases} \partial_t T - \sigma \Delta_x T + 4\pi\sigma(\rho T - \vartheta) = 0, \\ \partial_t \vartheta + 4\pi\sigma'(\vartheta - \rho T) = 0, \end{cases}$$

in the sense of distributions on $(0, +\infty) \times \Omega$, and satisfies the Neumann condition

$$\left. \frac{\partial T}{\partial n} \right|_{(0, \tau) \times \partial\Omega} = 0$$

in $H_{00}^{1/2}((0, \tau) \times \partial\Omega)'$ for each $\tau > 0$.

In fact, the existence of the solution of (24) follows from Theorem 2.6

2.4. The homogenization limit. Henceforth we assume that the material \mathcal{B} occupies N identical spherical inclusions with radius ϵ :

$$(25) \quad B_\epsilon = \bigcup_{i=1}^N B_i \quad \text{where} \quad B_i := \overline{B(x_i, \epsilon)}, \quad i = 1, \dots, N$$

and henceforth denote

$$(26) \quad A_\epsilon = \Omega \setminus B_\epsilon.$$

The number of inclusions N is assumed to scale as

$$(27) \quad N = 1/\epsilon.$$

The inclusion centers x_i are distributed so that their empirical distribution satisfies

$$(28) \quad \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \rightarrow \rho \mathcal{L}^3$$

in the weak topology of probability measures, where \mathcal{L}^3 designates the 3-dimensional Lebesgue measure and

$$(29) \quad \rho \text{ and } 1/\rho \in C_b(\overline{\Omega}), \quad \int_{\Omega} \rho(x) dx = 1.$$

Besides, we also assume that

$$(30) \quad \frac{1}{N} \sum_{i=1}^N |x_i|^2 \leq C^{in} \quad \text{for all } N \geq 1$$

for some positive constant C^{in} . Finally, we denote

$$(31) \quad r_\epsilon = \epsilon^{1/3}$$

and assume that the inclusion centers are chosen so that

$$(32) \quad |x_i - x_j| > 2r_\epsilon \quad \text{for all } i, j = 1, \dots, N.$$

For simplicity we assume that ρ_A, C_A and κ_A are constants, and define

$$(33) \quad \sigma = \kappa_A / \rho_A C_A.$$

We further assume that ρ_B and C_B are scaled with ϵ so that $\rho_B C_B \sim \text{Const.}/\epsilon^2$, and introduce the constant

$$(34) \quad \sigma' = 3\kappa_A / 4\pi\rho_B C_B \epsilon^2.$$

The scaled infinite heat conductivity problem takes the form

$$(35) \quad \begin{cases} \partial_t T_\epsilon(t, x) = \sigma \Delta_x T_\epsilon(t, x), & x \in A_\epsilon, t > 0, \\ \frac{\partial T_\epsilon}{\partial n}(t, x) = 0, & x \in \partial\Omega, t > 0, \\ T_\epsilon(t, x) = T_{i,\epsilon}(t), & x \in \partial B(x_i, \epsilon), t > 0, 1 \leq i \leq N, \\ \dot{T}_{i,\epsilon}(t) = \frac{\sigma'}{\epsilon} \int_{\partial B(x_i, \epsilon)} \frac{\partial T_\epsilon}{\partial n}(t, x) dS(x), & t > 0, 1 \leq i \leq N, \\ T_\epsilon(0, x) = T_\epsilon^{in}(x), & x \in \Omega. \end{cases}$$

The initial data $T_\epsilon^{in} \in \mathcal{H}_N$, so that T_ϵ^{in} is a.e. a constant in $B(x_i, \epsilon)$:

$$(36) \quad T_{i,\epsilon}^{in} := \frac{3}{4\pi\epsilon^3} \int_{B(x_i, \epsilon)} T^{in}(x) dx$$

Then

$$\begin{aligned} |T_\epsilon^{in}|_{\mathcal{H}_N}^2 &= \rho_A C_A \int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \sum_{i=1}^N \frac{4\pi}{3} \rho_B C_B \epsilon^3 |T_{i,\epsilon}^{in}|^2 \\ &= \rho_A C_A \left(\int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \frac{\sigma}{\sigma'} \epsilon \sum_{i=1}^N |T_{i,\epsilon}^{in}|^2 \right) \end{aligned}$$

We shall henceforth assume that the initial data satisfies

$$|T_\epsilon^{in}|_{\mathcal{H}_N}^2 = O(1)$$

i.e. that there exists a positive constant, taken equal to C^{in} for notational simplicity, such that

$$(37) \quad \int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \frac{\sigma}{\sigma'} \epsilon \sum_{i=1}^N |T_{i,\epsilon}^{in}|^2 \leq C^{in} \quad \text{for all } \epsilon > 0.$$

Theorem 2.6. *Assume that (27) holds, that the distribution of inclusion centers satisfies (28) and (30), that the volumetric heat capacity of the material in the inclusions scales as prescribed in (34), and that the initial data T_ϵ^{in} satisfies the bound (37). Assume further that*

$$T_\epsilon^{in} \rightarrow T^{in} \quad \text{in } L^2(\Omega) \text{ weak as } \epsilon \rightarrow 0$$

while¹

$$\frac{1}{N} \sum_{i=1}^N T_{i,\epsilon}^{in} \delta_{x_i} \rightarrow \vartheta^{in} \text{ in } \mathcal{M}_b(\Omega) \text{ weak-}^* \text{ as } \epsilon \rightarrow 0.$$

¹The notation $\mathcal{M}_b(\Omega)$ designates the set of bounded (signed) Radon measures on Ω .

Let $T_\epsilon \in C([0, +\infty); \mathcal{H}_N) \cap L^2(0, \tau; \mathcal{V}_N)$ for all $\tau > 0$ be the weak solution of the scaled infinite heat conductivity problem (35). Then, in the limit as $\epsilon \rightarrow 0$,

$$T_\epsilon \rightarrow T \quad \begin{cases} \text{in } L^2(0, \tau; H^1(\Omega)) \text{ weak for all } \tau > 0 \\ \text{and in } L^\infty([0, +\infty); L^2(\Omega)) \text{ weak-}^* \end{cases}$$

and

$$\vartheta_\epsilon := \frac{1}{N} \sum_{i=1}^N T_{i,\epsilon} \delta_{x_i} \rightarrow \vartheta \text{ in } L^\infty([0, +\infty); \mathcal{M}_b(\Omega)) \text{ weak-}^*$$

where

$$T_{i,\epsilon} := \frac{3}{4\pi\epsilon^3} \int_{B(x_i, \epsilon)} T_\epsilon(t, x) dx.$$

Besides

$$T \in C_b([0, +\infty); L^2(\Omega)) \times L^2(0, \tau; H^1(\Omega)) \text{ for each } \tau > 0$$

while

$$\vartheta \in C_b([0, +\infty); L^2(\Omega)).$$

Finally, the pair (T, ϑ) is the unique weak solution of the homogenized system (24) with initial condition

$$T|_{t=0} = T^{in}, \quad \vartheta|_{t=0} = \vartheta^{in}.$$

3. PROOFS OF PROPOSITIONS 2.1, 2.2 AND 2.3

Proof of Proposition 2.1. Consider the Hilbert spaces $\mathcal{H} = L^2(\Omega)$ and $\mathcal{V} = H^1(\Omega)$ equipped with the inner products

$$(u|v)_\mathcal{H} := \int_\Omega u(x)v(x)\rho(x)C(x)dx,$$

$$(u|v)_\mathcal{V} := \int_\Omega (u(x)v(x) + \nabla u(x) \cdot \nabla v(x))\rho(x)C(x)dx.$$

Let a be the bilinear form defined on $\mathcal{V} \times \mathcal{V}$ by

$$a(u, v) = \int_\Omega \kappa(x) \nabla_x u(x) \cdot \nabla_x v(x) dx;$$

observe that

$$|a(u, v)| \leq \frac{\kappa_M}{\rho_m C_m} (u|u)_\mathcal{V}^{1/2} (v|v)_\mathcal{V}^{1/2}$$

while

$$a(u, u) \geq \frac{\kappa_m}{\rho_M C_M} ((u|u)_\mathcal{V} - (u|u)_\mathcal{H}).$$

By Theorem X.9 in [3], there exists a unique $T \in L^2(0, \tau; \mathcal{V}) \cap C_b([0, \tau]; \mathcal{H})$ such that $\rho C \partial_t T \in L^2(0, \tau; \mathcal{V}')$ for each $\tau > 0$ such that the linear functional

$$L(t) : w \mapsto \partial_t (T(t, \cdot) | w)_\mathcal{H} + a(T(t, \cdot), w) = \langle \rho C \partial_t T(t, \cdot), w \rangle_{\mathcal{V}', \mathcal{V}} + a(T(t, \cdot), w)$$

satisfies

$$\langle L(t), w \rangle_{\mathcal{V}', \mathcal{V}} = 0 \text{ for a.e. } t \in [0, +\infty)$$

for all $w \in \mathcal{V}$. Equivalently, T is the unique weak solution of (14).

By Lemma A.2, this linear functional satisfies $L(t) = 0$ for a.e. $t \in [0, +\infty)$. In particular

$$0 = \langle L(s), T(s, \cdot) \rangle_{\mathcal{V}', \mathcal{V}} = \langle \rho C \partial_t T(s, \cdot), T(s, \cdot) \rangle_{\mathcal{V}', \mathcal{V}} + a(T(s, \cdot), T(s, \cdot))$$

for a.e. $s \in [0, +\infty)$. Integrating in $s \in [0, t]$ and applying statement b) of Lemma A.1 give the “energy identity”. \square

Proof of Proposition 2.2. Specializing (17) to the case where $w \in C_c^\infty(A)$ implies (18). In particular, the vector field

$$(0, \tau) \times A \ni (t, x) \mapsto (\rho_A(x)C_A(x)T(t, x), -\kappa_A(x)\nabla_x T(t, x))$$

is divergence free in $(0, \tau) \times A$. Applying statement b) in Lemma A.3 shows that, for each $w \in \mathcal{V}_N$, one has

$$\begin{aligned} 0 &= \frac{d}{dt} \int_{\Omega} \rho(x)C(x)T(t, x)w(x)dx + \int_A \kappa_A(x)\nabla_x T(t, x) \cdot \nabla w(x)dx \\ &= \frac{d}{dt} \int_A \rho_A(x)C_A(x)T(t, x)w(x)dx + \sum_{i=1}^N \beta_i w_i \dot{T}_i(t) \\ &\quad + \int_A \kappa_A(x)\nabla_x T(t, x) \cdot \nabla w(x)dx \\ &= \sum_{i=1}^N w_i \left(\beta_i \dot{T}_i(t) - \left\langle \kappa_A \frac{\partial T}{\partial n} \Big|_{\partial B_i}(t, \cdot), 1 \right\rangle_{H^{-1/2}(\partial B_i), H^{1/2}(\partial B_i)} \right) \\ &\quad + \left\langle \kappa_A \frac{\partial T}{\partial n} \Big|_{\partial \Omega}(t, \cdot), w|_{\partial \Omega} \right\rangle_{H^{-1/2}(\partial \Omega), H^{1/2}(\partial \Omega)}, \end{aligned}$$

where

$$w_i := \frac{1}{|B_i|} \int_{B_i} w(y)dy, \quad i = 1, \dots, N.$$

Since this is true for all $w \in \mathcal{V}_N$, and therefore for all $(w_1, \dots, w_N) \in \mathbf{R}^N$, one concludes that

$$\beta_i \dot{T}_i - \left\langle \kappa_A \frac{\partial T}{\partial n} \Big|_{\partial B_i}, 1 \right\rangle_{H^{-1/2}(\partial B_i), H^{1/2}(\partial B_i)} = 0$$

in $H^{-1}((0, \tau))$ for all $i = 1, \dots, N$, and

$$\kappa_A \frac{\partial T}{\partial n} \Big|_{\partial \Omega} = 0$$

in $H_0^{1/2}((0, \tau) \times \partial \Omega)'$.

Conversely, if T satisfies (18), (19) and (20), the equality above shows that (17) holds. \square

Proof of Proposition 2.3. Let b be the bilinear form defined on $\mathcal{V}_N \times \mathcal{V}_N$ by

$$b(u, v) = \int_A \kappa_A(x)\nabla_x u(x) \cdot \nabla_x v(x)dx;$$

observe that

$$|b(u, v)| \leq \frac{\kappa_M}{\rho_m C_m} (u|u)_{\mathcal{V}_N}^{1/2} (v|v)_{\mathcal{V}_N}^{1/2}$$

while

$$b(u, u) \geq \frac{\kappa_m}{\rho_M C_M} ((u|u)_{\mathcal{V}_N} - (u|u)_{\mathcal{H}_N}).$$

By the same argument as in the proof of Proposition 2.1, for each $T^{in} \in \mathcal{H}_N$, there exists a unique weak solution of (15), and this solution satisfies the energy identity in the statement of Proposition 2.3. \square

4. PROOF OF THEOREM 2.4

We keep the notation used in the proof of Proposition 2.1, especially with the same definitions of a, b, \mathcal{H} and \mathcal{V} .

For each $\eta > 0$, the weak solution T_η of (14) satisfies the energy identity

$$\begin{aligned} \frac{1}{2} \int_{\Omega} \rho(x) C(x) T_\eta(t, x)^2 dx + \int_0^t \int_A \kappa_A(x) |\nabla_x T_\eta(s, x)|^2 dx ds \\ + \frac{1}{\eta} \int_0^t \int_B \kappa_B(x) |\nabla_x T_\eta(s, x)|^2 dx ds \\ = \frac{1}{2} \int_{\Omega} \rho(x) C(x) T^{in}(x)^2 dx. \end{aligned}$$

Hence, for $\eta \in (0, 1)$, one has

$$|T_\eta(t, \cdot)|_{\mathcal{H}}^2 \leq |T^{in}|_{\mathcal{H}}^2 \quad \text{and} \quad \int_0^\infty |\nabla_x T_\eta(t, \cdot)|_{\mathcal{H}}^2 dt \leq \frac{\rho_M C_M}{2\kappa_m} |T^{in}|_{\mathcal{H}}^2.$$

Applying the Banach-Alaoglu theorem shows that the family T_η is relatively compact in $L^\infty([0, +\infty); \mathcal{H})$ weak-* and in $L^2([0, +\infty); \mathcal{V})$ weak. Let T be a limit point of T_η ; passing to the limit in the energy identity above shows that, by convexity and weak limit,

$$\int_0^\infty \int_B |\nabla_x T(t, x)|^2 dx dt = 0.$$

Thus the function $x \mapsto T(t, x)$ is constant on B_i for $i = 1, \dots, N$ for a.e. $t \geq 0$ and $T \in L^\infty([0, +\infty); \mathcal{H}_N) \cap L^2(0, \tau; \mathcal{V}_N)$.

Write the variational formulation of (14) for a test function $w \in \mathcal{V}_N \subset \mathcal{V}$:

$$\frac{d}{dt} (T_\eta |w)_{\mathcal{H}} + a(T_\eta, w) = 0 \quad \text{in } L^2([0, \tau]) \text{ for all } \tau > 0.$$

Passing to the limit in a subsequence of T_η converging to T in $L^\infty([0, +\infty); \mathcal{H})$ weak-* and in $L^2(0, \tau; \mathcal{V})$ weak, one finds that

$$\begin{aligned} a(T_\eta, w) &= \int_A \kappa_A(x) \nabla_x T_\eta(t, x) \cdot \nabla w(x) dx + \frac{1}{\eta} \int_B \kappa_B(x) \nabla_x T_\eta(t, x) \cdot \nabla w(x) dx \\ &= \int_A \kappa_A(x) \nabla_x T_\eta(t, x) \cdot \nabla w(x) dx \\ &\rightarrow \int_A \kappa_A(x) \nabla_x T(t, x) \cdot \nabla w(x) dx = b(T, w) \text{ weakly in } L^2([0, \tau]) \end{aligned}$$

since $\nabla_x T_\eta \rightarrow \nabla_x T$ weakly in $L^2([0, \tau] \times \Omega)$. (The second equality above come from the fact that $\nabla w = 0$ on B since $w \in \mathcal{V}_N$.) On the other hand, for each $w \in \mathcal{V}_N$

$$\begin{aligned} \int_0^\tau \left| \frac{d}{dt} (T_\eta(t, \cdot) | w)_{\mathcal{H}} \right|^2 dt &= \int_0^\tau |a(T_\eta(t, \cdot), w)|^2 dt \\ &= \int_0^\tau \left| \int_A \kappa_A(x) \nabla_x T_\eta(t, x) \cdot \nabla w(x) dx \right|^2 dt \\ &\leq \int_0^\tau \int_A \kappa_A(x) |\nabla_x T_\eta(t, x)|^2 dx dt \int_A \kappa_A(x) |\nabla w(x)|^2 dx \\ &\leq \frac{\kappa_M}{2\rho_m C_m} |T^{in}|_{\mathcal{H}}^2 |w|_{\mathcal{V}}^2 \end{aligned}$$

while

$$(T_\eta | w)_{\mathcal{H}} \rightarrow (T | w)_{\mathcal{H}} \text{ in } L^\infty([0, +\infty)) \text{ weak-}^*.$$

Therefore, for each $w \in \mathcal{V}_N$, one has

$$\frac{d}{dt} (T | w)_{\mathcal{H}} + b(T, w) = 0 \quad \text{in } L^2([0, \tau]) \text{ for all } \tau > 0,$$

which implies in particular that

$$\rho C \partial_t T \in L^2(0, \tau; \mathcal{V}'_N),$$

and therefore $T \in C_b([0, +\infty); \mathcal{H}_N)$ by statement a) of Lemma A.1. Besides, by the Ascoli-Arzelà theorem,

$$(T_\eta(t, \cdot) | w)_{\mathcal{H}} \rightarrow (T(t, \cdot) | w)_{\mathcal{H}} \quad \text{uniformly in } t \in [0, \tau] \text{ for all } \tau > 0.$$

In particular

$$(T_\eta(0, \cdot) | w)_{\mathcal{H}} = (T^{in} | w)_{\mathcal{H}} \rightarrow (T(0, \cdot) | w)_{\mathcal{H}}$$

so that

$$T(0, \cdot) = T^{in}.$$

In other words T is the weak solution of (15) with initial data T^{in} — the uniqueness of the weak solution following from Proposition 2.3. By compactness of the family T_η and uniqueness of the limit point, we conclude that

$$T_\eta \rightarrow T \text{ in } L^\infty([0, +\infty); \mathcal{H}) \text{ weak-}^* \text{ and in } L^2(0, \tau; \mathcal{V}) \text{ weak}$$

as $\eta \rightarrow 0$.

The energy identities in Propositions 2.1 and 2.3 are recast in the form

$$\begin{aligned} \frac{1}{2} \int_\Omega \rho(x) C(x) T_\eta(t, x)^2 dx + \int_0^t \int_A \kappa_A(x) |\nabla_x T_\eta(s, x)|^2 dx ds \\ + \frac{1}{\eta} \int_0^t \int_B \kappa_B(x) |\nabla_x T_\eta(s, x)|^2 dx ds \\ = \frac{1}{2} \int_\Omega \rho(x) C(x) T^{in}(x)^2 dx, \end{aligned}$$

and

$$\begin{aligned} \frac{1}{2} \int_\Omega \rho(x) C(x) T(t, x)^2 dx + \int_0^t \int_A \kappa_A(x) |\nabla_x T(s, x)|^2 dx ds \\ = \frac{1}{2} \int_\Omega \rho(x) C(x) T^{in}(x)^2 dx. \end{aligned}$$

(Notice that the condition $T^{in} \in \mathcal{H}_N$ is essential in order that

$$\frac{1}{2} \int_{\Omega} \rho(x) C(x) T^{in}(x)^2 dx = \frac{1}{2} \int_A \rho_A(x) C_A(x) T^{in}(x)^2 dx + \frac{1}{2} \sum_{i=1}^N \beta_i |T_i^{in}|^2;$$

likewise

$$\frac{1}{2} \int_{\Omega} \rho(x) C(x) T(t, x)^2 dx = \frac{1}{2} \int_A \rho_A(x) C_A(x) T(t, x)^2 dx + \frac{1}{2} \sum_{i=1}^N \beta_i T_i(t)^2$$

since $T(t, \cdot) \in \mathcal{H}_N$ for all $t > 0$.)

On the other hand, by convexity and weak convergence

$$\frac{1}{2} \int_{\Omega} \rho(x) C(x) T(t, x)^2 dx \leq \liminf_{\eta \rightarrow 0^+} \frac{1}{2} \int_{\Omega} \rho(x) C(x) T_{\eta}(t, x)^2 dx \text{ for all } t > 0,$$

and

$$\int_0^t \int_A \kappa_A(x) |\nabla_x T(s, x)|^2 dx ds \leq \liminf_{\eta \rightarrow 0^+} \int_0^t \int_A \kappa_A(x) |\nabla_x T_{\eta}(s, x)|^2 dx ds.$$

We conclude from the energy identities recalled above that

$$\frac{1}{2} \int_{\Omega} \rho(x) C(x) T_{\eta}(t, x)^2 dx \rightarrow \frac{1}{2} \int_{\Omega} \rho(x) C(x) T(t, x)^2 dx \text{ for all } t > 0,$$

while

$$\begin{cases} \int_0^t \int_A \kappa_A(x) |\nabla_x T_{\eta}(s, x)|^2 dx ds \rightarrow \int_0^t \int_A \kappa_A(x) |\nabla_x T(s, x)|^2 dx ds, \\ \frac{1}{\eta} \int_0^t \int_B \kappa_B(x) |\nabla_x T_{\eta}(s, x)|^2 dx ds \rightarrow 0, \end{cases}$$

for all $t > 0$.

Therefore

$$T_{\eta} \rightarrow T \text{ and } \nabla_x T_{\eta} \rightarrow \nabla_x T \text{ strongly in } L^2([0, \tau] \times \Omega)$$

as $\eta \rightarrow 0$.

5. PROOF OF PROPOSITION 2.5

Since

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} T(t, x) \phi(x) dx + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x) dx \\ + 4\pi\sigma \int_{\Omega} (\rho(x) T(t, x) - \vartheta(t, x)) \phi(x) dx = 0 \\ \frac{d}{dt} \int_{\Omega} \vartheta(t, x) \psi(x) dx + 4\pi\sigma' \int_{\Omega} (\vartheta(t, x) - \rho(x) T(t, x)) \psi(x) dx = 0 \end{aligned}$$

with

$$T \in L^2(0, \tau; H^1(\Omega)) \quad \text{and} \quad \vartheta \in L^2(0, \tau; L^2(\Omega))$$

one has

$$\left| \frac{d}{dt} \int_{\Omega} T(t, x) \phi(x) dx \right| \leq (4\pi + 1) \sigma (\|\rho\|_{L^\infty} \|T(t, \cdot)\|_{H^1(\Omega)} + \|\vartheta\|_{L^2(\Omega)}) \|\phi\|_{H^1(\Omega)}$$

and

$$\left| \frac{d}{dt} \int_{\Omega} \vartheta(t, x) \psi(x) dx \right| \leq 4\pi\sigma'(\|\rho\|_{L^\infty} \|T(t, \cdot)\|_{L^2(\Omega)} + \|\vartheta\|_{L^2(\Omega)}) \|\psi\|_{L^2(\Omega)}$$

so that the linear functionals

$$\phi \mapsto \frac{d}{dt} \int_{\Omega} T(t, x) \phi(x) dx \quad \text{and} \quad \psi \mapsto \frac{d}{dt} \int_{\Omega} \vartheta(t, x) \psi(x) dx$$

are continuous on $H^1(\Omega)$ and on $L^2(\Omega)$ respectively with values in $L^2([0, \tau])$. Therefore

$$\partial_t T \in L^2(0, \tau; H^1(\Omega)') \quad \text{and} \quad \partial_t \vartheta \in L^2([0, \tau] \times \Omega)$$

for each $\tau > 0$. Since $T \in L^2(0, \tau; H^1(\Omega))$ and $\vartheta \in L^2(0, \tau; L^2(\Omega))$, this implies that

$$T \text{ and } \vartheta \in C_b(\mathbf{R}_+; L^2(\Omega)).$$

Since the system (24) is linear, proving uniqueness reduces to proving that the only weak solution of (24) satisfying the initial condition $T^{in} = \vartheta^{in} = 0$ is the trivial solution $T = \vartheta = 0$.

By Lemma A.2, taking $\phi(x) = T(t, x)$ and $\psi(x) = \frac{\sigma}{\sigma'} \vartheta(t, x) / \rho(x)$, one has

$$\begin{aligned} & \langle \partial_t T(t, \cdot), T(t, \cdot) \rangle_{H^1(\Omega)', H^1(\Omega)} + \sigma \int_{\Omega} |\nabla_x T(t, x)|^2 dx \\ & \quad + 4\pi\sigma \int_{\Omega} (\rho(x)T(t, x) - \vartheta(t, x))T(t, x) dx = 0, \\ & \frac{\sigma}{\sigma'} \int_{\Omega} \frac{1}{\rho(x)} \vartheta(t, x) \partial_t \vartheta(t, x) dx + 4\pi\sigma \int_{\Omega} (\vartheta(t, x) - \rho(x)T(t, x)) \frac{\vartheta(t, x)}{\rho(x)} dx = 0. \end{aligned}$$

Adding both sides of the identities above, one finds that

$$\begin{aligned} & \langle \partial_t T(t, \cdot), T(t, \cdot) \rangle_{H^1(\Omega)', H^1(\Omega)} + \frac{\sigma}{\sigma'} \int_{\Omega} \frac{1}{\rho(x)} \vartheta(t, x) \partial_t \vartheta(t, x) dx \\ & \quad + \sigma \int_{\Omega} |\nabla_x T(t, x)|^2 dx = 0. \end{aligned}$$

Integrating both sides of the identity above on $[0, t]$ and applying Lemma A.1 leads to

$$\frac{1}{2} \int_{\Omega} T(t, x)^2 dx + \frac{\sigma}{\sigma'} \int_{\Omega} \frac{1}{\rho(x)} \vartheta(t, x)^2 dx + \sigma \int_0^t \int_{\Omega} |\nabla_x T(s, x)|^2 dx ds = 0$$

so that $T = \vartheta = 0$.

Specializing the variational formulation to $\phi, \psi \in C_c^\infty(\Omega)$ shows that T and ϑ satisfy

$$\begin{cases} \partial_t T - \sigma \Delta_x T + 4\pi\sigma(\rho T - \vartheta) = 0, \\ \partial_t \vartheta + 4\pi\sigma'(\vartheta - \rho T) = 0, \end{cases}$$

in the sense of distributions on $(0, +\infty) \times \Omega$.

Finally, we apply Lemma A.3 to the vector field

$$(t, x) \mapsto ((T(t, x) + \frac{\sigma}{\sigma'} \vartheta(t, x)), -\sigma \nabla_x T(t, x)).$$

Indeed,

$$T + \frac{\sigma}{\sigma'} \vartheta \in C_b([0, +\infty); L^2(\Omega)) \quad \text{and} \quad \nabla_x T \in L^2([0, \tau] \times \Omega)$$

for each $\tau > 0$. By linear combination of the two partial differential equations in (24), one has

$$\partial_t(T + \frac{\sigma}{\sigma'}\vartheta) + \operatorname{div}_x(-\sigma\nabla_x T) = 0$$

in the sense of distributions on $(0, +\infty) \times \Omega$, while

$$\frac{d}{dt} \int_{\Omega} (T(t, x) + \frac{\sigma}{\sigma'}\vartheta(t, x))\phi(x)dx - \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla\phi(x)dx = 0$$

for each $\phi \in H^1(\Omega)$. Therefore

$$\left. \frac{\partial T}{\partial n} \right|_{(0, \tau) \times \partial\Omega} = 0$$

in $H_{00}^{1/2}((0, \tau) \times \partial\Omega)'$ for each $\tau > 0$.

6. PROOF OF THE HOMOGENIZATION LIMIT

Proof of Theorem 2.6. The proof is decomposed in several steps and involves several auxiliary lemmas whose proofs belong to the next section.

Step 1: uniform bounds.

The energy identity for the scaled infinite conductivity problem is

$$\begin{aligned} \frac{1}{2} \int_{A_\epsilon} T_\epsilon(t, x)^2 dx + \frac{1}{2}\epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} T_{i,\epsilon}(t)^2 + \sigma \int_0^t \int_{A_\epsilon} |\nabla_x T_\epsilon(s, x)|^2 dx ds \\ = \frac{1}{2} \int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \frac{1}{2}\epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} |T_{i,\epsilon}^{in}|^2 \end{aligned}$$

for all $t \geq 0$ and $\epsilon > 0$.

As a first consequence of this energy identity, the function $T_\epsilon \in C_b([0, +\infty); \mathcal{H}_N)$ satisfies the bounds

$$\begin{aligned} \|T_\epsilon(t, \cdot)\|_{\mathcal{H}_N}^2 &= \int_{\Omega} \rho(x)C(x)T_\epsilon(t, x)^2 dx \\ &= \rho_A C_A \int_{A_\epsilon} T_\epsilon(t, x)^2 dx + \frac{4\pi}{3}\epsilon^3 \rho_B C_B \sum_{i=1}^N T_{i,\epsilon}(t)^2 \\ &\leq \rho_A C_A \left(\int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} |T_{i,\epsilon}^{in}|^2 \right) \leq \rho_A C_A C^{in} \end{aligned}$$

and

$$\sigma \int_0^t \int_{A_\epsilon} |\nabla_x T_\epsilon(s, x)|^2 dx ds \leq \frac{1}{2} \int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \frac{1}{2}\epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} |T_{i,\epsilon}^{in}|^2 \leq \frac{1}{2} C^{in}$$

since $T_\epsilon(t, x) = T_{i,\epsilon}(t)$ for a.e. $x \in B(x_i, \epsilon)$ and all $i = 1, \dots, N$.

A second consequence of the same energy identity is that

$$\epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} T_{i,\epsilon}(t)^2 \leq \int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} |T_{i,\epsilon}^{in}|^2 \leq C^{in}$$

for all $t \in [0, +\infty)$ and $\epsilon > 0$. To the weak solution T_ϵ of the scaled infinite conductivity problem we associate the empirical measure

$$\mu_\epsilon(t, dx d\theta) := \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \otimes \delta_{T_{i,\epsilon}(t)}, \quad N = 1/\epsilon.$$

Accordingly, we denote

$$\mu_\epsilon^{in}(dx d\theta) := \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \otimes \delta_{T_{i,\epsilon}^{in}}.$$

The estimate above is recast as

$$\iint_{\Omega \times \mathbf{R}} \theta^2 \mu_\epsilon(t, dx d\theta) = \epsilon \sum_{i=1}^N T_{i,\epsilon}(t)^2 \leq \frac{\sigma'}{\sigma} C^{in}.$$

On the other hand, by assumption (30)

$$\iint_{\Omega \times \mathbf{R}} |x|^2 \mu_\epsilon(t, dx d\theta) = \frac{1}{N} \sum_{i=1}^N |x_i|^2 \leq C^{in}.$$

Step 2: compactness properties.

These uniform bounds obviously imply that the family T_ϵ is relatively compact in $L^\infty([0, +\infty); L^2(\Omega))$ weak-* and in $L^2(0, \tau; H^1(\Omega))$ weak for all $\tau > 0$. Likewise the family $(1 + |x|^2 + \theta^2)\mu_\epsilon$ is relatively compact in $L^\infty([0, +\infty); \mathcal{M}_b(\Omega \times \mathbf{R}))$ viewed as the dual of the Banach space² $L^1([0, +\infty); C_0(\Omega \times \mathbf{R}))$ equipped with the weak-* topology.

Henceforth, we denote by (T, μ) a limit point of the family $(T_\epsilon, \mu_\epsilon)$ as $\epsilon \rightarrow 0$. Define

$$\rho(t, \cdot) := \int_{\mathbf{R}} \mu(t, \cdot, d\theta), \quad \vartheta(t, \cdot) := \int_{\mathbf{R}} \theta \mu(t, \cdot, d\theta).$$

Next we return to the energy identity in step 1 recast as follows

$$\begin{aligned} (38) \quad & \frac{1}{2} \int_{\Omega} T_\epsilon(t, x)^2 dx + \left(\frac{\sigma}{\sigma'} - \frac{4}{3}\pi\epsilon^2\right) \iint_{\Omega \times \mathbf{R}} \frac{1}{2} \theta^2 \mu_\epsilon(t, dx d\theta) \\ & + \sigma \int_0^t \int_{A_\epsilon} |\nabla_x T_\epsilon(s, x)|^2 dx ds \\ & = \frac{1}{2} \int_{A_\epsilon} T_\epsilon^{in}(x)^2 dx + \frac{1}{2} \epsilon \sum_{i=1}^N \frac{\sigma}{\sigma'} |T_{i,\epsilon}^{in}|^2, \end{aligned}$$

so that

$$\left(\frac{\sigma}{\sigma'} - \frac{4}{3}\pi\epsilon^2\right) \iint_{\Omega \times \mathbf{R}} \theta^2 \mu_\epsilon(t, dx d\theta) \leq C^{in}.$$

Thus, for each $R > 0$, using $(x, \theta) \mapsto \min(\theta^2, R)$ as test function and the weak-* convergence of the family of measures $(1 + |x|^2 + \theta^2)\mu_\epsilon$, passing to the limit in each side of the inequality above, we get

$$\frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \min(\theta^2, R) \mu(t, dx d\theta) \leq C^{in}.$$

²If X is a locally compact space, the notation $C_0(X)$ designates the set of real-valued continuous functions f defined on X such that f converges to 0 at infinity. This is a Banach space for the norm $\|f\| = \sup_{x \in X} |f(x)|$.

Letting $R \rightarrow +\infty$, by monotone convergence

$$\frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \theta^2 \mu(t, dx d\theta) \leq C^{in}.$$

By the Cauchy-Schwarz inequality

$$\vartheta(t, \cdot)^2 = \left(\int_{\mathbf{R}} \theta \mu(t, \cdot, d\theta) \right)^2 \leq \int_{\mathbf{R}} \mu(t, \cdot, d\theta) \int_{\mathbf{R}} \theta^2 \mu(t, \cdot, d\theta) = \rho \int_{\mathbf{R}} \theta^2 \mu(t, \cdot, d\theta)$$

so that

$$\int_{\Omega} \vartheta(t, x)^2 dx \leq \int_{\Omega} \rho(x) \int_{\mathbf{R}} \theta^2 \mu(t, dx d\theta) \leq \frac{\sigma'}{\sigma} C^{in} \|\rho\|_{L^\infty(\Omega)}.$$

Thus, going back to (38), we conclude that, for each $\tau > 0$,

$$T \in L^\infty([0, +\infty); L^2(\Omega)) \cap L^2(0, \tau; H^1(\Omega)) \quad \text{and} \quad \vartheta \in L^\infty([0, +\infty); L^2(\Omega)).$$

Step 3: passing to the limit in the variational formulation.

Start from the variational formulation of the scaled infinite conductivity problem: for each $\Phi_\epsilon \in \mathcal{V}_N$

$$\begin{aligned} \frac{d}{dt} \left(\int_{A_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx + \frac{3\sigma}{4\pi\sigma'} \frac{1}{\epsilon^2} \int_{B_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx \right) \\ + \sigma \int_{A_\epsilon} \nabla_x T_\epsilon(t, x) \cdot \nabla \Phi_\epsilon(x) dx = 0 \end{aligned}$$

for a.e. $t \in [0, +\infty)$.

Since $T_\epsilon(t, \cdot) \in \mathcal{V}_N$, assuming that $\Phi_\epsilon \in \mathcal{V}_N \cap C_b(\overline{\Omega})$,

$$\frac{3\sigma}{4\pi\sigma'} \frac{1}{\epsilon^2} \int_{B_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx = \frac{\sigma}{\sigma'} \epsilon \sum_{i=1}^N T_{i,\epsilon}(t) \Phi_{i,\epsilon} = \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon(t, dx d\theta).$$

On the other hand

$$\begin{aligned} \int_{A_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx &= \int_{\Omega} T_\epsilon(t, x) \Phi_\epsilon(x) dx - \int_{B_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx \\ &= \int_{\Omega} T_\epsilon(t, x) \Phi_\epsilon(x) dx - \frac{4\pi}{3} \epsilon^3 \sum_{i=1}^N T_{i,\epsilon}(t) \Phi_{i,\epsilon}(t) \\ &= \int_{\Omega} T_\epsilon(t, x) \Phi_\epsilon(x) dx - \frac{4\pi}{3} \epsilon^2 \iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon(t, dx d\theta) \end{aligned}$$

so that

$$\begin{aligned} \left| \int_{\Omega} T_\epsilon(t, x) \Phi_\epsilon(x) dx - \int_{A_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx \right| \\ \leq \frac{2\pi}{3} \epsilon^2 \|\Phi_\epsilon\|_{L^\infty(\Omega)} \iint_{\Omega \times \mathbf{R}} (1 + \theta^2) \mu_\epsilon(t, dx d\theta) \\ \leq \frac{2\pi}{3} \epsilon^2 \|\Phi_\epsilon\|_{L^\infty(\Omega)} (1 + \frac{\sigma'}{\sigma} C^{in}). \end{aligned}$$

Finally

$$\int_{A_\epsilon} \nabla_x T_\epsilon(t, x) \cdot \nabla \Phi_\epsilon(x) dx = \int_{\Omega} \nabla_x T_\epsilon(t, x) \cdot \nabla \Phi_\epsilon(x) dx.$$

We shall pass to the limit in the variational formulation above for two different classes of test functions Φ_ϵ .

Step 4: first class of test functions.

Let $\phi \in C_c^1(\overline{\Omega})$. By the mean value theorem

$$|\phi(x) - \phi(x_i)| \leq \epsilon \|D\phi\|_{L^\infty}$$

so that ϕ “almost” belongs to \mathcal{V}_N — but in general does not belong to \mathcal{V}_N . This difficulty is fixed by the following procedure.

For each $\psi \in C(\overline{B(0, \epsilon)})$, define $\chi[\psi]$ to be the solution of

$$(39) \quad \begin{cases} \Delta \chi[\psi](z) = 0, & \epsilon < |z| < r_\epsilon, \\ \chi[\psi](z) = \psi(z), & |z| \leq \epsilon, \\ \chi[\psi](z) = 0, & |z| = r_\epsilon. \end{cases}$$

Define

$$\mathcal{Q}_\epsilon(x) := \sum_{i=1}^N \chi[\phi(x_i + \cdot) - \phi(x_i)](x - x_i),$$

and let

$$\Phi_\epsilon(x) := \phi(x) - \mathcal{Q}_\epsilon(x).$$

Lemma 6.1. *For each $\epsilon > 0$, one has*

$$\|\mathcal{Q}_\epsilon\|_{L^\infty(\Omega)} \leq 2\|\phi\|_{L^\infty(\Omega)}.$$

Besides

$$\mathcal{Q}_\epsilon \rightarrow 0 \text{ in } H^1(\Omega) \text{ strong}$$

as $\epsilon \rightarrow 0$.

The proof of this lemma is postponed to the end of this section. Taking this for granted, one has

$$\Phi_\epsilon \rightarrow \phi \text{ in } H^1(\Omega) \text{ strong}$$

as $\epsilon \rightarrow 0$. Therefore

$$\begin{aligned} \int_{A_\epsilon} \nabla_x T_\epsilon(t, x) \cdot \nabla \Phi_\epsilon(x) dx &= \int_{\Omega} \nabla_x T_\epsilon(t, x) \cdot \nabla \Phi_\epsilon(x) dx \\ &\rightarrow \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x) dx \text{ weakly in } L^2([0, +\infty)) \end{aligned}$$

as $\epsilon \rightarrow 0$.

On the other hand

$$\begin{aligned} \int_{\Omega} T_\epsilon(t, x) \Phi_\epsilon(x) dx &= \int_{\Omega} T_\epsilon(t, x) \phi(x) dx - \int_{\Omega} T_\epsilon(t, x) \mathcal{Q}_\epsilon(x) dx \\ &\rightarrow \int_{\Omega} T(t, x) \phi(x) dx \text{ in } L^\infty([0, +\infty)) \text{ weak-}^* \end{aligned}$$

as $\epsilon \rightarrow 0$ since

$$\left| \int_{\Omega} T_\epsilon(t, x) \mathcal{Q}_\epsilon(x) dx \right| \leq \|T_\epsilon(t, \cdot)\|_{L^2} \|\mathcal{Q}_\epsilon\|_{L^2}.$$

Indeed

$$\sup_{t \geq 0} \|T_\epsilon(t, \cdot)\|_{L^2} < \infty, \quad \text{while } \|\mathcal{Q}_\epsilon\|_{L^2} \rightarrow 0 \text{ as } \epsilon \rightarrow 0$$

by Lemma 6.1.

Finally

$$\iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon(t, dx d\theta) = \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu_\epsilon(t, dx d\theta)$$

since $\phi(x_i) = \Phi_\epsilon(x_i)$ for $i = 1, \dots, N$, so that

$$\iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon(t, dx d\theta) \rightarrow \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta) \text{ in } L^\infty([0, +\infty)) \text{ weak-}^*$$

as $\epsilon \rightarrow 0$.

By construction $\Phi_\epsilon \in \mathcal{V}_N$, so that Φ_ϵ can be used as a test function in the variational formulation. Passing to the limit in the variational formulation of the scaled infinite heat conductivity problem in the sense of distributions gives

$$\begin{aligned} \frac{d}{dt} \left(\int_{\Omega} T(t, x) \phi(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta) \right) \\ + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x) dx = 0 \end{aligned}$$

in $L^2_{loc}([0, +\infty))$ for each $\phi \in C^1_c(\overline{\Omega})$.

Step 5: second class of test functions

In this step, we shall use a class of test functions $\Psi_\epsilon \in H^1(\Omega)$ such that $\Psi_\epsilon|_{B(x_i, \epsilon)} = 0$ for all $i = 1, \dots, N$. Given $\phi \in C^1_c(\overline{\Omega})$, define Ψ_ϵ as follows:

$$\Psi_\epsilon(x) := \phi(x) - \mathcal{P}_\epsilon(x)$$

where

$$\mathcal{P}_\epsilon(x) := \sum_{i=1}^N \chi[\phi(x_i + \cdot)](x - x_i).$$

We shall further decompose \mathcal{P}_ϵ as follows:

$$\begin{aligned} \mathcal{P}_\epsilon(x) &= \sum_{i=1}^N \chi[\phi(x_i + \cdot) - \phi(x_i)](x - x_i) + \sum_{i=1}^N \chi[\phi(x_i)](x - x_i) \\ &= \mathcal{Q}_\epsilon(x) + \mathcal{R}_\epsilon(x). \end{aligned}$$

Likewise, one associates to the solution T_ϵ of the scaled infinite heat conductivity problem

$$\Theta_\epsilon(t, x) := T_\epsilon(t, x) - \mathcal{S}_\epsilon(t, x)$$

where

$$\mathcal{S}_\epsilon(t, x) := \sum_{i=1}^N \chi[T_{i, \epsilon}(t)](x - x_i).$$

The variational formulation for the test function Ψ_ϵ becomes

$$\frac{d}{dt} \int_{\Omega} T_\epsilon(t, x) \Psi_\epsilon(x) dx + \sigma \int_{\Omega} \nabla_x T_\epsilon(t, x) \cdot \nabla \Psi_\epsilon(x) dx = 0$$

in $L^2_{loc}([0, +\infty))$, since $\Psi_\epsilon = 0$ on $B(x_i, \epsilon)$ for all $i = 1, \dots, N$.

Lemma 6.2. *One has*

$$\mathcal{R}_\epsilon \rightarrow 0 \quad \text{in } H^1(\Omega) \text{ weak,}$$

so that

$$\mathcal{P}_\epsilon \rightarrow 0 \text{ in } H^1(\Omega) \text{ weak,}$$

while

$$\mathcal{S}_\epsilon \rightarrow 0 \quad \text{in } L^\infty([0, +\infty); H^1(\Omega)) \text{ weak-}^*$$

as $\epsilon \rightarrow 0$.

Taking this lemma for granted, and observing that

$$\text{supp}(\mathcal{P}_\epsilon) \subset \text{supp}(\phi) + \overline{B(0, r_\epsilon)}$$

the Rellich compactness theorem implies that

$$\mathcal{P}_\epsilon \rightarrow 0 \quad \text{in } L^2(\Omega) \text{ strong}$$

as $\epsilon \rightarrow 0$, so that

$$\int_{\Omega} T_\epsilon(t, x) \Psi_\epsilon(x) dx \rightarrow \int_{\Omega} T(t, x) \phi(x) dx \text{ in } L^\infty([0, +\infty)) \text{ weak-}^*$$

as $\epsilon \rightarrow 0$.

Next, decompose

$$\begin{aligned} \int_{\Omega} \nabla_x T_\epsilon(t, x) \cdot \nabla \Psi_\epsilon(x) dx &= \int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \phi(x) dx \\ &\quad - \int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \mathcal{P}_\epsilon(x) dx \\ &\quad + \int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \Psi_\epsilon(x) dx \end{aligned}$$

Since $\mathcal{S}_\epsilon \rightarrow 0$ in $L^2(0, \tau; H_0^1(\Omega))$ weak as $\epsilon \rightarrow 0$, one has

$$\nabla_x \Theta_\epsilon = \nabla_x T_\epsilon - \nabla_x \mathcal{S}_\epsilon \rightarrow \nabla_x T \quad \text{in } L^2([0, \tau] \times \Omega) \text{ weak}$$

as $\epsilon \rightarrow 0$, so that

$$\int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \phi(x) dx \rightarrow \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x) dx \text{ in } L^2([0, \tau]) \text{ weak}$$

as $\epsilon \rightarrow 0$.

Furthermore, one has

$$\begin{aligned} \int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \Psi_\epsilon(x) dx &= \int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \phi(x) dx - \int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \mathcal{Q}_\epsilon(x) dx \\ &\quad - \sum_{i=1}^N \int_{B(x_i, r_\epsilon) \setminus B(x_i, \epsilon)} \nabla \chi[T_{i, \epsilon}(t)](z) \cdot \nabla \chi[\phi(x_i)](z) dz. \end{aligned}$$

As noticed above, $\nabla_x \mathcal{S}_\epsilon \rightarrow 0$ weakly in $L^2([0, \tau] \times \Omega)$ for all $\tau > 0$ as $\epsilon \rightarrow 0$, and therefore

$$\int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \phi(x) dx \rightarrow 0 \quad \text{in } L^2([0, \tau]) \text{ weak}$$

for all $\tau > 0$ as $\epsilon \rightarrow 0$, while

$$\int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \mathcal{Q}_\epsilon(x) dx \rightarrow 0 \quad \text{in } L^2([0, \tau]) \text{ strong}$$

for all $\tau > 0$ as $\epsilon \rightarrow 0$ by Lemma 6.1.

The third term on the right hand side of the last equality is handled with the following lemma.

Lemma 6.3. *One has*

$$\sum_{i=1}^N \int_{B(0, r_\epsilon) \setminus B(0, \epsilon)} \nabla \chi[T_{i, \epsilon}(t)](z) \cdot \nabla \chi[\phi(x_i)](z) dz \rightarrow 4\pi \int_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta)$$

in $L^\infty(\mathbf{R}_+)$ weak- * .

Therefore

$$\int_{\Omega} \nabla_x \mathcal{S}_\epsilon(t, x) \cdot \nabla \Psi_\epsilon(x) dx \rightarrow -4\pi \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta)$$

in $L^2([0, \tau])$ weak as $\epsilon \rightarrow 0$.

It remains to treat the term

$$\begin{aligned} \int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \mathcal{P}_\epsilon(x) dx &= \int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \mathcal{Q}_\epsilon(x) dx \\ &+ \sum_{i=1}^N \int_{B(x_i, r_\epsilon) \setminus B(x_i, \epsilon)} \nabla_x \Theta_\epsilon(t, z) \cdot \nabla \chi[\phi(x_i)](z) dz \end{aligned}$$

By Lemma 6.1, $\mathcal{Q}_\epsilon \rightarrow 0$ in $H^1(\Omega)$ strong; by the second convergence in Lemma 6.2, the family $\nabla_x \mathcal{S}_\epsilon$ is bounded in $L^2([0, \tau] \times \Omega)$ for each $\tau > 0$, while $\nabla_x \mathcal{T}_\epsilon$ is bounded in $L^2([0, \tau] \times \Omega)$ as explained in Step 1. Thus $\nabla_x \Theta_\epsilon$ is bounded in $L^2([0, \tau] \times \Omega)$ for all $\tau > 0$, so that

$$\int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \mathcal{Q}_\epsilon(x) dx \rightarrow 0 \text{ in } L^2([0, \tau]) \text{ strong}$$

for each $\tau > 0$ as $\epsilon \rightarrow 0$.

Next, by Green's formula

$$\int_{B(x_i, r_\epsilon) \setminus B(x_i, \epsilon)} \nabla_x \Theta_\epsilon(t, z) \cdot \nabla \chi[\phi(x_i)](z) dz = \int_{\partial B(x_i, r_\epsilon)} \Theta_\epsilon(t, z) \frac{\partial \chi[\phi(x_i)]}{\partial n}(z) dz$$

since $\chi[\phi(x_i)]$ is harmonic on $B(x_i, r_\epsilon) \setminus B(x_i, \epsilon)$ and $\Theta_\epsilon|_{\partial B(x_i, \epsilon)} = 0$.

Lemma 6.4. *For each $\phi \in C_b(\mathbf{R}^3)$, one has*

$$\sum_{i=1}^N \frac{\partial}{\partial n} (\chi[\phi(x_i)](x - x_i)) \delta_{\partial B(x_i, r_\epsilon)} = -\frac{\epsilon r_\epsilon}{r_\epsilon^2 (r_\epsilon - \epsilon)} \sum_{i=1}^N \phi(x_i) \delta_{\partial B(x_i, r_\epsilon)} \rightarrow -4\pi \rho \phi$$

in $H^{-1}(\mathbf{R}^3)$. We recall that $\rho \in C_b(\overline{\Omega})$ is defined as follows:

$$\frac{1}{N} \sum_{i=1}^N \delta_{x_i} \rightarrow \rho \mathcal{L}^3$$

weakly in the sense of probability measures on $\overline{\Omega}$, where \mathcal{L}^3 designates the 3-dimensional Lebesgue measure.

Taking this lemma for granted, we see that

$$\int_{\Omega} \nabla_x \Theta_\epsilon(t, x) \cdot \nabla \mathcal{R}_\epsilon(x) dx \rightarrow -4\pi \int_{\Omega} \rho(x) T(t, x) \phi(x) dx \text{ in } L^2([0, \tau]) \text{ weak}$$

for each $\tau > 0$ as $\epsilon \rightarrow 0$.

Summarizing the various limits established in this step, we conclude that, for each $\phi \in C_c^1(\overline{\Omega})$

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} T(t, x) \phi(x) dx + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x) dx + 4\pi \sigma \int_{\Omega} \rho(x) T(t, x) \phi(x) dx \\ - 4\pi \sigma \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta) = 0 \end{aligned}$$

in $L^2_{loc}([0, +\infty))$.

Step 6: initial conditions

As explained in steps 3-4, for each $\phi \in C_c^1(\overline{\Omega})$, defining

$$\Phi_\epsilon = \phi - \mathcal{Q}_\epsilon \in \mathcal{V}_N$$

one has

$$(40) \quad \begin{aligned} \int_{A_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon(t, dx d\theta) \\ \rightarrow \int_{\Omega} T(t, x) \phi(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta) \end{aligned}$$

in $L^2([0, \tau])$ weak as $\epsilon \rightarrow 0$, while

$$\begin{aligned} \int_0^\infty \left| \frac{d}{dt} \left(\int_{A_\epsilon} T_\epsilon(t, x) \Phi_\epsilon(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon(t, dx d\theta) \right) \right|^2 dt \\ \leq \sigma^2 \|\nabla \Phi_\epsilon\|_{L^2(\Omega)}^2 \int_0^\infty \|\nabla_x T_\epsilon(t, \cdot)\|_{L^2(\Omega)}^2 dt \\ \leq C^{in} \sigma (\|\nabla \phi\|_{L^2(\Omega)} + o(1))^2 \end{aligned}$$

By Ascoli-Arzelà's theorem, the convergence in (40) is uniform on $[0, \tau]$ for each τ .

In particular

$$\begin{aligned} \int_{A_\epsilon} T_\epsilon^{in}(x) \Phi_\epsilon(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \Phi_\epsilon(x) \theta \mu_\epsilon^{in}(dx d\theta) \\ \rightarrow \int_{\Omega} T^{in}(x) \phi(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu^{in}(dx d\theta) \\ = \int_{\Omega} T(0, x) \phi(x) dx + \frac{\sigma}{\sigma'} \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(0, dx d\theta) \end{aligned}$$

for each $\phi \in C_c^1(\overline{\Omega})$, so that

$$T(0, \cdot) + \frac{\sigma'}{\sigma} \int_{\mathbf{R}} \theta \mu(0, \cdot, d\theta) = T^{in} + \frac{\sigma'}{\sigma} \int_{\mathbf{R}} \theta \mu^{in}(\cdot, d\theta).$$

Likewise, we have seen in step 5 that, for each $\phi \in C_c^1(\overline{\Omega})$, defining Ψ_ϵ as

$$\Psi_\epsilon = \phi - \mathcal{P}_\epsilon,$$

one has

$$(41) \quad \int_{\Omega} T_\epsilon(t, x) \Psi_\epsilon(x) dx \rightarrow \int_{\Omega} T(t, x) \phi(x) dx \text{ in } L^\infty([0, +\infty)) \text{ weak-}^*$$

as $\epsilon \rightarrow 0$. Besides

$$\begin{aligned} \int_0^\infty \left| \frac{d}{dt} \int_{\Omega} T_\epsilon(t, x) \Psi_\epsilon(x) dx \right|^2 dt \leq \sigma^2 \|\nabla \Psi_\epsilon\|_{L^2(\Omega)}^2 \int_0^\infty \int_{\Omega} |\nabla_x T_\epsilon(t, x)|^2 dx dt \\ \leq C^{in} \sigma (\|\nabla \phi\|_{L^2(\Omega)} + o(1))^2 \end{aligned}$$

By the Ascoli-Arzelà theorem, the convergence in (41) is uniform in $[0, \tau]$ for each $\tau > 0$. In particular

$$\int_{\Omega} T_\epsilon^{in}(x) \Psi_\epsilon(x) dx \rightarrow \int_{\Omega} T^{in}(x) \phi(x) dx = \int_{\Omega} T(0, x) \phi(x) dx$$

so that

$$T(0, \cdot) = T^{in}.$$

Step 7: identification of the limiting system

In steps 4-5, we have proved that

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} T(t, x) \Phi(x) dx + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \Phi(x) dx \\ + 4\pi\sigma \int_{\Omega} (\rho(x)T(t, x) - \vartheta(t, x)) \Phi(x) dx = 0 \end{aligned}$$

and

$$\begin{aligned} \frac{d}{dt} \left(\int_{\Omega} T(t, x) \Phi(x) dx + \frac{\sigma}{\sigma'} \int_{\Omega} \vartheta(t, x) \Phi(x) dx \right) \\ + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \Phi(x) dx = 0 \end{aligned}$$

for each $\Phi \in C_c^1(\overline{\Omega})$. By linear combination, one finds that

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} T(t, x) \phi(x) dx + \sigma \int_{\Omega} \nabla_x T(t, x) \cdot \nabla \phi(x) dx \\ + 4\pi\sigma \int_{\Omega} (\rho(x)T(t, x) - \vartheta(t, x)) \phi(x) dx = 0 \\ \frac{d}{dt} \int_{\Omega} \vartheta(t, x) \psi(x) dx + 4\pi\sigma' \int_{\Omega} (\vartheta(t, x) - \rho(x)T(t, x)) \psi(x) dx = 0 \end{aligned}$$

for all $\phi, \psi \in C_c^1(\overline{\Omega})$. Since $T, \vartheta \in C_b([0, +\infty); L^2(\Omega))$ and $T \in L^2(0, \tau; H^1(\Omega))$ for each $\tau > 0$, the identities above hold for all $\phi, \psi \in H^1(\Omega)$ by a straightforward density argument.

Thus (T, ϑ) is the unique weak solution of (24) with initial data (T^{in}, ϑ^{in}) .

By compactness, this implies that

$$T_{\epsilon} \rightarrow T \text{ in } L^{\infty}([0, +\infty); L^2(\Omega)) \text{ weak-}^* \text{ and in } L^2(0, \tau; H^1(\Omega)) \text{ weak,}$$

while

$$\vartheta_{\epsilon} \rightarrow \vartheta \text{ in } L^{\infty}([0, +\infty); L^2(\Omega)) \text{ weak-}^*$$

without extracting subsequences. \square

7. PROOF OF LEMMAS 6.1, 6.2, 6.3 AND 6.4

When $\psi = 1$, the solution of the boundary value problem (39) is given by

$$\chi[1](z) = \frac{\epsilon r_{\epsilon}}{r_{\epsilon} - \epsilon} \left(\frac{1}{|z|} - \frac{1}{r_{\epsilon}} \right) \mathbf{1}_{B(0, r_{\epsilon}) \setminus B(0, \epsilon)}(z) + \mathbf{1}_{B(0, \epsilon)}(z)$$

for all $z \in \mathbf{R}^3$. In that case

$$\|\chi[1]\|_{L^2(\mathbf{R}^3)}^2 = \frac{4\pi}{3} \epsilon^2 r_{\epsilon}, \quad \|\nabla \chi[1]\|_{L^2(\mathbf{R}^3)}^2 = 4\pi \frac{\epsilon r_{\epsilon}}{r_{\epsilon} - \epsilon} \sim 4\pi \epsilon \quad \text{as } \epsilon \rightarrow 0.$$

Proof of Lemma 6.1. First, by the maximum principle and the mean value theorem, one has

$$\begin{aligned} \|\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^{\infty}(\mathbf{R}^3)} &\leq \|\phi(x_i + \cdot) - \phi(x_i)\|_{L^{\infty}(\mathbf{R}^3)} \\ &\leq \min(2\|\phi\|_{L^{\infty}(\mathbf{R}^3)}, \|\nabla \phi\|_{L^{\infty}(\mathbf{R}^3)} \epsilon). \end{aligned}$$

Since the functions $x \mapsto \chi[\phi(x_i + \cdot) - \phi(x_i)](x - x_i)$ have disjoint supports by (32), one has both

$$\|\mathcal{Q}_{\epsilon}\|_{L^{\infty}(\Omega)} \leq \sup_{1 \leq i \leq N} \|\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^{\infty}(\mathbf{R}^3)} \leq 2\|\phi\|_{L^{\infty}(\mathbf{R}^3)},$$

and

$$\begin{aligned} \|\mathcal{Q}_\epsilon\|_{L^2(\Omega)}^2 &\leq \sum_{i=1}^N \|\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(\mathbf{R}^3)}^2 \\ &\leq N|B(0, r_\epsilon)| \|\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^\infty(\mathbf{R}^3)}^2 \\ &\leq N \cdot \frac{4}{3}\pi r_\epsilon^3 \|\nabla\phi\|_{L^\infty(\mathbf{R}^3)}^2 \epsilon^2 = \frac{4}{3}\pi \|\nabla\phi\|_{L^\infty(\mathbf{R}^3)}^2 \epsilon^2 \rightarrow 0 \end{aligned}$$

as $\epsilon \rightarrow 0$. Next

$$\begin{aligned} \|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(\mathbf{R}^3)}^2 &= \|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(B(0, \epsilon))}^2 \\ &\quad + \|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(B(0, r_\epsilon) \setminus B(0, \epsilon))}^2. \end{aligned}$$

First

$$\begin{aligned} \|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(B(0, \epsilon))}^2 &\leq \frac{4}{3}\pi\epsilon^3 \|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^\infty(B(0, \epsilon))}^2 \\ &= \frac{4}{3}\pi\epsilon^3 \|\nabla\phi\|_{L^\infty(B(0, \epsilon))}^2. \end{aligned}$$

Since $\chi[\phi(x_i + \cdot) - \phi(x_i)]$ is a harmonic function on $B(0, r_\epsilon) \setminus B(0, \epsilon)$, it minimizes the Dirichlet integral among functions with the same boundary values. Thus

$$\|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(B(0, r_\epsilon) \setminus B(0, \epsilon))}^2 \leq \|\nabla\chi_{i, \epsilon}\|_{L^2(B(0, r_\epsilon) \setminus B(0, \epsilon))}^2$$

where

$$\chi_{i, \epsilon}(z) = \left(\phi\left(x_i + \epsilon \frac{z}{|z|}\right) - \phi(x_i) \right) \frac{r_\epsilon - |z|}{r_\epsilon - \epsilon}.$$

Straightforward computations show that

$$\begin{aligned} \nabla\chi_{i, \epsilon}(z) &= \left(I - \frac{z \otimes z}{|z|^2} \right) \nabla\phi\left(x_i + \epsilon \frac{z}{|z|}\right) \frac{\epsilon}{|z|} \frac{r_\epsilon - |z|}{r_\epsilon - \epsilon} \\ &\quad - \left(\phi\left(x_i + \epsilon \frac{z}{|z|}\right) - \phi(x_i) \right) \frac{1}{r_\epsilon - \epsilon} \frac{z}{|z|}, \end{aligned}$$

so that

$$\begin{aligned} |\nabla\chi_{i, \epsilon}(z)|^2 &\leq \left| \nabla\phi\left(x_i + \epsilon \frac{z}{|z|}\right) \right|^2 \frac{\epsilon^2}{|z|^2} \frac{(r_\epsilon - |z|)^2}{(r_\epsilon - \epsilon)^2} \\ &\quad + \left(\phi\left(x_i + \epsilon \frac{z}{|z|}\right) - \phi(x_i) \right)^2 \frac{1}{(r_\epsilon - \epsilon)^2}. \end{aligned}$$

Thus

$$\|\nabla\chi_{i, \epsilon}\|_{L^2(B(0, r_\epsilon) \setminus B(0, \epsilon))}^2 \leq \frac{8\pi}{3} \|\nabla\phi\|_{L^\infty(\mathbf{R}^3)}^2 \epsilon^2 r_\epsilon + O(\epsilon^3 r_\epsilon),$$

so that

$$\begin{aligned} \|\nabla\mathcal{Q}_\epsilon\|_{L^2(\Omega)}^2 &\leq \sum_{i=1}^N \|\nabla\chi[\phi(x_i + \cdot) - \phi(x_i)]\|_{L^2(\mathbf{R}^3)}^2 \\ &\leq \frac{4\pi}{3} \|\nabla\phi\|_{L^\infty(\mathbf{R}^3)}^2 N(\epsilon^3 + 2\epsilon^2 r_\epsilon) \\ &= \frac{4\pi}{3} \|\nabla\phi\|_{L^\infty(\mathbf{R}^3)}^2 (\epsilon^2 + 2\epsilon r_\epsilon) \rightarrow 0 \end{aligned}$$

as $\epsilon \rightarrow 0$. Hence $\mathcal{Q}_\epsilon \rightarrow 0$ in $H^1(\Omega)$ strong as $\epsilon \rightarrow 0$. □

Proof of Lemma 6.2. Assume that $0 < \epsilon < \frac{1}{8}$. Since the functions $x \mapsto \chi[1](x - x_i)$ have disjoint supports by (32), one has

$$\begin{aligned} \|\mathcal{S}_\epsilon(t, \cdot)\|_{L^2(\Omega)}^2 &\leq \sum_{i=1}^N T_{i,\epsilon}(t)^2 \|\chi[1]\|_{L^2(\mathbf{R}^3)}^2 \\ \|\nabla \mathcal{S}_\epsilon(t, \cdot)\|_{L^2(\Omega)}^2 &\leq \sum_{i=1}^N T_{i,\epsilon}(t)^2 \|\nabla \chi[1]\|_{L^2(\mathbf{R}^3)}^2. \end{aligned}$$

Thus

$$\|\mathcal{S}_\epsilon(t, \cdot)\|_{L^2(\Omega)}^2 \leq \frac{4\pi}{3} \epsilon^2 r_\epsilon \sum_{i=1}^N T_{i,\epsilon}(t)^2 \leq \frac{4\pi}{3} \frac{\sigma'}{\sigma} C^{in} \epsilon r_\epsilon \rightarrow 0$$

as $\epsilon \rightarrow 0$, while

$$\begin{aligned} \|\nabla \mathcal{S}_\epsilon(t, \cdot)\|_{L^2(\Omega)}^2 &\leq 4\pi \frac{\epsilon r_\epsilon}{r_\epsilon - \epsilon} \sum_{i=1}^N T_{i,\epsilon}(t)^2 \\ &\leq 8\pi \epsilon \sum_{i=1}^N T_{i,\epsilon}(t)^2 \leq 8\pi \frac{\sigma'}{\sigma} C^{in}. \end{aligned}$$

Hence $\mathcal{S}_\epsilon(t, \cdot) \rightarrow 0$ in $H^1(\Omega)$ weak, uniformly in $t \geq 0$ as $\epsilon \rightarrow 0$.

Now for \mathcal{R}_ϵ . First

$$\begin{aligned} \|\mathcal{R}_\epsilon\|_{L^2(\Omega)}^2 &= \sum_{i=1}^N \|\chi[\phi(x_i)]\|_{L^2(\mathbf{R}^3)}^2 \\ &\leq \sum_{i=1}^N \frac{4}{3} \pi \phi(x_i)^2 \epsilon^2 r_\epsilon \leq \frac{4}{3} \pi \|\phi\|_{L^\infty(\mathbf{R}^3)}^2 \epsilon r_\epsilon \rightarrow 0 \end{aligned}$$

as $\epsilon \rightarrow 0$, because the functions $x \mapsto \chi[1](x - x_i)$ have disjoint supports by (32). By the same token

$$\begin{aligned} \|\nabla \mathcal{R}_\epsilon\|_{L^2(\Omega)}^2 &= \sum_{i=1}^N \|\nabla \chi[\phi(x_i)]\|_{L^2(\mathbf{R}^3)}^2 \\ &\leq \sum_{i=1}^N 4\pi \phi(x_i)^2 \frac{\epsilon r_\epsilon}{r_\epsilon - \epsilon} \leq 4\pi \|\phi\|_{L^\infty(\mathbf{R}^3)}^2 \frac{N \epsilon r_\epsilon}{r_\epsilon - \epsilon} = O(1) \end{aligned}$$

as $\epsilon \rightarrow 0$. Thus $\mathcal{R}_\epsilon \rightarrow 0$ in $H^1(\Omega)$ weak as $\epsilon \rightarrow 0$. \square

Proof of Lemma 6.3. One has

$$\begin{aligned}
& \sum_{i=1}^N \int_{B(0,r_\epsilon) \setminus B(0,\epsilon)} \nabla \chi[T_{i,\epsilon}(t)](z) \cdot \nabla \chi[\phi(x_i)](z) dz \\
&= \|\nabla \chi[1]\|_{L^2(\mathbf{R}^N)}^2 \sum_{i=1}^N T_{i,\epsilon}(t) \phi(x_i) \\
&= 4\pi \frac{\epsilon r_\epsilon}{r_\epsilon - \epsilon} \sum_{i=1}^N T_{i,\epsilon}(t) \phi(x_i) \\
&= 4\pi \frac{r_\epsilon}{r_\epsilon - \epsilon} \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu_\epsilon(t, dx d\theta) \\
&\rightarrow 4\pi \iint_{\Omega \times \mathbf{R}} \phi(x) \theta \mu(t, dx d\theta) = 4\pi \int_{\Omega} \phi(x) \vartheta(t, x) dx
\end{aligned}$$

as $\epsilon \rightarrow 0$. □

Proof of Lemma 6.4. First

$$\begin{aligned}
& \sum_{i=1}^N \frac{\partial}{\partial n} (\chi[\phi(x_i)](x - x_i)) \delta_{\partial B(x_i, r_\epsilon)} \\
&= \sum_{i=1}^N \phi(x_i) \frac{x - x_i}{|x - x_i|} \cdot \nabla \chi[1](x - x_i) \delta_{\partial B(x_i, r_\epsilon)} \\
&= - \sum_{i=1}^N \phi(x_i) \frac{\epsilon r_\epsilon}{r_\epsilon - \epsilon} \frac{1}{|x - x_i|^2} \delta_{\partial B(x_i, r_\epsilon)} \\
&= - \frac{\epsilon r_\epsilon}{r_\epsilon^2 (r_\epsilon - \epsilon)} \sum_{i=1}^N \phi(x_i) \delta_{\partial B(x_i, r_\epsilon)}.
\end{aligned}$$

Next we recall that

$$\sum_{i=1}^N \phi(x_i) r_\epsilon \delta_{\partial B(x_i, r_\epsilon)} \rightarrow 4\pi \rho \phi \quad \text{strongly in } H^{-1}(\mathbf{R}^3)$$

as $\epsilon \rightarrow 0$. This result has been proved by Cioranescu-Murat [4] in the case where x_i are distributed periodically; see formula (64) and Appendix 1 in [5] for a proof adapted to the setting of the present paper.

With the explicit formula above and the fact that $\frac{\epsilon}{r_\epsilon^2 (r_\epsilon - \epsilon)} \rightarrow 1$ as $\epsilon \rightarrow 0$, this concludes the proof of Lemma 6.4. □

APPENDIX A. SOME LEMMAS ON EVOLUTION EQUATIONS

Let \mathcal{V} and \mathcal{H} be two separable Hilbert spaces such that $\mathcal{V} \subset \mathcal{H}$ with continuous inclusion and \mathcal{V} is dense in \mathcal{H} . The Hilbert space \mathcal{H} is identified with its dual and the map

$$\mathcal{H} \ni u \mapsto L_u \in \mathcal{V}',$$

where L_u is the linear functional

$$L_u : \mathcal{V} \ni v \mapsto (u|v)_{\mathcal{H}} \in \mathbf{R},$$

identifies \mathcal{H} with a dense subspace of \mathcal{V}' .

Lemma A.1. *Assume that*

$$v \in L^2(0, T; \mathcal{V}) \quad \text{and} \quad \frac{dL_v}{dt} \in L^2(0, T; \mathcal{V}').$$

Then

- a) *the function v is a.e. equal to a unique element of $C([0, T], \mathcal{H})$ still denoted v ;*
b) *this function $v \in C([0, T], \mathcal{H})$ satisfies*

$$\frac{1}{2}|v(t_2)|_{\mathcal{H}}^2 - \frac{1}{2}|v(t_1)|_{\mathcal{H}}^2 = \int_{t_1}^{t_2} \left\langle \frac{dL_v}{dt}(t), v(t) \right\rangle_{\mathcal{V}', \mathcal{V}} dt$$

for all $t_1, t_2 \in [0, T]$

Statement a) follows from Proposition 2.1 and Theorem 3.1 in chapter 1 of [8], and statement b) from Theorem II.5.12 of [1].

Lemma A.2. *Let $L \in L^2(0, T; \mathcal{V}')$ satisfy*

$$\langle L(t), w \rangle_{\mathcal{V}', \mathcal{V}} = 0 \quad \text{for a.e. } t \in [0, T]$$

for all $w \in \mathcal{V}$. Then

$$L(t) = 0 \quad \text{for a.e. } t \in [0, T].$$

Proof. Pick $\mathcal{N}_w \subset [0, T]$ negligible such that L is defined on $[0, T] \setminus \mathcal{N}_w$ and

$$\langle L(t), w \rangle_{\mathcal{V}', \mathcal{V}} = 0 \quad \text{for all } t \in [0, T] \setminus \mathcal{N}_w.$$

Let \mathcal{D} be a dense countable subset of \mathcal{V} and let

$$\tilde{\mathcal{N}} := \bigcup_{w \in \mathcal{D}} \mathcal{N}_w.$$

For all $t \in [0, T] \setminus \tilde{\mathcal{N}}$, one has

$$\langle L(t), w \rangle_{\mathcal{V}', \mathcal{V}} = 0 \quad \text{for all } w \in \mathcal{D} \quad \text{so that } L(t) = 0$$

because $L(t)$ is a continuous linear functional on \mathcal{V} and \mathcal{D} is dense in \mathcal{V} . \square

The next lemma recalls the functional background for Green's formula in the context of evolution equations.

Lemma A.3. *Let Ω be an open subset of \mathbf{R}^N with smooth boundary, and let $T > 0$. Denote by n the unit outward normal field on $\partial\Omega$. Let $\rho \in C([0, T]; L^2(\Omega))$ and $m \in L^2((0, T) \times \Omega, \mathbf{R}^N)$. Assume that*

$$\partial_t \rho + \operatorname{div}_x m = 0 \quad \text{in the sense of distributions in } (0, T) \times \Omega.$$

Then

- a) *the vector field m has a normal trace $m \cdot n|_{(0, T) \times \partial\Omega} \in H_{00}^{1/2}((0, T) \times \partial\Omega)'$;*
b) *for each $\psi \in H^1(\Omega)$*

$$\begin{aligned} \frac{d}{dt} \int_{\Omega} \rho(\cdot, x) \psi(x) dx - \int_{\Omega} m(\cdot, x) \cdot \nabla_x \psi(x) dx \\ = - \langle m \cdot n|_{\partial\Omega}, \psi|_{\partial\Omega} \rangle_{H^{-1/2}(\partial\Omega), H^{1/2}(\partial\Omega)} \end{aligned}$$

in $H^{-1}(0, T)$.

Proof. Let $\chi \in C_c^\infty(\mathbf{R})$ be such that

$$\chi(t) = 1 \text{ for } t \in [-1, T+1] \quad \text{and} \quad \text{supp}(\chi) \subset [-2, T+2].$$

Define

$$\bar{\rho}(t, x) := \begin{cases} \rho(t, x) & \text{if } 0 \leq t \leq T \\ \chi(t)\rho(0, x) & \text{if } t < 0 \\ \chi(t)\rho(T, x) & \text{if } t > T \end{cases}$$

and

$$\bar{m}(t, x) := \begin{cases} m(t, x) & \text{if } 0 \leq t \leq T \\ 0 & \text{if } t \notin [0, T] \end{cases}$$

so that the vector field $X := (\bar{\rho}, \bar{m})$ is an extension of (ρ, m) to $\mathbf{R} \times \Omega$ satisfying

$$X \in L^2(\mathbf{R} \times \Omega; \mathbf{R}^{N+1}).$$

Besides

$$(\partial_t \bar{\rho} + \text{div}_x \bar{m})(t, x) = \chi'(t)(\mathbf{1}_{t < 0}\rho(0, x) + \mathbf{1}_{t > T}\rho(T, x)) =: S(t, x)$$

with $S \in L^2(\mathbf{R} \times \Omega)$ so that

$$\text{div}_{t,x} X = S \in L^2(\mathbf{R} \times \Omega).$$

Therefore X has a normal trace on the boundary $\partial(\mathbf{R} \times \Omega) = \mathbf{R} \times \partial\Omega$, denoted $X \cdot n|_{\mathbf{R} \times \partial\Omega} \in H^{-1/2}(\mathbf{R} \times \partial\Omega)$.

Let $\phi \in H_{00}^{1/2}((0, T) \times \partial\Omega)$; denote by $\bar{\phi}$ its extension by 0 to $\mathbf{R} \times \partial\Omega$. Thus $\bar{\phi} \in H^{1/2}(\mathbf{R} \times \partial\Omega)$ and there exists $\bar{\Phi} \in H^1(\mathbf{R} \times \Omega)$ such that $\bar{\phi} = \bar{\Phi}|_{\mathbf{R} \times \partial\Omega}$. The normal trace of m is then defined as follows: by Green's formula

$$\begin{aligned} & \langle m \cdot n|_{\mathbf{R} \times \partial\Omega}, \phi \rangle_{H_{00}^{1/2}((0, T) \times \partial\Omega)', H_{00}^{1/2}((0, T) \times \partial\Omega)} \\ & := \langle X \cdot n|_{\mathbf{R} \times \partial\Omega}, \bar{\phi} \rangle_{H^{1/2}(\mathbf{R} \times \partial\Omega)', H^{1/2}(\mathbf{R} \times \partial\Omega)} \\ & = \iint_{\mathbf{R} \times \Omega} (\bar{\rho} \partial_t \bar{\Phi} + \bar{m} \cdot \nabla_x \bar{\Phi} + S \bar{\Phi})(t, x) dx dt. \end{aligned}$$

Applying Green's formula on $(0, T) \times \Omega$ shows that two different extensions of the vector field (ρ, m) define the same distribution $m \cdot n|_{(0, T) \times \partial\Omega}$ on $(0, T) \times \partial\Omega$. This completes the proof of statement a).

As for statement b), let $\kappa \in H_0^1(0, T)$ and $\psi \in H^1(\Omega)$, define $\Phi(t, x) := \kappa(t)\psi(x)$ and let $\bar{\Phi}$ be the extension of Φ by 0 to $\mathbf{R} \times \Omega$, so that $\bar{\Phi} \in H^1(\mathbf{R} \times \Omega)$. Thus

$$\begin{aligned}
\phi &= \Phi|_{(0,T)\times\partial\Omega} \in H_{00}^{1/2}((0,T)\times\partial\Omega) \text{ and} \\
&\left\langle \langle m \cdot n|_{\partial\Omega}, \psi|_{\partial\Omega} \rangle_{H^{-1/2}(\partial\Omega), H^{1/2}(\partial\Omega)}, \kappa \right\rangle_{H^{-1}(0,T), H_0^1(0,T)} \\
&:= \langle m \cdot n|_{(0,T)\times\partial\Omega}, \phi \rangle_{H_{00}^{1/2}((0,T)\times\partial\Omega)', H_{00}^{1/2}((0,T)\times\partial\Omega)} \\
&= \iint_{\mathbf{R}\times\Omega} (\bar{\rho}\partial_t\bar{\Phi} + \bar{m} \cdot \nabla_x\bar{\Phi} + S\bar{\Phi})(t, x) dx dt \\
&= \int_0^T \int_{\Omega} (\rho(t, x)\kappa'(t)\psi(x) + m(t, x) \cdot \nabla\psi(x)\kappa(t)) dx dt \\
&= -\left\langle \frac{d}{dt} \int_{\Omega} \rho(t, x)\psi(x) dx, \kappa \right\rangle_{H^{-1}(0,T), H_0^1(0,T)} \\
&\quad + \int_0^T \int_{\Omega} m(t, x) \cdot \nabla\psi(x)\kappa(t) dx dt
\end{aligned}$$

which is precisely the identity in statement b). \square

Acknowledgment: We thank Philippe Villedieu for suggesting the problem of deriving the temperature equations for multiphase flows in thermal local non-equilibrium.

REFERENCES

- [1] F. Boyer, P. Fabrie: “Eléments d’analyse pour l’étude de quelques modèles d’écoulements de fluides visqueux incompressibles”. Mathématiques et Applications Vol. 52. Springer Verlag, Berlin, Heidelberg 2006.
- [2] C.E. Brennen: “Fundamentals of Multiphase Flows”. Cambridge University Press, 2005.
- [3] H. Brezis: “Analyse Fonctionnelle. Théorie et Applications.” Masson, Paris 1987.
- [4] D. Cioranescu, F. Murat: *Un terme étrange venu d’ailleurs*. In “Nonlinear Partial Differential Equations and their Applications”, Collège de France Seminar vol. 2. (Paris 1979-1980) Res. Notes in Math. vol. 60, pp. 98–138. Pitman, Boston, London, 1982.
- [5] L. Desvillettes, F. Golse, V. Ricci: *The Mean-Field Limit for Solid Particles in a Navier-Stokes Flow*. J. Stat. Phys. **131** (2008), 941–967.
- [6] F. Fichot, F. Duval, N. Tréguerès, C. Béchaud, M. Quintard: *The impact of thermal non-equilibrium and large-scale 2D/3D effects on debris bed reflooding and coolability*. Nuclear Engineering and Design **236** (2006), 2144–2163.
- [7] V.A. L’vov, E.Ya. Hruslov: *Perturbations of a viscous incompressible fluid by small particles*. Theor. Appl. Quest. Differ. Equ. Algebra **267** (1978), 173–177.
- [8] J.-L. Lions, E. Magenes: “Problèmes aux limites non homogènes et applications. Volume 1”. Dunod, Paris 1968.
- [9] F. Petit, F. Fichot, M. Quintard: *Écoulement diphasique en milieu poreux: modèle à non-équilibre local*. Int. J. Therm. Sci. **38** (1999), 239–249.

(L.D.) ECOLE NORMALE SUPÉRIEURE DE CACHAN, CMLA, 61, AV. DU PdT. WILSON, 94235 CACHAN CEDEX, FRANCE

E-mail address: `desville@cmla.ens-cachan.fr`

(F.G.) ECOLE POLYTECHNIQUE, CENTRE DE MATHÉMATIQUES L. SCHWARTZ, 91128 PALAISEAU CEDEX, FRANCE

E-mail address: `francois.golse@math.polytechnique.fr`

(V.R.) DIPARTIMENTO DI METODI E MODELLI MATEMATICI, UNIVERSITÀ DI PALERMO, VIALE DELLE SCIENZE EDIFICIO 8, 90128 PALERMO, ITALY

E-mail address: `ricci@unipa.it`