Experimental study of forced stratified turbulence

P. Augier, P. Billant, Maria-Eletta Negretti, J.-M. Chomaz

► To cite this version:

P. Augier, P. Billant, Maria-Eletta Negretti, J.-M. Chomaz. Experimental study of forced stratified turbulence. 12th EUROMECH European Turbulence Conference, Sep 2009, Marburg, Germany. pp.397-400, 10.1007/978-3-642-03085-7_95 . hal-01025993

HAL Id: hal-01025993

<https://hal-polytechnique.archives-ouvertes.fr/hal-01025993>

Submitted on 23 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution| 4.0 International License

Experimental study of forced stratified turbulence

P. Augier, P. Billant, E. Negretti and J.-M. Chomaz

LadHyX, Ecole polytechnique, 91128 Palaiseau, France
 pierre.augier@ladhyx.polytechnique.fr

Recent results have shown that strongly stratified turbulence has a three dimensional dynamic instead of a quasi-two dimensional dynamic conjectured previously. The structures are strongly anisotropic with an aspect ratio scaling like the Froude number $l_v/l_h \sim F_h$ [1]. A direct cascade of energy from large scales to small scales associated with a $k_h^{-5/3}$ horizontal kinetic energy spectrum has been predicted and observed in DNS of forced stratified turbulence when the buoyancy Reynolds number $\mathcal{R}^t$ is sufficiently large [2]. In constrast, for small $\mathcal{R}^t$, the flow is dominated by vertical viscous effects even if the Reynolds number is large [3, 4].

Fig. 1. Sketch of the experimental apparatus.

In order to investigate the turbulent regime experimentally with a sufficiently high buoyancy Reynolds number, we have set-up a new experiment where the flow is generated by 12 vortex generators (flaps) placed on the side of a large stratified tank (Fig. 1). Each generator produces periodically and independently a counter-rotating vertical columnar vortex pair which propagates toward the central part of the tank. The interactions between all the

randomly produced vortex pairs give rise to a forced turbulent flow with a low Froude number F_h and a large Reynolds number Re . The buoyancy Reynolds number based on the characteristics of the vortices generated by the flaps is $\mathcal{R}^s \equiv ReF_h^2 \sim 100$ while the buoyancy Reynolds number based on the injection rate of energy ϵ : $\mathcal{R}^t \equiv \epsilon/(\nu N^2)$, is order unity. As seen on figure 2, the mean kinetic energy of the flow is statistically stationary but with high fluctuations typical of turbulent flows.

Fig. 2. Mean kinetic energy as a function of time.

Fig. 3. Vertical cross-section of the velocity field. The background color represents the horizontal vorticity $|\omega_y|$ perpendicular to the cross-section. In heavy shaded regions, $|\omega_y| \sim N$.

PIV measurements in vertical cross-sections show that the flow is organised into horizontal layers (Fig. 3). This layering arises spontaneously via the zigzag instability of individual vortex pairs [5]. When the buoyancy Reynolds number is increased, shear instabilities and intermittent overturning become more and more visible between layers in agreement with [3, 6].

Horizontal velocity spectra exhibit a narrow $k_h^{-5/3}$ inertial domain at intermediate scales ($0.2 \text{ rad/mm} \lesssim k_h \lesssim 0.7 \text{ rad/mm}$) even for the lowest $\mathcal{R}^t$ investigated (Fig. 4). This inertial range becomes more and more flat as $\mathcal{R}^t$ increases as observed in DNS of forced stratified turbulence [3].

Fig. 4. Compensated 1D kinetic energy spectrum : $E(k_h)k_h^{5/3}$ as function of the horizontal wave number k_h for different buoyancy Reynolds numbers.

As shown in figure 5, the rotational and divergent parts of the horizontal velocity are of the same order at all scales even if the forcing is purely vortical. This is consistent with the scaling law $l_v/l_h \sim F_h$ for which the dynamic of waves and vortices can not be separated [1, 3]

We have set-up a novel experiment on forced strongly stratified turbulence with buoyancy Reynolds number of order one. For the first time, a $k_h^{-5/3}$ inertial range has been observed experimentally in agreement with theoretical predictions, DNS and atmospheric measurements.

Fig. 5. Compensated 2D kinetic spectrum of the vortical and divergent parts of the horizontal velocity.

References

1. P. Billant and J.-M. Chomaz, Self-similarity of strongly stratified inviscid flows, *Phys. Fluids*, 13 (2001)
2. E. Lindborg. The energy cascade in a strongly stratified fluid, *J. Fluid Mech.*, 550 (2006)
3. G. Brethouwer, and P. Billant, and E. Lindborg, and J.-M. Chomaz. Scaling analysis and simulation of strongly stratified turbulent flows, *J. Fluid Mech.*, 585 (2007)
4. O. Praud and A.M. Fincham and J. Sommeria, Decaying grid turbulence in a strongly stratified fluid, *J. Fluid Mech.*, 522 (2005)
5. P. Billant and J.-M. Chomaz. Experimental evidence for a new instability of a vertical columnar vortex pair in a strongly stratified fluid, *J. Fluid Mech.*, 418 (2000)
6. A. Deloncle, P. Billant and J.-M. Chomaz. Nonlinear evolution of the zigzag instability in stratified fluids : a shortcut on the route to dissipation, *J. Fluid Mech.*, 599 (2008)