

LINEAR BOLTZMANN EQUATION AND FRACTIONAL DIFFUSION

Claude Bardos, François Golse, Iván Moyano

► **To cite this version:**

Claude Bardos, François Golse, Iván Moyano. LINEAR BOLTZMANN EQUATION AND FRACTIONAL DIFFUSION. Kinetic and Related Models , AIMS, 2018, 11 (4), pp.1011-1036. <10.3934/krm.2018039>. <hal-01579874>

HAL Id: hal-01579874

<https://hal-polytechnique.archives-ouvertes.fr/hal-01579874>

Submitted on 31 Aug 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LINEAR BOLTZMANN EQUATION AND FRACTIONAL DIFFUSION

CLAUDE BARDOS, FRANÇOIS GOLSE, AND IVAN MOYANO

ABSTRACT. Consider the linear Boltzmann equation of radiative transfer in a half-space, with constant scattering coefficient σ . Assume that, on the boundary of the half-space, the radiation intensity satisfies the Lambert (i.e. diffuse) reflection law with albedo coefficient α . Moreover, assume that there is a temperature gradient on the boundary of the half-space, which radiates energy in the half-space according to the Stefan-Boltzmann law. In the asymptotic regime where $\sigma \rightarrow +\infty$ and $1 - \alpha \sim C/\sigma$, we prove that the radiation pressure exerted on the boundary of the half-space is governed by a fractional diffusion equation. This result provides an example of fractional diffusion asymptotic limit of a kinetic model which is based on the harmonic extension definition of $\sqrt{-\Delta}$. This fractional diffusion limit therefore differs from most of other such limits for kinetic models reported in the literature, which are based on specific properties of the equilibrium distributions (“heavy tails”) or of the scattering coefficient as in [U. Frisch-H. Frisch: Mon. Not. R. Astr. Not. **181** (1977), 273–280].

1. INTRODUCTION

The diffusion approximation for the linear Boltzmann equation has been known for a long time and in very different contexts, such as nuclear engineering (see chapter IX in [23]), or radiative transfer (see chapter III.2 in [21]). There are several proofs of the validity of the diffusion approximation, involving rather different mathematical methods: stochastic processes [15, 20, 5], Hilbert expansion [17, 7], or chapter XXI, §5 in [10], moment method [2, 3] . . .

However, there are several situations where the same type of scaling limits of the linear Boltzmann equation lead to fractional (or more generally nonlocal) diffusion equation. A first class of linear Boltzmann equations leading to a fractional diffusion equation includes situations where the scattering rate is not uniformly large over the whole energy range of the particle system considered. See [12] for an example in radiative transfer.

A second class of linear Boltzmann equations leading to a fractional diffusion equation includes the case of collision integrals whose equilibrium solutions have infinite second order moments: see for instance [4, 18, 19, 1].

In the present work, we give an example of a completely different type of asymptotic limit of a linear Boltzmann equation leading to a fractional diffusion equation.

Before describing more precisely the physical problem considered in the present paper, we recall the following well-known fact from classical analysis, which the reader might find convenient to keep in mind.

1991 *Mathematics Subject Classification.* 45K05, 45M05, 35R11 (82C70, 85A25).

Key words and phrases. Linear Boltzmann equation, Radiative transfer equation, Diffusion approximation, Fractional diffusion.

1.1. Fractional Diffusion and Harmonic Extensions. Let¹ $f \in L^p(\mathbf{T}^d)$, with $1 \leq p < \infty$, and let $0 < \gamma < 2$. Then

$$(-\Delta_x)^{\gamma/2} f(x) = -\partial_y F(x, 0),$$

where $F \equiv F(x, y) \in C_b([0, \infty); L^p(\mathbf{T}^d))$ is the unique solution to the boundary value problem

$$\begin{cases} -\Delta_x F(x, y) - \gamma^2 c_\gamma^{\gamma/2} y^{2-2/\gamma} \partial_y^2 F(x, y) = 0, & (x, y) \in \mathbf{T}^d \times (0, +\infty), \\ F(x, 0) = f(x), & x \in \mathbf{T}^d, \end{cases}$$

where

$$c_\gamma = 2^{-\gamma} |\Gamma(-\gamma/2)| / \Gamma(\gamma/2)$$

(See for instance [16], Theorem 1.1 (j) for the analogous result, assuming that the function f satisfies $f \in L^p(\mathbf{R}^d)$ instead of $L^p(\mathbf{T}^d)$.)

1.2. The Model. We seek to describe radiative transfer in a homogeneous medium filling the half-space

$$\{z := (x, y) : y > 0 \text{ and } x \in \mathbf{R}^2\}.$$

We assume that true absorption and emission are negligible effects in this medium. We assume that the only process of physical importance in the medium is scattering (of Thomson type), which means that the elementary scattering process at the level of particles is independent of, and does not modify the frequency of photons. The scattering coefficient $\sigma > 0$ and scattering transition probability (also called “phase function” in §3 of [8]) $p \equiv p(\omega, \omega')$ are assumed to be independent of the position variable z .

Under these assumptions, the radiative intensity (integrated in the frequency variable) at the position z and in the direction ω , henceforth denoted $f \equiv f(z, \omega)$, satisfies the radiative transfer equation (see chapter I in [8]):

$$(1) \quad \omega \cdot \nabla_z f(z, \omega) = \sigma \int_{\mathbf{S}^2} p(\omega, \omega') (f(z, \omega') - f(z, \omega)) d\omega, \quad y > 0, \quad |\omega| = 1.$$

On the plane of equation $y = 0$ (the boundary of the half-space), we assume that the radiation field obeys the Lambert reflection law, with albedo α (see formula (81) in §47 on p. 146 in [8]). On the other hand, we assume that the temperature at the point x of the plane of equation $y = 0$ is $T(x)$, so that the surface emits a radiative intensity $a c T(x)^4$ in all directions at the point x , according to the Stefan-Boltzmann law. The constant a is

$$a := \frac{8\pi^5 k_B^4}{15c^2 \hbar^3},$$

where k_B is the Boltzmann constant, c is the speed of light in the vacuum, and \hbar is the reduced Planck constant.

In other words, the radiative intensity f satisfies the boundary condition

$$(2) \quad f(x, 0, \omega) = a T(x)^4 + \frac{\alpha}{\pi} \int_{\mathbf{S}^2} f(x, 0, \omega') (\omega'_y)^- d\omega', \quad \omega_y > 0.$$

¹We denote by \mathbf{T}^d the d -dimensional torus $\mathbf{T}^d = \mathbf{R}^d / \mathbf{Z}^d$.

1.3. Scaling and Other Assumptions. We are concerned with the following limit of the radiative transfer equation (1) with the boundary condition (2):

- (a) $\sigma \gg 1$ (high scattering regime);
- (b) $0 < 1 - \alpha \ll 1$ (high albedo surface).

We shall adopt the following mathematical setting. For any integer $d \geq 1$, set $Z := \mathbf{T}^d \times (0, +\infty)$; points in Z are denoted by $z = (x, y)$ with $x \in \mathbf{T}^d$ and $y > 0$. Accordingly, unit tangent vectors to Z are denoted by² $\omega = (\omega_x, \omega_y) \in \mathbf{S}^d$, with $\omega_x \in \mathbf{R}^d$ and $\omega_y \in \mathbf{R}$. One has

$$\int_{\mathbf{S}^d} \omega_y^\pm d\omega = \int_0^{\pi/2} \cos \theta (\sin \theta)^{d-1} d\theta = \frac{1}{d} |\mathbf{S}^{d-1}| = |\mathbf{B}^d|.$$

We shall henceforth use the following notation:

$$\langle \phi \rangle := \frac{1}{|\mathbf{S}^d|} \int_{\mathbf{S}^d} \phi(\omega) d\omega, \quad \phi \in C(\mathbf{S}^d),$$

and

$$\langle\langle \psi \rangle\rangle_\pm := \frac{1}{|\mathbf{B}^d|} \int_{\mathbf{S}^d} \psi(\omega) \omega_y^\pm d\omega, \quad \psi \in C(\mathbf{S}^d).$$

We assume that condition (b) is realized by taking the albedo coefficient α of ∂Z of the form

$$\alpha := \frac{\kappa\sigma}{1 + \kappa\sigma}$$

where $\kappa > 0$ is a constant, and we set

$$S(x) := (1 + \kappa\sigma) aT(x)^4.$$

Since $\beta_2 = \pi$, the boundary condition (2) is recast as

$$(3) \quad f(x, 0, \omega) = \frac{S(x)}{1 + \kappa\sigma} + \frac{\kappa\sigma}{1 + \kappa\sigma} \langle\langle f(x, 0, \cdot) \rangle\rangle_-, \quad \omega_y > 0.$$

Observing that the right hand side of (3) is independent of ω , this boundary condition implies that $f(x, 0, \omega)$ is independent of ω for $\omega_y > 0$, so that

$$f(x, 0, \omega) = \frac{f(x, 0, \omega)}{1 + \kappa\sigma} + \frac{\kappa\sigma}{1 + \kappa\sigma} \langle\langle f(x, 0, \cdot) \rangle\rangle_+.$$

Substituting this expression on the left hand side of (3), we arrive at the following equivalent formulation of the boundary condition (3):

$$(4) \quad f(x, 0, \omega) = S(x) - \frac{\kappa\sigma}{|\mathbf{B}^d|} \int_{\mathbf{S}^d} f(x, 0, \omega') \omega'_y d\omega', \quad \omega_y > 0.$$

We further assume that $p \equiv p(\omega, \omega')$ is a measurable function defined a.e. on $\mathbf{S}^d \times \mathbf{S}^d$ satisfying the following condition:

$$(5) \quad p(\omega, \omega') = p(\omega', \omega) > 0 \text{ a.e. on } \mathbf{S}^d \times \mathbf{S}^d, \quad \int_{\mathbf{S}^d} p(\omega, \omega') d\omega' = 1 \text{ for a.e. } \omega \in \mathbf{S}^d.$$

We shall denote by \mathcal{L} the bounded linear operator defined on $L^\infty(\mathbf{S}^d)$ by the formula

$$\mathcal{L}\phi(\omega) = \int_{\mathbf{S}^d} p(\omega, \omega') (\phi(\omega) - \phi(\omega')) d\omega' = \phi(\omega) - \int_{\mathbf{S}^d} p(\omega, \omega') \phi(\omega') d\omega'.$$

²We denote by \mathbf{S}^d the unit sphere in \mathbf{R}^{d+1} , and by \mathbf{B}^d the unit ball in \mathbf{R}^d . We also denote by $|\mathbf{S}^d|$ the d -dimensional area of \mathbf{S}^d , and by $|\mathbf{B}^d|$ the d -dimensional Lebesgue measure of \mathbf{B}^d . We recall that $d|\mathbf{B}^d| = |\mathbf{S}^{d-1}|$ for all $d \geq 1$.

With these elements of notation, the boundary value problem for the radiative transfer equation (1) with boundary condition (2) takes the form

$$(6) \quad \begin{cases} \omega \cdot \nabla_z f_\sigma(z, \omega) + \sigma(\mathcal{L}f_\sigma)(z, \cdot) = 0, & (z, \omega) \in Z \times \mathbf{S}^d, \\ f_\sigma(x, 0, \omega) = \frac{S(x)}{1 + \kappa\sigma} + \frac{\kappa\sigma}{1 + \kappa\sigma} \langle\langle f_\sigma(x, 0, \cdot) \rangle\rangle_-, & x \in \mathbf{T}^d, \omega_y > 0, \end{cases}$$

where

$$(\mathcal{L}f_\sigma)(z, \omega) := (\mathcal{L}f_\sigma(z, \cdot))(\omega).$$

The purpose of the present work is to study the boundary value problem (6) in the asymptotic regime $\sigma \rightarrow +\infty$.

2. MAIN RESULT

2.1. Heuristic approach. In order to gain some intuition on this problem, we apply the Hilbert expansion method. This method is named after Hilbert on the basis of [14], where it has been used for the first time on the Boltzmann equation in the context of the kinetic theory of gases. Its application to the linear Boltzmann equation is discussed in detail in [17, 7].

Thus, we seek f_σ as the formal power series

$$(7) \quad f_\sigma(z, \omega) = \sum_{j \geq 0} \sigma^{-j} f_j(z, \omega), \quad \nabla_z^m f_j \in C(Z \times \mathbf{S}^d) \text{ for all integers } j, m \geq 0.$$

Substituting this ansatz in the radiative transfer equation leads to the sequence of integral equations:

Order $O(\sigma)$:

$$\mathcal{L}f_0(z, \omega) = 0,$$

Order $O(1)$:

$$\mathcal{L}f_1(z, \omega) = -\omega \cdot \nabla_z f_0,$$

.....

Order $O(\sigma^{-j})$:

$$\mathcal{L}f_{j+1}(z, \omega) = -\omega \cdot \nabla_z f_j.$$

.....

Lemma 2.1. *Under assumption (5), the linear operator \mathcal{L} is bounded on $L^2(\mathbf{S}^d; \mathbf{C}^d)$ and satisfies*

$$\|\mathcal{L}\| \leq 2, \quad \text{and} \quad \text{Ker } \mathcal{L} = \{ \text{constants} \} = \mathbf{C}.$$

Proof. Because of the last equality in (5), one has $\mathbf{R} \subset \text{Ker } \mathcal{L}$.

For each $\phi \in L^2(\mathbf{S}^d)$, one has

$$\begin{aligned}
 (8) \quad \int_{\mathbf{S}^d} \phi(\omega)(\mathcal{L}\psi)(\omega) \, d\omega &= \iint_{\mathbf{S}^d \times \mathbf{S}^d} \phi(\omega)p(\omega, \omega')(\psi(\omega) - \psi(\omega')) \, d\omega \, d\omega' \\
 &= \iint_{\mathbf{S}^d \times \mathbf{S}^d} \phi(\omega')p(\omega, \omega')(\psi(\omega') - \psi(\omega)) \, d\omega \, d\omega' \\
 &= \iint_{\mathbf{S}^d \times \mathbf{S}^d} \frac{1}{2}p(\omega, \omega')(\phi(\omega) - \phi(\omega'))(\psi(\omega) - \psi(\omega')) \, d\omega \, d\omega'.
 \end{aligned}$$

The second equality above follows from the symmetry of p in (5), while the third equality is obtained by adding the right hand sides of the first and the second equalities.

If $\phi \in \text{Ker } \mathcal{L}$, then

$$\iint_{\mathbf{S}^d \times \mathbf{S}^d} p(\omega, \omega')(\phi(\omega) - \phi(\omega'))^2 \, d\omega \, d\omega' = 0$$

because of (8) with $\phi = \psi$. Since $p > 0$ a.e. on $\mathbf{S}^d \times \mathbf{S}^d$,

$$\phi(\omega) - \phi(\omega') = 0 \text{ for a.e. } (\omega, \omega') \in \mathbf{S}^d \times \mathbf{S}^d.$$

Averaging both sides of this equality in ω' implies that

$$\phi(\omega) = \langle \phi \rangle \text{ for a.e. } \omega \in \mathbf{S}^d.$$

Thus ϕ is a.e. a constant, and this proves that $\text{Ker } \mathcal{L} = \{ \text{constants} \}$.

Finally

$$\begin{aligned}
 \int_{\mathbf{S}^d} |(\mathcal{L}\phi)(\omega)|^2 \, d\omega &= \int_{\mathbf{S}^d} \left| \int_{\mathbf{S}^d} p(\omega, \omega')(\phi(\omega) - \phi(\omega')) \, d\omega' \right|^2 \, d\omega \\
 &\leq \int_{\mathbf{S}^d} \left(\int_{\mathbf{S}^d} p(\omega, \omega') \, d\omega' \right) \left(\int_{\mathbf{S}^d} p(\omega, \omega') |\phi(\omega) - \phi(\omega')|^2 \, d\omega' \right) \, d\omega \\
 &= \iint_{\mathbf{S}^d} p(\omega, \omega') |\phi(\omega) - \phi(\omega')|^2 \, d\omega \, d\omega' = 2 \int_{\mathbf{S}^d} \overline{\phi(\omega)} (\mathcal{L}\phi)(\omega) \, d\omega,
 \end{aligned}$$

where the inequality follows from the Cauchy-Schwarz inequality applied to the inner integral, and the last equality from (8) with $\overline{\phi} = \psi$. Applying the Cauchy-Schwarz inequality to the last right hand side, one obtains

$$\int_{\mathbf{S}^d} |(\mathcal{L}\phi)(\omega)|^2 \, d\omega \leq 2 \|\phi\|_{L^2(\mathbf{S}^d)} \|\mathcal{L}\phi\|_{L^2(\mathbf{S}^d)},$$

which implies that $\|\mathcal{L}\| \leq 2$. □

We conclude from this lemma the solution to the equation at order $O(\sigma)$:

Order $O(\sigma)$:

$$f_0(z, \omega) = \rho_0(z), \quad \text{for a.e. } (z, \omega) \in Z \times \mathbf{S}^d.$$

Next we study the equation at order $O(1)$ for f_1 . We shall need the following additional assumption on p :

$$(9) \quad p \in L^2(\mathbf{S}^d \times \mathbf{S}^d).$$

Denoting by \mathcal{K} the linear operator defined by

$$(10) \quad (\mathcal{K}\phi)(\omega) := \int_{\mathbf{S}^d} p(\omega, \omega') \phi(\omega') \, d\omega', \quad \omega \in \mathbf{S}^d,$$

one has

$$\mathcal{L} = \mathbf{I} - \mathcal{K}.$$

Lemma 2.2. *The operator \mathcal{L} is bounded and of Fredholm type on $L^2(\mathbf{S}^d; \mathbf{C}^d)$, and satisfies*

$$\mathcal{L} = \mathcal{L}^*, \quad \text{Im } \mathcal{L} = \mathbf{C}^\perp = \left\{ \psi \in L^2(\mathbf{S}^d; \mathbf{C}^d) \text{ s.t. } \int_{\mathbf{S}^d} \psi(\omega) \, d\omega = 0 \right\}.$$

Proof. The operator \mathcal{K} is a self-adjoint Hilbert-Schmidt operator on $L^2(\mathbf{S}^d; \mathbf{C}^d)$ since its integral kernel p is a symmetric real-valued square-integrable function by (5)-(9). In particular \mathcal{K} is a compact operator on $L^2(\mathbf{S}^d; \mathbf{C}^d)$ (see section 6.4.2 and Theorem 6.12 in [6]). This implies that \mathcal{L} is a self-adjoint, bounded operator on $L^2(\mathbf{S}^d; \mathbf{C}^d)$ of Fredholm type (see section 6.4.1 in [6]).

Fredholm's alternative (Theorem 6.6 in [6]) states that

$$\text{Im } \mathcal{L} = (\text{Ker } \mathcal{L})^\perp$$

and we conclude the proof with the characterization of $\text{Ker } \mathcal{L}$ in Lemma 2.1. \square

Since $\langle \omega \rangle = 0$, Fredholm's alternative implies that there exists a unique function $\Omega \equiv \Omega(\omega) \in L^2(\mathbf{S}^d)$ such that

$$(11) \quad \langle \Omega \rangle = 0 \quad \text{and} \quad (\mathcal{L}\Omega)(\omega) = \omega, \quad \text{for a.e. } \omega \in \mathbf{S}^d.$$

Combining Lemmas 2.1 and 2.2 leads to the following result for the solution of the equation at order $O(1)$:

Order $O(1)$:

$$f_1(z, \omega) = \rho_1(z) - \Omega(\omega) \cdot \nabla \rho_0(z), \quad \text{for a.e. } (z, \omega) \in Z \times \mathbf{S}^d.$$

Higher order terms in the expansion are found in the same way, by solving successively for f_{j+1} the integral equations

$$\mathcal{L}f_{j+1} = -\omega \cdot \nabla_z f_j$$

for $j \geq 1$. We shall not pursue this line of investigation, since only f_0 and f_1 will be used in the present section.

At this point, we introduce a last assumption on the scattering transition probability p , specifically, we require that p is rotationally invariant: for each Q in the orthogonal group $O_{d+1}(\mathbf{R})$,

$$(12) \quad p(Q\omega, Q\omega') = p(\omega, \omega'), \quad \text{for a.e. } (\omega, \omega') \in \mathbf{S}^d \times \mathbf{S}^d.$$

Under this assumption, it has been proved in [7] that the diffusion matrix³

$$\langle \omega \otimes \Omega \rangle \text{ is of the form } \langle \omega \otimes \Omega \rangle = \frac{\langle \omega \cdot \Omega \rangle}{d+1} \mathbf{I}.$$

Besides

$$\langle \omega \cdot \Omega \rangle = \langle (\mathcal{L}\Omega) \cdot \Omega \rangle = \frac{1}{2} \iint_{\mathbf{S}^d \times \mathbf{S}^d} p(\omega, \omega') |\Omega(\omega) - \Omega(\omega')|^2 \, d\omega \, d\omega' > 0.$$

Notice that this inequality is strict: otherwise, one would have

$$p(\omega, \omega') (\Omega(\omega) - \Omega(\omega')) = 0 \quad \text{for a.e. } (\omega, \omega') \in \mathbf{S}^d \times \mathbf{S}^d.$$

³See formulas (40)-(44) on p. 624 in [7]. Actually, one can deduce from equation (42) in [7] that Ω is of the form $\Omega(\omega) = \lambda\omega$: see Lemma 3 in [11], or Lemma 8 in Appendix 1 of [13]. With Ω of this form, one immediately concludes that $\langle \omega \otimes \Omega \rangle = \frac{\lambda}{d+1} \mathbf{I}$.

This would imply that

$$0 = \int_{\mathbf{S}^d} p(\omega, \omega') (\Omega(\omega) - \Omega(\omega')) d\omega' = (\mathcal{L}\Omega)(\omega) = \omega$$

which is obviously a contradiction.

From the equation at order $O(\sigma)$, i.e.

$$\mathcal{L}f_2 = -\omega \cdot \nabla_z f_1,$$

we conclude that

$$\operatorname{div}_z \langle \omega f_1 \rangle = 0.$$

Otherwise, one would have $\omega \cdot \nabla_z f_1 \notin \operatorname{Im} \mathcal{L}$ according to Lemma 2.2. With the expression for f_1 obtained above, this equality takes the form

$$\operatorname{div}_z \left(-\frac{\langle \omega \cdot \Omega \rangle}{d+1} \nabla_z \rho_0 \right) = 0,$$

or equivalently

$$(13) \quad -\Delta \rho_0(z) = 0, \quad z \in Z.$$

Moreover, if the Hilbert expansion (7) holds at the boundary, then we deduce from the equivalent formulation (4) of the boundary condition in (6) that

$$\begin{aligned} \rho_0(x, 0) &= S(x) - \frac{\kappa\sigma}{|\mathbf{B}^d|} \int_{\mathbf{S}^d} \left(\rho_1(x, 0) - \frac{1}{\sigma} \Omega(\omega') \cdot \nabla \rho_0(x, 0) \right) \omega'_y d\omega' \\ (14) \quad &= S(x) + \frac{\kappa|\mathbf{S}^d|}{|\mathbf{B}^d|} \langle \omega_y \Omega \rangle \cdot \nabla \rho_0(x, 0) \\ &= S(x) + \frac{\kappa|\mathbf{S}^d| \langle \omega \cdot \Omega \rangle}{|\mathbf{B}^d|(d+1)} \partial_y \rho_0(x, 0), \quad x \in \mathbf{T}^d. \end{aligned}$$

Returning to the original variables, and to the case of physical interest where $d = 2$, we recall that the radiation pressure $\mathcal{P}(x)$ at the point x of ∂Z is defined by the identity

$$\frac{1}{c} \int_{\mathbf{S}^2} \omega \otimes \omega f(x, 0, \omega) d\omega = \mathcal{P}(x) \mathbf{I}.$$

(See for instance formula (1.19) in [21]). Therefore, to leading order in $1/\sigma$, one has

$$\mathcal{P}(x) \simeq \frac{4\pi}{3c} \rho_0(x, 0).$$

Since $\frac{4\pi}{3c} \rho_0$ is an harmonic extension of \mathcal{P} , applying the formula for the fractional diffusion operator recalled in section 1.1 leads to the following equation for the radiation pressure field on the boundary:

$$(15) \quad \frac{3}{4}(1 - \alpha)\mathcal{P}(x) + \alpha \langle \omega \cdot \Omega \rangle \frac{1}{\sigma} (-\Delta_x)^{1/2} \mathcal{P}(x) = \frac{a\pi T(x)^4}{c}, \quad x \in \mathbf{T}^2.$$

Assumption (12) is verified by most of the scattering transition probabilities p used in practice, such as

$$\begin{aligned} p(\omega, \omega') &= \frac{1}{|\mathbf{S}^d|}, & (\text{isotropic scattering}) \\ p(\omega, \omega') &= \frac{3}{16\pi} (1 + (\omega \cdot \omega')^2), & (\text{Rayleigh phase function for } d = 2) \end{aligned}$$

(see formulas (30)-(31)-(33) in §3 or formula (192) in §16 of [8]). More generally, all scattering transition probabilities of the form

$$(16) \quad p(\omega, \omega') = P((\omega \cdot \omega')^2),$$

under the normalizing condition

$$\int_0^1 P(\mu^2)(1 - \mu^2)^{\frac{d}{2}-1} d\mu = \frac{1}{2|\mathbf{S}^{d-1}|},$$

satisfy (12). In that case

$$\int_{\mathbf{S}^d} P((\omega \cdot \omega')^2) \omega' d\omega' = 0$$

(since the integral the right hand side of this last equality must be invariant under the substitution $\omega' \mapsto -\omega'$), so that

$$\mathcal{L}\omega = \omega, \quad \text{and therefore} \quad \Omega(\omega) = \omega.$$

In that case

$$\langle \omega \cdot \Omega \rangle = 1,$$

and the diffusion coefficient in the usual diffusion approximation of the linear Boltzmann equation is

$$\frac{\langle \omega \cdot \Omega \rangle}{(d+1)\sigma} = \frac{1}{(d+1)\sigma}.$$

In particular, the equation (15) satisfied by the radiation pressure field on the boundary ∂Z is, in this case,

$$\frac{3}{4}(1 - \alpha)\mathcal{P}(x) + \frac{\alpha}{\sigma}(-\Delta_x)^{1/2}\mathcal{P}(x) = \frac{a\pi T(x)^4}{c}, \quad x \in \mathbf{T}^2.$$

2.2. The Limit Theorem. The analysis based on Hilbert's expansion presented in the previous section is only formal. A rigorous analysis of the problem based on the moment method for kinetic models leads to the following result.

Theorem 2.3. *Assume that $p \equiv p(\omega, \omega')$ is a measurable function defined a.e. on $\mathbf{S}^d \times \mathbf{S}^d$, satisfying (5)-(9)-(12). Denote by Ω the unique element of $L^2(\mathbf{S}^d)$ defined by (11). Let $S \in W^{1,\infty}(\mathbf{T}^d)$.*

(a) *For each $\sigma > 0$, the boundary value problem (6) has a unique solution*

$$f_\sigma \in L^\infty(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d);$$

this solution satisfies

$$\|f_\sigma\|_{L^\infty(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d)} \leq \|S\|_{L^\infty(\mathbf{T}^d)},$$

and

$$\|\partial_{x_j} f_\sigma\|_{L^\infty(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d)} \leq \|\partial_{x_j} S\|_{L^\infty(\mathbf{T}^d)}$$

for all $j = 1, \dots, d$.

(b) *In the limit as $\sigma \rightarrow +\infty$, one has*

$$f_\sigma \rightarrow \rho \equiv \rho(z) \text{ in } L^\infty(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d) \text{ weak-}^*$$

and

$$\sigma(f_\sigma - \langle f_\sigma \rangle) \rightarrow -\Omega \cdot \nabla \rho \text{ in } L^2(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d) \text{ weak,}$$

where $\rho \equiv \rho(z)$ satisfies the properties

$$\rho \in L^\infty(\mathbf{T}^d \times (0, +\infty)), \quad \nabla \rho \in L^2(\mathbf{T}^d \times (0, +\infty)),$$

and is the weak solution of the boundary value problem

$$(17) \quad \begin{cases} -\Delta_z \rho(z) = 0, & z = (x, y) \in \mathbf{T}^d \times (0, +\infty), \\ \left(\rho - \kappa \langle \omega \cdot \Omega \rangle \frac{|\mathbf{B}^{d+1}|}{|\mathbf{B}^d|} \partial_y \rho \right) \Big|_{y=0} = S, & x \in \mathbf{T}^d. \end{cases}$$

(c) In the limit as $\sigma \rightarrow +\infty$, one has

$$f_\sigma \Big|_{y=0} \rightarrow R \quad \text{in } L^2(\mathbf{T}^d \times \mathbf{S}^d; |\omega_y| \, dx \, d\omega) \text{ weak,}$$

and

$$\sigma \langle \omega_y f_\sigma \rangle \Big|_{y=0} \rightarrow -\frac{\langle \omega \cdot \Omega \rangle}{d+1} \partial_y \rho \Big|_{y=0} \text{ in } H^{-1/2}(\mathbf{T}^d) \text{ weak,}$$

where $R \equiv R(x)$ is the solution to the fractional diffusion equation

$$(18) \quad R(x) + \kappa \langle \omega \cdot \Omega \rangle \frac{|\mathbf{B}^{d+1}|}{|\mathbf{B}^d|} (-\Delta_x)^{1/2} R(x) = S(x), \quad x \in \mathbf{T}^d.$$

3. PROOF OF THEOREM 2.3

3.1. **Step 1.** Consider the boundary value problem

$$\begin{cases} \lambda h + \omega \cdot \nabla_z h + \sigma h = \sigma Q(z, \omega), \\ h(x, 0, \omega) = H(x), \quad \omega_y > 0, \end{cases}$$

where $Q \in L^\infty(Z \times \mathbf{S}^d)$ and $H \in L^\infty(\mathbf{T}^d)$.

Its unique solution is given by the method of characteristics

$$\begin{aligned} h(z, \omega) &= H\left(x - \frac{y}{\omega_y} \omega_x\right) e^{-(\lambda+\sigma)y/\omega_y} \\ &\quad + \int_0^{y/\omega_y} \sigma e^{-(\lambda+\sigma)t} Q(z - t\omega, \omega) \, dt, \quad \omega_y > 0, \\ h(z, \omega) &= \int_0^\infty \sigma e^{-(\lambda+\sigma)t} Q(z - t\omega, \omega) \, dt, \quad \omega_y < 0. \end{aligned}$$

Setting

$$\theta := e^{-(\lambda+\sigma)y/\omega_y}$$

we find that, for a.e. $(z, \omega) \in Z \times \mathbf{S}^d$,

$$\begin{aligned} |h(z, \omega)| &\leq \max(\theta \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} (1-\theta) \|Q\|_{L^\infty}, \frac{\sigma}{\sigma+\lambda} \|Q\|_{L^\infty}) \\ &= \max\left(\frac{\lambda\theta}{\sigma+\lambda} \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} ((1-\theta) \|Q\|_{L^\infty} + \theta \|H\|_{L^\infty}), \frac{\sigma}{\sigma+\lambda} \|Q\|_{L^\infty}\right) \\ &\leq \max\left(\frac{\lambda}{\sigma+\lambda} \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} \max(\|Q\|_{L^\infty}, \|H\|_{L^\infty}), \frac{\sigma}{\sigma+\lambda} \|Q\|_{L^\infty}\right). \end{aligned}$$

This inequality implies, on the one hand, that, for each $H \in L^\infty(\mathbf{T}^d)$, the map

$$Q \mapsto \mathcal{K}h$$

is a contraction in $L^\infty(Z \times \mathbf{S}^d)$ with Lipschitz constant $\leq \frac{\sigma}{\lambda+\sigma}$. Indeed, denoting by h_1 and h_2 corresponding to Q_1 and Q_2 respectively, one has

$$|(h_1 - h_2)(z, \omega)| \leq \frac{\sigma}{\sigma+\lambda} \|Q_1 - Q_2\|_{L^\infty},$$

as a consequence of the previous inequality. Indeed, by linearity $h_1 - h_2$ is a solution of the boundary value problem above with source term $Q_1 - Q_2$ and boundary data $H = 0$.

Using assumption (5), we see that the operator \mathcal{K} defined in (10) is a bounded operator on $L^\infty(\mathbf{S}^d)$, satisfying

$$\|\mathcal{K}\|_{\mathcal{L}(L^\infty(\mathbf{S}^d))} \leq 1.$$

Hence

$$\|\mathcal{K}h_1 - \mathcal{K}h_2\|_{L^\infty} \leq \|h_1 - h_2\|_{L^\infty} \leq \frac{\sigma}{\sigma+\lambda} \|Q_1 - Q_2\|_{L^\infty}.$$

Therefore, the map $Q \mapsto \mathcal{K}h$ has a unique fixed point in $L^\infty(Z \times \mathbf{S}^d)$.

In other words, there exists a unique solution $g \in L^\infty(Z \times \mathbf{S}^d)$ to the boundary value problem

$$(19) \quad \begin{cases} \lambda g + \omega \cdot \nabla_z g + \sigma \mathcal{L}g = 0, & (z, \omega) \in Z \times \mathbf{S}^d, \\ g(x, 0, \omega) = H(x), & \omega_y > 0, \end{cases}$$

and this solution satisfies the bound

$$\|g\|_{L^\infty} \leq \max\left(\frac{\lambda}{\sigma+\lambda} \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} \max(\|\mathcal{K}g\|_{L^\infty}, \|H\|_{L^\infty}), \frac{\sigma}{\sigma+\lambda} \|\mathcal{K}g\|_{L^\infty}\right).$$

It is obviously impossible that

$$\|\mathcal{K}g\|_{L^\infty} > \|H\|_{L^\infty}.$$

Indeed, since

$$\|\mathcal{K}g\|_{L^\infty} \leq \|g\|_{L^\infty},$$

the inequality above would imply that

$$\|\mathcal{K}g\|_{L^\infty} \leq \frac{\lambda}{\sigma+\lambda} \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} \|\mathcal{K}g\|_{L^\infty},$$

which would imply in turn

$$\|\mathcal{K}g\|_{L^\infty} \leq \|H\|_{L^\infty},$$

in contradiction with our assumption. Hence

$$\|\mathcal{K}g\|_{L^\infty} \leq \|H\|_{L^\infty},$$

and therefore

$$\begin{aligned} \|g\|_{L^\infty} &\leq \max\left(\frac{\lambda}{\sigma+\lambda} \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} \max(\|\mathcal{K}g\|_{L^\infty}, \|H\|_{L^\infty}), \frac{\sigma}{\sigma+\lambda} \|\mathcal{K}g\|_{L^\infty}\right) \\ &= \max\left(\frac{\lambda}{\sigma+\lambda} \|H\|_{L^\infty} + \frac{\sigma}{\sigma+\lambda} \|H\|_{L^\infty}, \frac{\sigma}{\sigma+\lambda} \|\mathcal{K}g\|_{L^\infty}\right) = \|H\|_{L^\infty}. \end{aligned}$$

3.2. Step 2. With step 1, for each $\lambda > 0$ and $\sigma > 0$, we have constructed a linear map

$$\mathcal{T}_{\lambda, \sigma} : L^\infty(\mathbf{T}^d) \mapsto L^\infty(Z \times \mathbf{S}^d)$$

defined by the formula

$$\mathcal{T}_{\lambda, \sigma} H = g,$$

where g is the solution to the boundary value problem (19). We have also proved that

$$\|\mathcal{T}_{\lambda, \sigma} H\|_{L^\infty} \leq \|H\|_{L^\infty}.$$

Now we seek to solve the boundary value problem

$$(20) \quad \begin{cases} \lambda f + \omega \cdot \nabla_z f + \sigma \mathcal{L}f = 0, & (z, \omega) \in Z \times \mathbf{S}^d, \\ f(x, 0, \omega) = \frac{1}{1+\beta} S(x) + \frac{\beta}{1+\beta} \langle\langle f(x, 0, \cdot) \rangle\rangle_-, & \omega_y > 0, \end{cases}$$

Consider the map

$$\mathcal{A}_{\lambda, \sigma, \beta} : L^\infty(\mathbf{T}^d) \rightarrow L^\infty(\mathbf{T}^d)$$

defined by the formula

$$\mathcal{A}_{\lambda, \sigma, \beta} F(x) := \sigma \left\langle\langle \int_0^\infty e^{-(\lambda+\sigma)t} \mathcal{K} \left(\mathcal{T}_{\lambda, \sigma} \left(\frac{S}{1+\beta} + \frac{\beta F}{1+\beta} \right) \right) (x - t\omega, \omega) dt \right\rangle\rangle_-.$$

In terms of the operator \mathcal{K} defined in (10), one has obviously

$$(\mathcal{A}_{\lambda,\sigma,\beta}F_1 - \mathcal{A}_{\lambda,\sigma,\beta}F_2)(x) = \frac{\beta\sigma}{1+\beta} \left\langle \int_0^\infty e^{-(\lambda+\sigma)t} \mathcal{K}(\mathcal{T}_{\lambda,\sigma}(F_1 - F_2))(x - t\omega, \omega) dt \right\rangle_{-},$$

so that, for a.e. $x \in \mathbf{T}^d$,

$$\begin{aligned} |(\mathcal{A}_{\lambda,\sigma,\beta}F_1 - \mathcal{A}_{\lambda,\sigma,\beta}F_2)(x)| &\leq \frac{\beta\sigma}{1+\beta} \|\mathcal{T}_{\lambda,\sigma}(F_1 - F_2)\|_{L^\infty} \int_0^\infty e^{-(\lambda+\sigma)t} dt \\ &\leq \frac{\beta\sigma}{1+\beta} \|F_1 - F_2\|_{L^\infty} \int_0^\infty e^{-(\lambda+\sigma)t} dt \\ &= \frac{\beta}{1+\beta} \frac{\sigma}{\lambda+\sigma} \|F_1 - F_2\|_{L^\infty}. \end{aligned}$$

In other words, $\mathcal{A}_{\lambda,\sigma,\beta}$ is a contraction in the Banach space $L^\infty(\mathbf{T}^d)$ with Lipschitz constant $\leq \frac{\beta}{1+\beta} \frac{\sigma}{\lambda+\sigma} < 1$.

By the fixed point theorem, there exists a unique $F \in L^\infty(\mathbf{T}^d)$ such that

$$\mathcal{A}_{\lambda,\sigma,\beta}F = F.$$

The solution to the boundary value problem (20) is given by the formula

$$f := \mathcal{T}_{\lambda,\sigma} \left(\frac{S}{1+\beta} + \frac{\beta F}{1+\beta} \right).$$

The fixed point F satisfies in particular the bound

$$\|F - \mathcal{A}_{\lambda,\sigma,\beta}0\|_{L^\infty} \leq \frac{\beta}{1+\beta} \frac{\sigma}{\lambda+\sigma} \|F\|_{L^\infty}.$$

Since

$$|\mathcal{A}_{\lambda,\sigma,\beta}0(x)| = \sigma \left\langle \int_0^\infty e^{-(\lambda+\sigma)t} \mathcal{K} \left(\mathcal{T}_{\lambda,\sigma} \frac{S}{1+\beta} \right) (x - t\omega, \omega) dt \right\rangle_{-},$$

one has

$$\|\mathcal{A}_{\lambda,\sigma,\beta}0\|_{L^\infty} \leq \frac{\sigma}{1+\beta} \|S\|_{L^\infty} \int_0^\infty e^{-(\lambda+\sigma)t} dt = \frac{1}{1+\beta} \frac{\sigma}{\lambda+\sigma} \|S\|_{L^\infty},$$

so that

$$\|F\|_{L^\infty} \leq \frac{1}{1+\beta} \frac{\sigma}{\lambda+\sigma} \|S\|_{L^\infty} + \frac{\beta}{1+\beta} \frac{\sigma}{\lambda+\sigma} \|F\|_{L^\infty}.$$

Therefore

$$\|F\|_{L^\infty} \leq \frac{\sigma}{\lambda(1+\beta) + \sigma} \|S\|_{L^\infty},$$

which implies in turn

$$\begin{aligned} \|f\|_{L^\infty} &= \left\| \mathcal{T}_{\lambda,\sigma} \left(\frac{S}{1+\beta} + \frac{\beta F}{1+\beta} \right) \right\|_{L^\infty} \\ &\leq \left(\frac{1}{1+\beta} + \frac{\beta}{1+\beta} \frac{\sigma}{\lambda(1+\beta) + \sigma} \right) \|S\|_{L^\infty} \leq \|S\|_{L^\infty}. \end{aligned}$$

3.3. Step 3. Call f_λ the unique solution of the boundary value problem (20) obtained in step 2, for which we have obtained the bound

$$(21) \quad \|f_\lambda\|_{L^\infty} \leq \|S\|_{L^\infty}, \quad \lambda > 0,$$

which is uniform in $\lambda > 0$.

Now we consider the boundary value problem

$$(22) \quad \begin{cases} \omega \cdot \nabla_z f + \sigma \mathcal{L}f = 0, & (z, \omega) \in Z \times \mathbf{S}^d, \\ f(x, 0, \omega) = \frac{1}{1+\beta} S(x) + \frac{\beta}{1+\beta} \langle\langle f(x, 0, \cdot) \rangle\rangle_-, & \omega_y > 0, \end{cases}$$

of which we seek a solution by passing to the limit in f_λ as $\lambda \rightarrow 0$.

On account of the uniform bound (21), the Banach-Alaoglu theorem implies the existence of a sequence $\lambda_n \rightarrow 0$ such that

$$f_{\lambda_n} \rightarrow f \quad \text{in } L^\infty(Z \times \mathbf{S}^d) \text{ weak} - *.$$

One has obviously

$$\lambda_n f_{\lambda_n} + \omega \cdot \nabla_z f_{\lambda_n} + \sigma \mathcal{L}f_{\lambda_n} \rightarrow \omega \cdot \nabla_z f + \sigma \mathcal{L}f$$

in $\mathcal{D}'(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d)$, so that

$$\omega \cdot \nabla_z f + \sigma \mathcal{L}f = 0 \quad \text{dans } \mathcal{D}'(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d).$$

For each $\phi \in W^{1,1}(\mathbf{T}^d)$, one has

$$\begin{aligned} \partial_y \left(\omega_y \int_{\mathbf{T}^d} f_{\lambda_n}(x, y, \omega) \phi(x) dx \right) &= -\lambda_n \int_{\mathbf{T}^d} f_{\lambda_n}(x, y, \omega) dx \\ &\quad + \int_{\mathbf{T}^d} \omega_x \cdot \nabla \phi(x) f_{\lambda_n}(x, y, \omega) dx \\ &\quad - \sigma \int_{\mathbf{T}^d} \mathcal{L}f_{\lambda_n}(x, y, \omega) \phi(x) dx. \end{aligned}$$

Therefore

$$\int_{\mathbf{T}^d} f_{\lambda_n}(x, y, \omega) \phi(x) dx \quad \text{et} \quad \partial_y \left(\omega_y \int_{\mathbf{T}^d} f_{\lambda_n}(x, y, \omega) \phi(x) dx \right)$$

are bounded in $L^\infty((0, +\infty) \times \mathbf{S}^d)$. By Ascoli-Arzelà's theorem,

$$\omega_y f_{\lambda_n}(x, 0, \omega) \rightarrow \omega_y f(x, 0, \omega) \quad \text{in } L^\infty(\mathbf{S}^d; W^{-1,\infty}(\mathbf{T}^d)) \text{ weak} - *,$$

and

$$\langle\langle f_{\lambda_n} \rangle\rangle_-(\cdot, 0) \rightarrow \langle\langle f \rangle\rangle_-(\cdot, 0) \quad \text{dans } W^{-1,\infty}(\mathbf{T}^d) \text{ weak} - *.$$

By passing to the limit in the boundary condition

$$f_{\lambda_n}(x, 0, \omega) = \frac{1}{1+\beta} S(x) + \frac{\beta}{1+\beta} \langle\langle f_{\lambda_n}(x, 0, \cdot) \rangle\rangle_-, \quad \omega_y > 0, \quad x \in \mathbf{T}^d,$$

we conclude that

$$\omega_y^+ f|_{y=0} = \frac{\omega_y^+ S}{1+\beta} + \frac{\beta \omega_y^+}{1+\beta} \langle\langle f \rangle\rangle_-|_{y=0} \quad \text{in } L^\infty(\mathbf{S}^d; W^{-1,\infty}(\mathbf{T}^d)).$$

In particular

$$f(\cdot, 0, \omega) = \frac{S}{1+\beta} + \frac{\beta}{1+\beta} \langle\langle f \rangle\rangle_-(\cdot, 0) \quad \text{in } W^{-1,\infty}(\mathbf{T}^d)$$

for a.e. $\omega \in \mathbf{S}^d$ such that $\omega_y > 0$.

The uniform bound (21) obviously implies that the solution to the boundary value problem (22) satisfies the bound

$$\|f\|_{L^\infty} \leq \|S\|_{L^\infty}.$$

3.4. Step 4. In this step, we check that, for each $S \in L^\infty(\mathbf{T}^d)$, there exists at most one weak solution f of the boundary value problem

$$\begin{cases} \omega \cdot \nabla_z f + \sigma \mathcal{L}f = 0, & (z, \omega) \in Z \times \mathbf{S}^d, \\ f(x, 0, \omega) = \frac{1}{1+\beta} S(x) + \frac{\beta}{1+\beta} \langle\langle f(x, 0, \cdot) \rangle\rangle_-, & \omega_y > 0, \end{cases}$$

in the space $L^\infty(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d)$.

By linearity, it is enough to show that, if $g \in L^\infty(\mathbf{T}^d \times (0, +\infty) \times \mathbf{S}^d)$ is a solution to the boundary value problem

$$\begin{cases} \omega \cdot \nabla_z g + \sigma \mathcal{L}g = 0, & (z, \omega) \in Z \times \mathbf{S}^d, \\ g(x, 0, \omega) = \frac{\beta}{1+\beta} \langle\langle g(x, 0, \cdot) \rangle\rangle_-, & \omega_y > 0, \end{cases}$$

then $g = 0$, a.e..

Denote the sequence of Fourier coefficients of g in the y variable by

$$\hat{g}(k, y, \omega) = \int_{\mathbf{T}^d} e^{-2\pi i k \cdot x} g(x, y, \omega) dx, \quad k \in \mathbf{Z}^d.$$

Then, for each $k \in \mathbf{Z}^d$, on a

$$\begin{cases} \omega_y \partial_y \hat{g}(k, y, \omega) + i2\pi k \cdot \omega_x \hat{g}(k, y, \omega) + \sigma \mathcal{L}g(k, y, \omega) = 0, & y > 0, |\omega| = 1, \\ \hat{g}(k, 0, \omega) = \frac{\beta}{1+\beta} \langle\langle g \rangle\rangle_-(k, 0), & \omega_y > 0. \end{cases}$$

For each k and for a.e. $\omega \in \mathbf{S}^d$, the function $y \mapsto g(k, y, \omega)$ belongs therefore to $W^{1,\infty}(0, +\infty)$.

Multiplying both sides of the differential equation above by $\overline{\hat{g}(k, y, \omega)}$ leads to the identity

$$\overline{(\hat{g} \omega_y \partial_y \hat{g})}(k, y, \omega) + i2\pi k \cdot \omega_x |\hat{g}(k, y, \omega)|^2 + \sigma \overline{\hat{g}(k, y, \omega)} \mathcal{L} \hat{g}(k, y, \omega) = 0.$$

Taking the real part of the right hand side of the equality above, we arrive at the identity

$$\begin{aligned} \omega_y \partial_y |\hat{g}(k, y, \omega)|^2 + 2\sigma |\hat{g}(k, y, \omega) - \langle \hat{g} \rangle(k, y)|^2 \\ = -2\sigma \Re \left(\overline{\hat{g}(k, y, \omega)} \mathcal{L} \hat{g}(k, y, \omega) \right). \end{aligned}$$

Now we average both sides of this equality in ω : for each $k \in \mathbf{Z}^d$,

$$\partial_y \langle \omega_y |\hat{g}|^2 \rangle(k, y) + 2\sigma \Re \langle \overline{\hat{g}(k, y, \cdot)} \mathcal{L} \hat{g}(k, y, \cdot) \rangle = 0, \quad y > 0.$$

At this point, we recall the following property of \mathcal{L} .

Lemma 3.1. *Under assumptions (5)-(9) on the function p , the operator \mathcal{L} satisfies the following properties. For each $\phi \in L^2(\mathbf{S}^d; \mathbf{C})$, one has*

$$\langle \overline{\phi} \mathcal{L} \phi \rangle = \frac{1}{2|\mathbf{S}^d|} \iint_{\mathbf{S}^d \times \mathbf{S}^d} p(\omega, \omega') |\phi(\omega) - \phi(\omega')|^2 d\omega d\omega'.$$

Moreover, there exists $\mu > 0$ such that

$$\langle \overline{\phi} \mathcal{L} \phi \rangle \geq \mu \|\phi - \langle \phi \rangle\|_{L^2(\mathbf{S}^d)}^2.$$

The proof of this lemma is deferred until the end of the present step.

Thus, for each $k \in \mathbf{Z}^d$, the function

$$y \mapsto \langle \omega_y |\hat{g}|^2 \rangle(k, y)$$

is bounded and nonincreasing on $(0, +\infty)$, and applying Lemma 3.1 shows that

$$2\sigma\mu \int_0^\infty \left\langle |\hat{g} - \langle \hat{g} \rangle|^2 \right\rangle(k, y) \, dy \leq \lim_{Y \rightarrow +\infty} [\langle \omega_y |\hat{g}|^2 \rangle(k, \cdot)]_{y=Y}^{y=0} < +\infty.$$

Next observe that

$$(23) \quad \langle \omega_y |\hat{g}|^2 \rangle(k, y) = \langle \omega_y |\hat{g} - \langle \hat{g} \rangle|^2 \rangle(k, y) + 2\Re \left(\langle \omega_y (\hat{g} - \langle \hat{g} \rangle) \rangle(k, y) \overline{\langle \hat{g} \rangle(k, y)} \right).$$

In view of the previous estimate, the first term on the right hand side of this equality belongs to $L^1(0, +\infty)$ since $|\omega_y| \leq 1$, while the second term is bounded by the Cauchy-Schwarz inequality:

$$2\Re \left(\langle \omega_y (\hat{g} - \langle \hat{g} \rangle) \rangle(k, y) \overline{\langle \hat{g} \rangle(k, y)} \right) \leq \|g\|_{L^\infty} \langle \omega_y^2 |g - \langle g \rangle|^2 \rangle(k, y)^{1/2}.$$

Hence the second term on the right hand side of (23) belongs to $L^2(0, +\infty)$, and since the first term is also bounded in $L^\infty(0, +\infty)$ because g is assumed to belong to $L^\infty(Z \times \mathbf{S}^d)$, we conclude that

$$y \mapsto \langle \omega_y |\hat{g}|^2 \rangle(k, y) \text{ belongs to } L^2(0, +\infty)$$

for each $k \in \mathbf{Z}^d$. Since this function is moreover nonincreasing, we conclude that

$$0 = \lim_{Y \rightarrow +\infty} \langle \omega_y |\hat{g}|^2 \rangle(k, Y) \leq \langle \omega_y |\hat{g}|^2 \rangle(k, y) \quad \text{for all } y > 0 \text{ and all } k \in \mathbf{Z}^d.$$

Thus, for each $Y > 0$, one has

$$\begin{cases} \langle \omega_y |\hat{g}|^2 \rangle(k, Y) + 2\sigma\mu \int_0^Y \left\langle |\hat{g} - \langle \hat{g} \rangle|^2 \right\rangle(k, y) \, dy \leq \langle \omega_y |\hat{g}|^2 \rangle(k, 0), \\ \langle \omega_y |\hat{g}|^2 \rangle(k, Y) \geq 0. \end{cases}$$

Moreover, the right hand side of this inequality satisfies

$$\begin{aligned} \langle \omega_y |\hat{g}|^2 \rangle(k, 0) &= \frac{|\mathbf{B}^d|}{|\mathbf{S}^d|} \left(\langle\langle |\hat{g}|^2 \rangle\rangle_+(k, 0) - \langle\langle |\hat{g}|^2 \rangle\rangle_-(k, 0) \right) \\ &= \frac{|\mathbf{B}^d|}{|\mathbf{S}^d|} \left(\left(\frac{\beta}{1+\beta} \right)^2 |\langle\langle g \rangle\rangle_-(k, 0)|^2 \langle\langle 1 \rangle\rangle_+ - \langle\langle |\hat{g}|^2 \rangle\rangle_-(k, 0) \right) \\ &= \frac{|\mathbf{B}^d|}{|\mathbf{S}^d|} \left(\left(\frac{\beta}{1+\beta} \right)^2 |\langle\langle g \rangle\rangle_-(k, 0)|^2 - \langle\langle |\hat{g}|^2 \rangle\rangle_-(k, 0) \right) \\ &\leq \frac{|\mathbf{B}^d|}{|\mathbf{S}^d|} \left(\left(\frac{\beta}{1+\beta} \right)^2 - 1 \right) \langle\langle |\hat{g}|^2 \rangle\rangle_-(k, 0) \end{aligned}$$

by the Cauchy-Schwarz inequality.

Summarizing, we have proved that, for all $k \in \mathbf{Z}^d$ and all $Y > 0$

$$0 = \langle \omega_y |\hat{g}|^2 \rangle(k, Y) = \int_0^Y \left\langle |\hat{g} - \langle \hat{g} \rangle|^2 \right\rangle(k, y) \, dy = \langle\langle |\hat{g}|^2 \rangle\rangle_-(k, 0).$$

In particular, since $\mathcal{L}1 = 0$, one has

$$\mathcal{L}\hat{g} = \mathcal{L}(\hat{g} - \langle \hat{g} \rangle) = 0,$$

so that

$$\begin{cases} \omega_y \partial_y \hat{g}(k, y, \omega) + i2\pi k \cdot \omega_x \hat{g}(k, y, \omega) = g(k, y, \omega) - \langle \hat{g} \rangle(k, y) = 0, \\ \hat{g}(k, 0, \omega) = \frac{\beta}{1+\beta} \langle\langle g \rangle\rangle_-(k, 0) = 0, \quad \omega_y > 0. \end{cases}$$

The second equality above implies that, for a.e. $(k, y) \in \mathbf{Z}^d \times (0, +\infty)$, the function $\omega \mapsto \hat{g}(k, y, \omega)$ is a.e. constant on \mathbf{S}^d . Hence

$$\omega_y \partial_y \hat{g}(k, y) + i2\pi k \cdot \omega_x \hat{g}(k, y) = 0, \text{ for all } \omega \in \mathbf{S}^d, \text{ and a.e. } (k, y) \in \mathbf{Z} \times (0, +\infty).$$

Setting $\omega_x = k/|k|$ and $\omega_y = 0$ shows that

$$k \neq 0 \Rightarrow \hat{g}(k, y) = 0 \text{ for a.e. } y \geq 0.$$

On the other hand

$$\partial_y \hat{g}(0, y) = 0$$

by choosing $\omega_x = 0$ and $\omega_y = 1$, so that

$$\hat{g}(0, y) = \hat{g}(0, 0).$$

Finally, the boundary condition implies that

$$\hat{g}(0, y) = \hat{g}(0, 0) = 0, \quad y > 0.$$

Therefore, we have proved that $\hat{g}(k, y, \omega) = 0$ for a.e. $(y, \omega) \in (0, +\infty) \times \mathbf{S}^d$, for all $k \in \mathbf{Z}^d$. Since the Fourier transform is one-to-one, this implies that $g = 0$ in $L^\infty(Z \times \mathbf{S}^d)$.

The discussion above shows the existence and uniqueness of the solution $f_\sigma \in L^\infty(Z \times \mathbf{S}^d)$ to the boundary value problem (6) in the case where $\beta = \kappa\sigma$; besides we have seen at the end of Step 3 that

$$(24) \quad \|f_\sigma\|_{L^\infty} \leq \|S\|_{L^\infty}.$$

If moreover $S \in W^{1,\infty}(\mathbf{T}^d)$, we may apply the results of Steps 1-4 to $\partial_{x_j} S$; this shows that

$$f_\sigma \in L^\infty((0, +\infty) \times \mathbf{S}^d; W^{1,\infty}(\mathbf{T}^d)),$$

and satisfies the bound

$$(25) \quad \|\nabla_x f_\sigma\|_{L^\infty} \leq \|\nabla_x S\|_{L^\infty}, \quad \sigma > 0.$$

Notice that this bound is uniform in $\sigma > 0$. In other words, statement (a) in Theorem 2.3 is implied by steps 1-4 with $\beta = \kappa\sigma$.

Proof of Lemma 3.1. The formula for $\langle \bar{\phi} \mathcal{L} \phi \rangle$ is (8) where ϕ is replaced with $\bar{\phi}$ and ψ with ϕ .

On the other hand, we already know from Lemma 2.2 that \mathcal{L} is self-adjoint and of Fredholm type on $L^2(\mathbf{S}^d; \mathbf{C}^d)$. The Fredholm alternative implies that the continuous, one-to-one linear map

$$\mathcal{L}|_{(\text{Ker } \mathcal{L})^\perp} : (\text{Ker } \mathcal{L})^\perp \rightarrow (\text{Ker } \mathcal{L})^\perp$$

is onto. By Banach's open mapping theorem (see Theorem 2.6 and Corollary 2.7 in [6]), this map is bicontinuous, which implies the existence of the positive constant μ . \square

3.5. **Step 5.** Set

$$F_\sigma(x) := \frac{S(x)}{1 + \sigma\nu} + \frac{\kappa\sigma}{1 + \kappa\sigma} \langle\langle f_\sigma \rangle\rangle_-(x, 0);$$

arguing as at the end of step 2, we find that

$$\|F_\sigma\|_{L^\infty} \leq \frac{\|S\|_{L^\infty}}{1 + \kappa\sigma} + \frac{\kappa\sigma}{1 + \kappa\sigma} \lim_{\lambda \rightarrow 0^+} \|\mathcal{A}_{\lambda, \sigma, \kappa\sigma} F_\sigma\|_{L^\infty},$$

and then

$$\|\mathcal{A}_{\lambda, \sigma, \kappa\sigma} F_\sigma\|_{L^\infty} \leq \|\mathcal{A}_{\lambda, \sigma, \kappa\sigma} F_\sigma - \mathcal{A}_{\lambda, \sigma, \kappa\sigma} 0\|_{L^\infty} + \|\mathcal{A}_{\lambda, \sigma, \kappa\sigma} 0\|_{L^\infty} \leq \frac{\kappa\sigma}{1 + \kappa\sigma} \|F_\sigma\|_{L^\infty},$$

while

$$\|\mathcal{A}_{\lambda, \sigma, \kappa\sigma} 0\|_{L^\infty} \leq \frac{\|S\|_{L^\infty}}{1 + \kappa\sigma}.$$

Therefore

$$\|F_\sigma\|_{L^\infty} \leq \frac{\|S\|_{L^\infty}}{1 + \kappa\sigma} + \frac{\kappa\sigma}{1 + \kappa\sigma} \left(\frac{\kappa\sigma}{1 + \kappa\sigma} \|F_\sigma\|_{L^\infty} + \frac{\|S\|_{L^\infty}}{1 + \kappa\sigma} \right)$$

and hence

$$\left(1 - \left(\frac{\kappa\sigma}{1 + \kappa\sigma} \right)^2 \right) \|F_\sigma\|_{L^\infty} \leq \left(1 + \frac{\kappa\sigma}{1 + \kappa\sigma} \right) \frac{\|S\|_{L^\infty}}{1 + \kappa\sigma}.$$

Since

$$\left(1 - \left(\frac{\kappa\sigma}{1 + \kappa\sigma} \right)^2 \right) = \left(1 + \frac{\kappa\sigma}{1 + \kappa\sigma} \right) \left(1 - \frac{\kappa\sigma}{1 + \kappa\sigma} \right) = \left(1 + \frac{\kappa\sigma}{1 + \kappa\sigma} \right) \frac{1}{1 + \kappa\sigma}$$

the inequality above implies that

$$\|F_\sigma\|_{L^\infty} \leq \|S\|_{L^\infty}.$$

By the same token

$$\|\partial_{x_j} F_\sigma\|_{L^\infty} \leq \|\partial_{x_j} S\|_{L^\infty}, \quad j = 1, \dots, d.$$

Let $\chi(y) := (1 - y)_+^2$, so that $\chi \in C^1([0, \infty))$ has its support equal to $[0, 1]$, and set

$$g_\sigma(x, y) := F_\sigma(x) \chi(y);$$

define further the function

$$h_\sigma(x, y, \omega) := f_\sigma(x, y, \omega) - g_\sigma(x, y).$$

It is easily seen that h_σ is a solution to the boundary value problem

$$(26) \quad \begin{cases} \omega \cdot \nabla_z h_\sigma + \sigma \mathcal{L} h_\sigma = -\omega \cdot \nabla g_\sigma(z), \\ h_\sigma(x, 0, \omega) = 0, \quad \omega_y > 0. \end{cases}$$

Notice that $h_\sigma \in L^\infty((0, +\infty) \times \mathbf{S}^d; W^{1, \infty}(\mathbf{T}^d))$.

Because of the regularity of h_σ in x , we deduce from the equation that the function $z \mapsto h_\sigma(z, \omega)$ belongs to $W^{1, \infty}(\mathbf{T}^d \times (0, +\infty))$. Thus, we can multiply both sides of the first equation in (26) by $2\sigma h_\sigma$ and integrate in ω , to obtain

$$\sigma \operatorname{div}_z \langle \omega h_\sigma^2 \rangle + 2\sigma^2 \langle h_\sigma \mathcal{L} h_\sigma \rangle = -2\sigma \langle \omega h_\sigma \rangle \cdot \nabla g_\sigma(z) = -2\sigma \langle \omega (h_\sigma - \langle h_\sigma \rangle) \rangle \cdot \nabla g_\sigma(z).$$

After integrating in $x \in \mathbf{T}^d$ both sides of this equality, we see that

$$\begin{aligned} \sigma \partial_y \int_{\mathbf{T}^d} \langle \omega_y h_\sigma^2 \rangle(x, y) dx + 2\sigma^2 \int_{\mathbf{T}^d} \langle h_\sigma \mathcal{L} h_\sigma \rangle(x, y) dx \\ = -2\sigma \int_{\mathbf{T}^d} \langle \omega(h_\sigma - \langle h_\sigma \rangle) \rangle \cdot \nabla g_\sigma(x, y) dx. \end{aligned}$$

By construction

$$f_\sigma(x, y, \omega) = h_\sigma(x, y, \omega) \quad \text{whenever } y > 1,$$

so that

$$\langle \omega_y h_\sigma^2 \rangle(x, y) = \langle \omega_y f_\sigma^2 \rangle(x, y) \text{ and } g_\sigma(x, y) = 0, \quad \text{for all } y > 1.$$

On the other hand, Parseval's theorem implies that

$$\int_{\mathbf{T}^d} \langle \omega_y f_\sigma^2 \rangle(x, y) dx = \left\langle \omega_y \int_{\mathbf{T}^d} f_\sigma^2(x, y, \cdot) dx \right\rangle = \sum_{k \in \mathbf{Z}^d} \langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y).$$

Proceeding as in step 4 and applying Lemma 3.1, we see that

$$\partial_y \langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y) + 2\sigma\mu \left\langle \left| \hat{f}_\sigma - \langle \hat{f}_\sigma \rangle \right|^2 \right\rangle(k, y) \leq 0, \quad y > 1, \quad k \in \mathbf{Z}^d.$$

For all $k \in \mathbf{Z}^d$, the function

$$y \mapsto \langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y)$$

is bounded and Lipschitz continuous on $(0, +\infty)$, and nonincreasing on $(1, +\infty)$. Hence

$$\int_1^\infty \left\langle \left| \hat{f}_\sigma - \langle \hat{f}_\sigma \rangle \right|^2 \right\rangle(k, y) dy \leq \frac{1}{2\sigma\mu} \left[\langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y) \right]_{y \rightarrow \infty}^{y=1} < \infty.$$

With the decomposition

$$\langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y) = \langle \omega_y \hat{f}_\sigma - \langle \hat{f}_\sigma \rangle \rangle^2(k, y) + 2\Re \left(\langle \omega_y (\hat{f}_\sigma - \langle \hat{f}_\sigma \rangle) \rangle(k, y) \overline{\langle \hat{f}_\sigma \rangle(k, y)} \right),$$

we conclude that

$$y \mapsto \langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y) \text{ belongs to } L^2(0, +\infty)$$

for all $k \in \mathbf{Z}^d$. (Notice that we do not seek a uniform in σ estimate in L^2 for $\hat{f}_\sigma - \langle \hat{f}_\sigma \rangle$ at this stage in the argument, although this is our ultimate goal.) Since this function is nonincreasing on $(1, +\infty)$ for all $k \in \mathbf{Z}^d$, we find that

$$0 = \lim_{Y \rightarrow +\infty} \langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, Y) \leq \langle \omega_y |\hat{f}_\sigma|^2 \rangle(k, y) \quad \text{for all } y > 1 \text{ and all } k \in \mathbf{Z}^d.$$

Thus, for each $Y > 1$, one has

$$\begin{aligned} \sigma \int_{\mathbf{T}^d} \langle \omega_y f_\sigma^2 \rangle(x, Y) dx + 2\sigma^2 \mu \int_0^Y \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) dx dy \\ = -2\sigma \int_0^1 \int_{\mathbf{T}^d} \langle \omega(h_\sigma - \langle h_\sigma \rangle) \rangle \cdot \nabla g_\sigma(x, y) dx dy + \sigma \int_{\mathbf{T}^d} \langle \omega_y h_\sigma^2 \rangle(x, 0) dx, \end{aligned}$$

so that

$$(27) \quad \begin{aligned} & 2\sigma^2\mu \int_0^Y \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) \, dx \, dy \\ & \leq -2\sigma \int_0^1 \int_{\mathbf{T}^d} \langle \omega(h_\sigma - \langle h_\sigma \rangle) \rangle \cdot \nabla g_\sigma(x, y) \, dx \, dy. \end{aligned}$$

Indeed

$$\langle \omega_y h_\sigma^2 \rangle(x, 0) = -\frac{|\mathbf{B}^d|}{|\mathbf{S}^d|} \langle h_\sigma^2 \rangle_-(x, 0) \leq 0,$$

since $h_\sigma(x, 0, \omega) = 0$ for $\omega_y > 0$ by construction, and on the other hand, as explained above,

$$\int_{\mathbf{T}^d} \langle \omega_y f_\sigma^2 \rangle(x, Y) \, dx \geq 0 \quad \text{for all } Y > 1.$$

Notice that the first integral on the right hand side of (27) involves only $y \in [0, 1]$ since $g(x, y) = 0$ for all $y > 1$ by construction.

By the Cauchy-Schwarz inequality, for all $Y > 1$, one has

$$\begin{aligned} & \sigma^2\mu \int_0^Y \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) \, dx \, dy \\ & \leq \left(\sigma^2 \int_0^1 \int_{\mathbf{T}^d} |\langle \omega(h_\sigma - \langle h_\sigma \rangle) \rangle|^2(x, y) \, dx \, dy \right)^{1/2} \left(\int_0^1 \int_{\mathbf{T}^d} |\nabla g_\sigma(x, y)|^2 \, dx \, dy \right)^{1/2} \\ & \leq \left(\sigma^2 \int_0^Y \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) \, dx \, dy \right)^{1/2} \left(\int_0^1 \int_{\mathbf{T}^d} |\nabla g_\sigma(x, y)|^2 \, dx \, dy \right)^{1/2}, \end{aligned}$$

so that

$$\sigma^2 \int_0^Y \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) \, dx \, dy \leq \int_0^1 \int_{\mathbf{T}^d} |\nabla g_\sigma(x, y)|^2 \, dx \, dy.$$

Observe that

$$h_\sigma - \langle h_\sigma \rangle = f_\sigma - \langle f_\sigma \rangle$$

since $g_\sigma = f_\sigma - h_\sigma$ is independent of ω by construction. Hence

$$\sigma^2\mu \int_0^Y \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) \, dx \, dy \leq \int_0^1 \int_{\mathbf{T}^d} |\nabla g_\sigma(x, y)|^2 \, dx \, dy,$$

and since this inequality holds for all $Y > 0$, we conclude that

$$\sigma^2\mu \int_0^\infty \int_{\mathbf{T}^d} \left\langle (h_\sigma - \langle h_\sigma \rangle)^2 \right\rangle(x, y) \, dx \, dy \leq \int_0^1 \int_{\mathbf{T}^d} |\nabla g_\sigma(x, y)|^2 \, dx \, dy.$$

3.6. Step 6. Summarizing, for each $S \in W^{1,\infty}(\mathbf{T}^d)$, the boundary value problem (6) has a unique (by Step 4) solution $f_\sigma \in L^\infty(Z \times \mathbf{S}^d)$, and this solution satisfies

$$\|f_\sigma\|_{L^\infty} \leq \|S\|_{L^\infty},$$

according to Step 3, and

$$\|\sigma(f_\sigma - \langle f_\sigma \rangle)\|_{L^2(Z \times \mathbf{S}^d)}^2 \leq \frac{2}{\mu} |\mathbf{S}^d| \left(\|\nabla S\|_{L^\infty(\mathbf{T}^d)}^2 + \frac{4}{3} \|S\|_{L^\infty(\mathbf{T}^d)}^2 \right),$$

by Step 5.

Therefore, there exists a sequence $\sigma_n \rightarrow +\infty$ such that

$$f_{\sigma_n} \rightarrow f \quad \text{in } L^\infty(Z \times \mathbf{S}^d) \text{ weak } -*,$$

and

$$J_n := \sigma_n(f_{\sigma_n} - \langle f_{\sigma_n} \rangle) \rightarrow J \quad \text{in } L^2(Z \times \mathbf{S}^d) \text{ weak.}$$

In particular, the fact that J_n is bounded in $L^2(Z \times \mathbf{S}^d)$ implies that

$$(28) \quad f(z, \omega) = \rho(z) \text{ for a.e. } (z, \omega) \in Z \times \mathbf{S}^d.$$

In order to compute J , observe that, since $\mathcal{L}1 = 0$,

$$\mathcal{L}J_n = \sigma_n \mathcal{L}f_{\sigma_n} = -\omega \cdot \nabla_z f_{\sigma_n}.$$

Since the linear operator \mathcal{L} is continuous on $L^2(\mathbf{S}^d)$, one has

$$\mathcal{L}J_n \rightarrow \mathcal{L}J \text{ in } L^2(Z \times \mathbf{S}^d) \text{ weak,}$$

and since

$$\omega \cdot \nabla_z f_{\sigma_n} \rightarrow \omega \cdot \nabla \rho \quad \text{in } \mathcal{D}'(Z \times \mathbf{S}^d),$$

we conclude that

$$\mathcal{L}J(z, \omega) = \omega \cdot \nabla \rho(z) \text{ for a.e. } (z, \omega) \in Z \times \mathbf{S}^d.$$

Beside, since $J_n(z, \cdot) \in (\text{Ker } \mathcal{L})^\perp$ for all n and a.e. $z \in Z$, one has also

$$J(z, \cdot) \in (\text{Ker } \mathcal{L})^\perp, \quad \text{for a.e. } z \in Z.$$

Applying the Fredholm alternative to \mathcal{L} , we conclude that

$$(29) \quad J(z, \omega) = -\Omega(\omega) \cdot \nabla \rho(z), \quad \text{for a.e. } (z, \omega) \in Z \times \mathbf{S}^d.$$

Notice that the formula above for J and the L^2 bound on J_n imply that

$$\nabla \rho \in L^2(Z).$$

Let now $\phi \in C_c^\infty(\mathbf{T}^d \times (0, +\infty))$. Multiplying both sides of the equation for f_{σ_n} in (6) by

$$\sigma_n \phi(z) + \Omega(\omega) \cdot \nabla \phi(z)$$

and integrating both sides in (z, ω) , we obtain

$$(30) \quad \int_0^\infty \int_{\mathbf{T}^d} (\phi(z) \text{div}_z(\sigma_n \langle \omega f_{\sigma_n} \rangle) + \nabla \phi(z) \cdot \langle \Omega(\omega) \cdot \nabla_z f_{\sigma_n} \rangle) dx dy \\ + \int_0^\infty \int_{\mathbf{T}^d} \sigma_n^2 \phi(z) \langle \mathcal{L}f_{\sigma_n} \rangle dx dy + \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle \sigma_n \Omega \mathcal{L}f_{\sigma_n} \rangle dx dy = 0.$$

Since \mathcal{L} is self-adjoint on $L^2(\mathbf{S}^d)$ (see Lemma 2.2), one has

$$\langle \mathcal{L}f_{\sigma_n} \rangle = \langle f_{\sigma_n} \mathcal{L}^* 1 \rangle = 0,$$

and

$$\langle \sigma_n \Omega \mathcal{L}f_{\sigma_n} \rangle = \langle \sigma_n f_{\sigma_n} \mathcal{L}\Omega \rangle = \langle \sigma_n \omega f_{\sigma_n} \rangle.$$

Hence the third term on the left hand side of (30) is 0, while the first and last terms on the left hand side of (30) combine to give

$$\int_0^\infty \int_{\mathbf{T}^d} (\text{div}_z(\sigma_n \langle \omega f_{\sigma_n} \rangle \phi(z)) + \nabla \phi(z) \cdot \langle \Omega(\omega) \cdot \nabla_z f_{\sigma_n} \rangle) dx dy = 0.$$

The first integral on the left hand side is simplified by using Green's formula:

$$\int_0^\infty \int_{\mathbf{T}^d} \text{div}_z(\sigma_n \langle \omega f_{\sigma_n} \rangle \phi(z)) dx dy = - \int_{\mathbf{T}^d} \sigma_n \langle \omega f_{\sigma_n} \rangle \phi(x, 0) dx.$$

Next we use the boundary condition verified by f_{σ_n} , i.e.

$$f_{\sigma_n}(x, 0, \omega) = \frac{S(x)}{1 + \kappa\sigma_n} + \frac{\kappa\sigma_n}{1 + \kappa\sigma_n} \langle\langle f_{\sigma_n} \rangle\rangle_-(x), \quad \omega_y > 0,$$

or equivalently

$$f_{\sigma_n}(x, 0, \omega) = S(x) - \kappa\sigma_n \frac{|\mathbf{S}^d|}{|\mathbf{B}^d|} \langle \omega f_{\sigma_n} \rangle(x, 0), \quad \omega_y > 0.$$

Since the right hand side of this identity is independent of ω , one has also

$$(31) \quad \langle\langle f_{\sigma_n} \rangle\rangle_+(x, 0) = S(x) - \kappa\sigma_n \frac{|\mathbf{S}^d|}{|\mathbf{B}^d|} \langle \omega f_{\sigma_n} \rangle(x, 0), \quad x \in \mathbf{T}^d.$$

Thus

$$\begin{aligned} \int_0^\infty \int_{\mathbf{T}^d} \operatorname{div}_z(\sigma_n \langle \omega f_{\sigma_n} \rangle \phi(z)) \, dx \, dy &= - \int_{\mathbf{T}^d} \sigma_n \langle \omega f_{\sigma_n} \rangle \phi(x, 0) \, dx \\ &= \frac{|\mathbf{B}^d|}{\kappa|\mathbf{S}^d|} \int_{\mathbf{T}^d} (S(x) - \langle\langle f_{\sigma_n} \rangle\rangle_+(x, 0)) \phi(x, 0) \, dx. \end{aligned}$$

In other words

$$\begin{aligned} &\frac{|\mathbf{B}^d|}{\kappa|\mathbf{S}^d|} \int_{\mathbf{T}^d} (S(x) - \langle\langle f_{\sigma_n} \rangle\rangle_+(x, 0)) \phi(x, 0) \, dx \\ &= \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle \Omega(\omega \cdot \nabla_z f_{\sigma_n}) \rangle \, dx \, dy \end{aligned}$$

for all $\phi \in C_c^\infty(\mathbf{T}^d \times (0, +\infty))$.

As $n \rightarrow \infty$, the integral on the right hand side satisfies

$$\begin{aligned} &\int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle \Omega(\omega \cdot \nabla_z f_{\sigma_n}) \rangle \, dx \, dy = - \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle \Omega \mathcal{L} J_n \rangle \, dx \, dy \\ &\rightarrow - \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle \Omega \mathcal{L} J \rangle \, dx \, dy = - \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle (\mathcal{L}^* \Omega) J \rangle \, dx \, dy \\ &= - \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \langle \omega J \rangle \, dx \, dy = \int_0^\infty \int_{\mathbf{T}^d} \langle \omega \otimes \Omega \rangle : \nabla \phi(z) \otimes \nabla_z \rho(z) \, dx \, dy \\ &= \frac{\langle \omega \cdot \Omega \rangle}{d+1} \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \nabla_z \rho(z) \, dx \, dy. \end{aligned}$$

On the other hand, arguing as in Step 3 shows that

$$\begin{aligned} &\frac{|\mathbf{B}^d|}{|\mathbf{S}^d|} \partial_y \int_{\mathbf{T}^d} \langle\langle f_{\sigma_n} \rangle\rangle_+(x, y) \psi(x) \, dx = \partial_y \int_{\mathbf{T}^d} \langle \omega_y^+ f_{\sigma_n} \rangle(x, y) \psi(x) \, dx \\ &= \int_{\mathbf{T}^d} \langle \mathbf{1}_{\omega_y > 0} \omega_x f_{\sigma_n} \rangle(x, y) \psi'(x) \, dx - \int_{\mathbf{T}^d} \langle \mathbf{1}_{\omega_y > 0} \mathcal{L} J_n \rangle(x, y) \psi(x) \, dx \\ &= O(1) \text{ in } L^2 + L^\infty(0, +\infty). \end{aligned}$$

By Ascoli-Arzelà theorem, we conclude that

$$\int_{\mathbf{T}^d} \langle\langle f_{\sigma_n} \rangle\rangle_+(x, 0) \phi(x, 0) \, dx \rightarrow \int_{\mathbf{T}^d} \rho(x, 0) \phi(x, 0) \, dx.$$

Summarizing, we have proved that

$$\begin{aligned} &\rho \in L^\infty(Z), \quad \nabla \rho \in L^2(Z) \\ &\frac{|\mathbf{B}^d|}{\kappa|\mathbf{S}^d|} \int_{\mathbf{T}^d} (S(x) - \rho(x, 0)) \phi(x, 0) \, dx = \frac{\langle \omega \cdot \Omega \rangle}{d+1} \int_0^\infty \int_{\mathbf{T}^d} \nabla \phi(z) \cdot \nabla_z \rho(z) \, dx \, dy \end{aligned}$$

for all $\phi \in C_c^\infty(\mathbf{T}^d \times (0, +\infty))$. One easily checks that this is the variational formulation of the boundary value problem (17).

This proves statement (b) of Theorem 2.3.

3.7. Step 7. Set $Y = \mathbf{T}^d \times (0, 1)$. The results obtained in Step 6 imply that

$$f_{\sigma_n}|_{Y \times \mathbf{S}^d} \rightarrow \rho \quad \text{and} \quad \omega \cdot \nabla_z f_{\sigma_n}|_{Y \times \mathbf{S}^d} \rightarrow \omega \cdot \nabla_z \rho$$

weakly in $L^2(Y \times \mathbf{S}^d)$.

By applying Cessenat's trace theorem (Theorem 1 in [9]), we find that

$$(32) \quad f_{\sigma_n}(x, 0, \omega) \rightarrow R(x) \text{ weakly in } L^2(\mathbf{T}^d \times \mathbf{S}^d; \omega_y^+ d\omega dx),$$

where

$$R = \rho|_{y=0}.$$

This last point requires additional explanations. At this point, we use Cessenat's notation in [9]. Observe that, for all $x \in \mathbf{T}^d$, the exit time from Y , starting from y in the direction ω , is

$$\tau_{(x,0),\omega} = 1/|\omega_y| \geq 1, \quad \text{since } |\omega| = 1.$$

Choosing the arbitrary parameter $K = 1$ in the definition of the measure $d\xi$ on p. 832 in [9], viz.

$$d\xi|_{\mathbf{T}^d \times \{0\} \times \mathbf{S}^d} = |\omega_y| \min(1, \tau_{(x,0),\omega}) dx d\omega,$$

has its restriction to $\mathbf{T}^d \times \{0\}$ given by

$$d\xi|_{\mathbf{T}^d \times \{0\} \times \mathbf{S}^d} = |\omega_y| dx d\omega.$$

This observation justifies that the convergence holds as stated in (32).

Since

$$\rho \in L^\infty(0, +\infty; L^2(\mathbf{T}^d)) \quad \text{and} \quad \nabla \rho \in L^2(Z),$$

one has in particular $\rho|_Y \in H^1(Y)$, so that

$$R = \rho|_{y=0} \in H^{1/2}(\mathbf{T}^d).$$

Since

$$\nabla \rho \in L^2(Z) \quad \text{and} \quad \operatorname{div}(\nabla \rho) = 0,$$

we deduce from the trace theorem of J.-L. Lions for the space $H(Y, \operatorname{div})$ of vector fields in $L^2(Y; \mathbf{R}^{d+1})$ with divergence in $L^2(Y)$ (see Lemma 20.2 in [22]) that

$$\partial_y \rho|_{y=0} \in H^{-1/2}(\mathbf{T}^d).$$

On the other hand

$$\sigma_n \langle \omega f_{\sigma_n} \rangle|_Y = \sigma \langle \omega J_n \rangle|_Y \text{ is bounded in } L^\infty(0, 1; L^2(\mathbf{T}^d)) \subset L^2(Y),$$

and

$$\operatorname{div}_z(\sigma_n \langle \omega f_{\sigma_n} \rangle) = 0.$$

Applying again the Lions trace theorem in $H(Y, \operatorname{div})$, we conclude that

$$\sigma_n \langle \omega_y f_{\sigma_n} \rangle|_{y=0} \rightarrow -\frac{\langle \omega \cdot \Omega \rangle}{d+1} \partial_y \rho|_{y=0} \text{ in } H^{-1/2}(\mathbf{T}^d) \text{ weak.}$$

Since $\nabla \rho \in L^2(Z)$, we find that

$$\partial_y \rho \in L^2(\mathbf{T}^d \times (0, \infty)),$$

so that

$$\begin{aligned} \int_{\mathbf{T}^d} |\rho(x, y_2) - \rho(x, y_1)|^2 dx &= \int_{\mathbf{T}^d} \left(\int_{y_1}^{y_2} \partial_y \rho(x, z) dz \right)^2 dx \\ &\leq |y_2 - y_1| \int_{\mathbf{T}^d} \int_{y_1}^{y_2} |\partial_y \rho(x, z)|^2 dz dx \leq |y_2 - y_1| \|\nabla \rho\|_{L^2(Z)}. \end{aligned}$$

Hence

$$\rho \in C_b(0, +\infty; L^2(\mathbf{T}^d)).$$

Since ρ is a harmonic extension of $R = \rho|_{y=0}$, we deduce from Kwasnicki's Theorem 1.1 (j) that

$$\partial_y \rho|_{y=0} = -(-\Delta_x)^{1/2} R.$$

This concludes the proof of statement (c) of Theorem 2.3.

4. OPEN PROBLEMS

The result obtained in the present paper suggests various questions, still open at the time of this writing.

(a) Can one extend Theorem 2.3 to other kinetic models — for instance, to the linearized Boltzmann equation, or even to the linearized BGK model? For instance, one could consider the Boltzmann equation for a vapor in a half-space over its liquid, condensed phase, with a linear combination of diffuse reflection and condensation or evaporation at the boundary, assuming that the Knudsen number in the vapor is small, and that the accommodation coefficient is close to one. At present, the proof of Theorem 2.3 is based on the maximum principle, which the linearized Boltzmann equation does not satisfy. Before treating the case of the linearized Boltzmann equation, it would be necessary to have a proof of Theorem 2.3 based on an L^2 energy estimate.

(b) Can one obtain in this way powers of the Laplacian other than $(-\Delta)^{1/2}$? The harmonic extension result recalled in section 1.1 suggests that one should seek a linearized Boltzmann equation leading to the diffusion operator

$$y^{\frac{2}{\gamma}-2} \Delta_x + \gamma^2 c_\gamma^{1/2} \partial_y^2.$$

The scattering operator of such a linearized Boltzmann equation must be strongly anisotropic; besides, the scattering coefficient should vanish either as $y \rightarrow +\infty$ or as $y \rightarrow 0$ depending on whether $\gamma \in (0, 1)$ or $\gamma \in (1, 2)$.

(c) Can one extend the result in Theorem 2.3 to other domains than half-spaces? Assuming that one can derive from the linearized Boltzmann equation in a bounded (spatial) domain Ω with smooth boundary the diffusion problem

$$\begin{cases} \Delta \rho(x) = 0, & x \in \Omega, \\ \left(\rho + \kappa \frac{\partial \rho}{\partial n} \right) \Big|_{\partial \Omega} = S, \end{cases}$$

the trace $\rho|_{\partial \Omega}$ must satisfy the equation

$$\rho|_{\partial \Omega} + \kappa \Lambda \rho|_{\partial \Omega} = S,$$

where Λ is the Dirichlet-to-Neumann operator, defined as follows.

For $u \in H^{1/2}(\partial\Omega)$, we set

$$\Lambda u := \frac{\partial U}{\partial n} \Big|_{\partial\Omega}$$

where $U \in H^1(\Omega)$ is the unique solution of

$$-\Delta U = 0 \text{ in } \Omega, \quad U|_{\partial\Omega} = u.$$

We recall that the linear map $u \mapsto U$ is continuous from $H^{1/2}(\partial\Omega)$ to $H^1(\Omega)$. By Green's formula

$$\left\langle \frac{\partial U}{\partial n}, \phi \right\rangle := \int_{\Omega} \nabla U(x) \cdot \nabla \phi(x) \, dx,$$

so that, for all $\phi \in H^1(\Omega)$

$$\left| \left\langle \frac{\partial U}{\partial n}, \phi \right\rangle \right| \leq \|\nabla U\|_{L^2(\Omega)} \|\nabla \phi\|_{L^2(\Omega)} \leq C \|u\|_{H^{1/2}(\partial\Omega)} \|\nabla \phi\|_{L^2(\Omega)}.$$

This defines $\partial U / \partial n|_{\partial\Omega}$ as an element of $H^{-1/2}(\partial\Omega)$, and shows that the linear map Λ is continuous from $H^{1/2}(\partial\Omega)$ to $H^{-1/2}(\partial\Omega)$. Observe that Λ is self-adjoint since

$$\langle \mathcal{L}u, \phi \rangle = \int_{\Omega} \nabla u(x) \cdot \nabla \phi(x) \, dx = \langle \mathcal{L}\phi, u \rangle.$$

Besides, $\Lambda \geq 0$ and $\text{Ker } \mathcal{L} = \mathbf{R} = \{ \text{constants} \}$, since

$$\langle \mathcal{L}u, u \rangle = \|\nabla u\|_{L^2(\Omega)}^2.$$

Call $\Delta^{\partial\Omega}$ the Laplace-Beltrami operator⁴ for the restriction of the Euclidean metric to $\partial\Omega$. Both Λ and $(-\Delta^{\partial\Omega})^{1/2}$ are unbounded self-adjoint operators on $L^2(\partial\Omega)$, mapping $H^{1/2}(\partial\Omega)$ to $H^{-1/2}(\partial\Omega)$.

However, Λ and $(-\Delta^{\partial\Omega})^{1/2}$ are different in general, as shown by the following example.

Example. Let Ω be the open unit ball of \mathbf{R}^{d+1} , so that $\partial\Omega = \mathbf{S}^d$ is the d -dimensional unit sphere. Using spherical coordinates, i.e. setting $r = |z|$ and $\omega := z/|z|$ for $z \neq 0$, one has

$$\Delta_z f(z) = r^{-d} \partial_r (r^d \partial_r f(r\omega)) + r^{-2} \Delta_{\omega}^{\mathbf{S}^d} f(r\omega).$$

Denote by $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \dots$ the sequence of eigenvalues of $\Delta^{\mathbf{S}^d}$ counted with multiplicities, and let $(e_n)_{n \geq 0}$ be a orthonormal and complete system in $L^2(\partial\Omega)$, with $-\Delta^{\mathbf{S}^d} e_n = \lambda_n e_n$. For each $f \equiv f(r, \omega) \in L^2(\partial\Omega)$, set

$$f_n(r) := (e_n | f(r, \cdot))_{L^2(\partial\Omega)}, \quad n \in \mathbf{N}, \, r > 0.$$

Then $\Delta_z f = 0$ on Ω if and only if

$$r^{2-d} \partial_r (r^d \partial_r f_n(r)) - \lambda_n f_n(r) = 0, \quad n \in \mathbf{N}, \, r > 0,$$

or equivalently

$$r^2 f_n''(r) + dr f_n'(r) - \lambda_n f_n(r) = 0, \quad n \in \mathbf{N}, \, r > 0.$$

This differential equation has a two-dimensional space of solutions over the half-line $(0, +\infty)$, viz.

$$\{ C_+ r^{\alpha_+(n)} + C_- r^{\alpha_-(n)}, \, C_{\pm} \in \mathbf{R} \}$$

⁴At variance with the definition commonly used in geometry, we normalize this operator so that $\Delta^{\partial\Omega} \leq 0$.

where

$$\alpha_{\pm}(n) := -\frac{d-1}{2} \pm \sqrt{\frac{(d-1)^2}{4} + \lambda_n}, \quad n \geq 0.$$

Observe that $\alpha_- \leq 1-d$ for all $n \in \mathbf{N}$, since $\lambda_n \geq 0$. Hence $2(\alpha_- - 1) + d \leq -d \leq 1$, so that

$$\int_0^1 |r^{\alpha_- - 1}|^2 r^d dr = +\infty.$$

Hence all harmonic functions in $H^1(\Omega)$ in the unit ball Ω are of the form

$$f(r\omega) = \sum_{n \in \mathbf{N}} c_n r^{\alpha_+(n)} e_n(\omega),$$

and

$$\partial_r f(r\omega)|_{r=1} = \sum_{n \in \mathbf{N}} c_n \alpha_-(n) e_n(\omega) \quad \text{while} \quad f(\omega) = \sum_{n \in \mathbf{N}} c_n e_n(\omega).$$

The formula for $\alpha_+(n)$ indicates that, in this case

$$\Lambda = -\frac{d-1}{2} + \sqrt{\frac{(d-1)^2}{4} - \Delta \mathbf{S}^d}.$$

In particular

$$\Lambda \neq (-\Delta \mathbf{S}^d)^{1/2}, \quad \text{unless } d = 1.$$

REFERENCES

- [1] P. Aceves-Sánchez, C. Schmeiser: *Fractional diffusion limit of a linear kinetic equation in a bounded domain*, arXiv: 1607.00855 [math.AP]
- [2] C. Bardos, F. Golse, B. Perthame, R. Sentis: *The nonaccretive radiative transfer equations, existence of solutions and Rosseland approximation*, J. Funct. Anal. **77** (1988), 434–460.
- [3] C. Bardos, E. Bernard, F. Golse, R. Sentis: *The Diffusion Approximation for the Linear Boltzmann Equation with Vanishing Scattering Coefficient*, Commun. Math. Sci. **13** (2015), 641–671.
- [4] N. Ben Abdallah, A. Mellet, M. Puel: *Anomalous diffusion limit for kinetic equations with degenerate collision frequency*, Math. Models and Methods Appl. Sci., **21** (2011), 2249–2262.
- [5] A. Bensoussan, J.-L. Lions, G. C. Papanicolaou: *Boundary layers and homogenization of transport processes*, Publ. Res. Inst. Math. Sci. **15** (1979), 53–157.
- [6] H. Brezis: “Functional Analysis, Sobolev Spaces and Partial Differential Equations”, Springer New-York, Dordrecht, Heidelberg, London, 2011.
- [7] C. Bardos, R. Santos, R. Sentis: *Diffusion approximation and computation of the critical size*, Trans. Amer. Math. Soc. **284** (1984), 617–649.
- [8] S. Chandrasekhar: “Radiative Transfer”, Oxford, Clarendon Press, 1950.
- [9] M. Cessenat: *Théorèmes de trace L^p pour des espaces de fonctions de la neutronique*, C.R. Acad. Sci. Paris Sér. I, **299** (1984), 831–834.
- [10] R. Dautray, J.-L. Lions: “Analyse mathématique et calcul numérique pour les sciences et les techniques”, Masson, Paris, 1985.
- [11] L. Desvillettes, F. Golse: *A remark concerning the Chapman-Enskog asymptotics*, In ‘Advances in kinetic theory and computing’, B; Perthame ed., 191–203, Ser. Adv. Math. Appl. Sci., **22**, World Sci. Publ., River Edge, NJ, 1994.
- [12] U. Frisch, H. Frisch: *Non LTE Transfer. Asymptotic Expansion for Small ϵ* , Mon. Not. R. Astr. Not. **181** (1977), 273–280.
- [13] F. Golse: *Fluid Dynamic Limits of the Kinetic Theory of Gases*, in “From Particle Systems to Partial Differential Equations”, C. Bernardin and Patrícia Gonçalves eds., 3–91, Springer Proc. in Math. and Statist. **75**, Springer Verlag, Berlin, Heidelberg, 2014.
- [14] D. Hilbert: *Begründung der kinetischen Gastheorie*, Math. Ann. **72** (1912), 562–577.
- [15] A. M. Il’in, R. Z. Has’minskii (Khasminskii): *On the equations of Brownian motion* (Russian), Teor. Veroyatnost. i Primenen. **9** (1964), 466–491.
- [16] M. Kwasnicki: *Ten Equivalent Definitions of the Fractional Laplace Operator*, Fract. Calc. Appl. Anal. **20** (2017), 7–51.

- [17] E. W. Larsen, J. B. Keller: *Asymptotics solutions of neutron transport problems for small mean free paths*, J. Math. Phys. **15** (1974), 75–81.
- [18] A. Mellet: *Fractional diffusion limit for collisional kinetic equations: a moments method*, Indiana Univ. Math. J., **59** (2010), 1333–1360.
- [19] A. Mellet, S. Mischler, and C. Mouhot: *Fractional diffusion limit for collisional kinetic equations*, Arch. Ration. Mech. Anal., **199** (2011), 493–525.
- [20] G. C. Papanicolaou: *Asymptotic analysis of transport processes*, Bull. Amer. Math. Soc. **81** (1975), 330–392.
- [21] G. C. Pomraning: “The Equations of Radiation Hydrodynamics”, Pergamon Press, 1973.
- [22] L. Tartar: “An Introduction to Sobolev Spaces and Interpolation Spaces”; Springer-Verlag, Berlin, Heidelberg, 2007.
- [23] A. Weinberg, E. Wigner: “The Physical Theory of Neutron Chain Reactors”, The University of Chicago Press, 1958.

(C.B.) LABORATOIRE J.-L. LIONS, BP 187, 75252 PARIS CEDEX 05, FRANCE
E-mail address: `claude.bardos@gmail.com`

(F.G.) CMLS, ÉCOLE POLYTECHNIQUE, CNRS, UNIVERSITÉ PARIS-SACLAY , 91128 PALAISEAU
CEDEX, FRANCE
E-mail address: `francois.golse@polytechnique.edu`

(I.M.) LABORATOIRE J.-L. LIONS, BP 187, 75252 PARIS CEDEX 05, FRANCE
E-mail address: `ivan.moyano@math.cnrs.fr`